
Chyung-Ru Wang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7102114/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Mitochondria Are Required for Antigen-Specific T Cell Activation through Reactive Oxygen Species Signaling. Immunity, 2013, 38, 225-236.	14.3	981
2	Tracking the Response of Natural Killer T Cells to a Glycolipid Antigen Using Cd1d Tetramers. Journal of Experimental Medicine, 2000, 192, 741-754.	8.5	818
3	Impaired NK1+ T Cell Development and Early IL-4 Production in CD1-Deficient Mice. Immunity, 1997, 6, 459-467.	14.3	440
4	Glycolipid antigen induces long-term natural killer T cell anergy in mice. Journal of Clinical Investigation, 2005, 115, 2572-2583.	8.2	386
5	Susceptibility of Mice Deficient in CD1D or TAP1 to Infection with Mycobacterium tuberculosis. Journal of Experimental Medicine, 1999, 189, 1973-1980.	8.5	329
6	Non-classical Immunity Controls Microbiota Impact on Skin Immunity and Tissue Repair. Cell, 2018, 172, 784-796.e18.	28.9	323
7	The response of natural killer T cells to glycolipid antigens is characterized by surface receptor down-modulation and expansion. Proceedings of the National Academy of Sciences of the United States of America, 2003, 100, 10913-10918.	7.1	306
8	Cd1-Restricted Nk T Cells Protect Nonobese Diabetic Mice from Developing Diabetes. Journal of Experimental Medicine, 2001, 194, 313-320.	8.5	251
9	Nonclassical binding of formylated peptide in crystal structure of the MHC class lb molecule H2-M3. Cell, 1995, 82, 655-664.	28.9	151
10	Quantitative and Qualitative Differences in the In Vivo Response of NKT Cells to Distinct α- and β-Anomeric Glycolipids. Journal of Immunology, 2004, 173, 3693-3706.	0.8	136
11	H2-M3, A FULL-SERVICE CLASS IbHISTOCOMPATIBILITY ANTIGEN. Annual Review of Immunology, 1997, 15, 851-879.	21.8	125
12	CD1d-expressing Dendritic Cells but Not Thymic Epithelial Cells Can Mediate Negative Selection of NKT Cells. Journal of Experimental Medicine, 2003, 197, 907-918.	8.5	122
13	H-2M3 encodes the MHC Class I molecule presenting the maternally transmitted antigen of the mouse. Cell, 1991, 66, 335-345.	28.9	117
14	Selective inhibition of anthrax edema factor by adefovir, a drug for chronic hepatitis B virus infection. Proceedings of the National Academy of Sciences of the United States of America, 2004, 101, 3242-3247.	7.1	109
15	Serine Protease Inhibitor 6 Protects Cytotoxic T Cells from Self-Inflicted Injury by Ensuring the Integrity of Cytotoxic Granules. Immunity, 2006, 24, 451-461.	14.3	107
16	CD1-restricted adaptive immune responses to <i>Mycobacteria</i> in human group 1 CD1 transgenic mice. Journal of Experimental Medicine, 2009, 206, 2497-2509.	8.5	105
17	Selection and Expansion of CD8α/α1 T Cell Receptor α/β1 Intestinal Intraepithelial Lymphocytes in the Absence of Both Classical Major Histocompatibility Complex Class I and Nonclassical Cd1 Molecules. Journal of Experimental Medicine, 1999, 190, 885-890.	8.5	92
18	Cutting Edge: The Ets1 Transcription Factor Is Required for the Development of NK T Cells in Mice. Journal of Immunology, 2000, 164, 2857-2860.	0.8	86

#	Article	IF	CITATIONS
19	CD1d-Expressing Breast Cancer Cells Modulate NKT Cell-Mediated Antitumor Immunity in a Murine Model of Breast Cancer Metastasis. PLoS ONE, 2011, 6, e20702.	2.5	85
20	Prevention of Allograft Tolerance by Bacterial Infection with <i>Listeria monocytogenes</i> . Journal of Immunology, 2008, 180, 5991-5999.	0.8	83
21	Tissue distribution, regulation and intracellular localization of murine CD1 molecules. Molecular Immunology, 1998, 35, 525-536.	2.2	82
22	The natural killer T?cell ligand ?-galactosylceramide prevents or promotes pristane-induced lupus in mice. European Journal of Immunology, 2005, 35, 1143-1154.	2.9	81
23	CD1d-Specific NK1.1+ T Cells with a Transgenic Variant TCR. Journal of Immunology, 2000, 165, 168-174.	0.8	74
24	MHC Class Ib-Restricted CTL Provide Protection Against Primary and Secondary <i>Listeria monocytogenes</i> Infection. Journal of Immunology, 2000, 165, 5192-5201.	0.8	73
25	Characterization of the Natural Killer T-Cell Response in an Adoptive Transfer Model of Atherosclerosis. American Journal of Pathology, 2007, 170, 1100-1107.	3.8	71
26	Induction of M3-Restricted Cytotoxic T Lymphocyte Responses by N-Formylated Peptides Derived from Mycobacterium tuberculosis. Journal of Experimental Medicine, 2001, 193, 1213-1220.	8.5	65
27	Dysregulation of CD1d-Restricted Type II Natural Killer T Cells Leads to Spontaneous Development of Colitis in Mice. Gastroenterology, 2012, 142, 326-334.e2.	1.3	65
28	The Majority of H2-M3 Is Retained Intracellularly in a Peptide-Receptive State and Traffics to the Cell Surface in the Presence of N-Formylated Peptides. Journal of Experimental Medicine, 1999, 190, 423-434.	8.5	64
29	Essential role of TNF family molecule LIGHT as a cytokine in the pathogenesis of hepatitis. Journal of Clinical Investigation, 2006, 116, 1045-1051.	8.2	62
30	Polyclonal type II natural killer T cells require PLZF and SAP for their development and contribute to CpG-mediated antitumor response. Proceedings of the National Academy of Sciences of the United States of America, 2014, 111, 2674-2679.	7.1	61
31	CD1b-autoreactive T cells contribute to hyperlipidemia-induced skin inflammation in mice. Journal of Clinical Investigation, 2017, 127, 2339-2352.	8.2	59
32	CD1d deficiency exacerbates inflammatory dermatitis in MRL-lpr/lpr mice. European Journal of Immunology, 2004, 34, 1723-1732.	2.9	58
33	Mycolic acid-specific T cells protect against Mycobacterium tuberculosis infection in a humanized transgenic mouse model. ELife, 2015, 4, .	6.0	55
34	The Functions of Type I and Type II Natural Killer T Cells in Inflammatory Bowel Diseases. Inflammatory Bowel Diseases, 2013, 19, 1330-1338.	1.9	53
35	Impaired response to Listeria in H2-M3–deficient mice reveals a nonredundant role of MHC class Ib–specific T cells in host defense. Journal of Experimental Medicine, 2006, 203, 449-459.	8.5	52
36	Recognition of the nonclassical MHC class I molecule H2-M3 by the receptor Ly49A regulates the licensing and activation of NK cells. Nature Immunology, 2012, 13, 1171-1177.	14.5	49

#	Article	IF	CITATIONS
37	Differential requirements for the Ets transcription factor Elf-1 in the development of NKT cells and NK cells. Blood, 2011, 117, 1880-1887.	1.4	48
38	MHC lb molecule Qa-1 presents Mycobacterium tuberculosis peptide antigens to CD8+ T cells and contributes to protection against infection. PLoS Pathogens, 2017, 13, e1006384.	4.7	47
39	Long-term loss of canonical NKT cells following an acute virus infection. European Journal of Immunology, 2005, 35, 879-889.	2.9	45
40	IFN-β-Mediated Up-Regulation of CD1d in Bacteria-Infected APCs. Journal of Immunology, 2006, 177, 7841-7848.	0.8	43
41	A Cell-Type Specific CD1d Expression Program Modulates Invariant NKT Cell Development and Function. Journal of Immunology, 2006, 176, 1421-1430.	0.8	40
42	The Selection of M3-Restricted T Cells Is Dependent on M3 Expression and Presentation of N-Formylated Peptides in the Thymus. Journal of Experimental Medicine, 1999, 190, 1869-1878.	8.5	39
43	Role of Group 1 CD1-Restricted T Cells in Infectious Disease. Frontiers in Immunology, 2015, 6, 337.	4.8	39
44	Autoreactive CD1b-restricted T cells: a new innate-like T-cell population that contributes to immunity against infection. Blood, 2011, 118, 3870-3878.	1.4	38
45	Tapasin Enhances Peptide-Induced Expression of H2-M3 Molecules, but Is Not Required for the Retention of Open Conformers. Journal of Immunology, 2001, 167, 2097-2105.	0.8	37
46	Induction of Mycobacterium Tuberculosis Lipid-Specific T Cell Responses by Pulmonary Delivery of Mycolic Acid-Loaded Polymeric Micellar Nanocarriers. Frontiers in Immunology, 2018, 9, 2709.	4.8	37
47	Polymorphisms in CD1d affect antigen presentation and the activation of CD1d-restricted T cells. Proceedings of the National Academy of Sciences of the United States of America, 2009, 106, 1909-1914.	7.1	33
48	Functional Roles of TAP and Tapasin in the Assembly of M3- <i>N</i> -Formylated Peptide Complexes. Journal of Immunology, 2001, 167, 1507-1514.	0.8	32
49	Transcriptional Regulation of <i>CD1D1</i> by Ets Family Transcription Factors. Journal of Immunology, 2005, 175, 1022-1029.	0.8	32
50	Organization and structure of the H-2M4-M8 class I genes in the mouse major histocompatibility complex. Immunogenetics, 1993, 38, 258-71.	2.4	31
51	Crosstalk between type II NKT cells and T cells leads to spontaneous chronic inflammatory liver disease. Journal of Hepatology, 2017, 67, 791-800.	3.7	31
52	An MHC class Ib–restricted CD8 T cell response confers antiviral immunity. Journal of Experimental Medicine, 2008, 205, 1647-1657.	8.5	30
53	The Lysine Acetyltransferase GCN5 Is Required for iNKT Cell Development through EGR2 Acetylation. Cell Reports, 2017, 20, 600-612.	6.4	30
54	Rat RT1 orthologs of mouse H2-M class lb genes. Immunogenetics, 1995, 42, 63-67.	2.4	29

#	Article	IF	CITATIONS
55	Positive selecting cell type determines the phenotype of MHC class Ib-restricted CD8 ⁺ T cells. Proceedings of the National Academy of Sciences of the United States of America, 2011, 108, 13241-13246.	7.1	27
56	Expression of CD1d Under the Control of a MHC Class Ia Promoter Skews the Development of NKT Cells, But Not CD8+ T Cells. Journal of Immunology, 2003, 171, 4105-4112.	0.8	25
57	H2–M3-restricted T cell response to infection. Microbes and Infection, 2006, 8, 2277-2283.	1.9	25
58	Modulation of NKT Cell Development by B7-CD28 Interaction: An Expanding Horizon for Costimulation. PLoS ONE, 2008, 3, e2703.	2.5	24
59	Bacterial infection alters the kinetics and function of iNKT cell responses. Journal of Leukocyte Biology, 2008, 84, 1462-1471.	3.3	22
60	CD1b-autoreactive T cells recognize phospholipid antigens and contribute to antitumor immunity against a CD1b ⁺ T cell lymphoma. OncoImmunology, 2016, 5, e1213932.	4.6	22
61	Mitochondrial metabolism is essential for invariant natural killer T cell development and function. Proceedings of the National Academy of Sciences of the United States of America, 2021, 118, .	7.1	20
62	Nonclassical MHC Ib-restricted CD8+ T Cells Recognize Mycobacterium tuberculosis-Derived Protein Antigens and Contribute to Protection Against Infection. PLoS Pathogens, 2016, 12, e1005688.	4.7	20
63	Affinity of thymic self-peptides for the TCR determines the selection of CD8+ T lymphocytes in the thymus. International Immunology, 2000, 12, 1353-1363.	4.0	19
64	Activating Transcription Factor/cAMP Response Element Binding Protein Family Member Regulated Transcription of CD1A. Journal of Immunology, 2006, 177, 7024-7032.	0.8	17
65	Antagonistic Effect of Toll-Like Receptor Signaling and Bacterial Infections on Transplantation Tolerance. Transplantation, 2009, 87, S77-S79.	1.0	17
66	Nonconventional CD8+ T Cell Responses to <i>Listeria</i> Infection in Mice Lacking MHC Class Ia and H2-M3. Journal of Immunology, 2011, 186, 489-498.	0.8	17
67	Linking CD1-Restricted T Cells With Autoimmunity and Dyslipidemia: Lipid Levels Matter. Frontiers in Immunology, 2018, 9, 1616.	4.8	17
68	Human CD1d Functions as a Transplantation Antigen and a Restriction Element in Mice. Journal of Immunology, 2001, 166, 3829-3836.	0.8	15
69	CD1d Expression in Paneth Cells and Rat Exocrine Pancreas Revealed by Novel Monoclonal Antibodies Which Differentially Affect NKT Cell Activation. PLoS ONE, 2010, 5, e13089.	2.5	15
70	USP22 controls iNKT immunity through MED1 suppression of histone H2A monoubiquitination. Journal of Experimental Medicine, 2020, 217, .	8.5	15
71	The adaptor protein SAP regulates type II NKTâ€cell development, cytokine production, and cytotoxicity against lymphoma. European Journal of Immunology, 2014, 44, 3646-3657.	2.9	11
72	Group 1 CD1-restricted T cells contribute to control of systemic Staphylococcus aureus infection. PLoS Pathogens, 2020, 16, e1008443.	4.7	11

#	Article	IF	CITATIONS
73	Comparative Contribution of CD1 on the Development of CD4+ and CD8+ T Cell Compartments. Journal of Immunology, 2000, 164, 739-745.	0.8	10
74	SAP Is Required for the Development of Innate Phenotype in H2-M3–Restricted CD8+ T Cells. Journal of Immunology, 2012, 189, 4787-4796.	0.8	10
75	Pak2 Controls Acquisition of NKT Cell Fate by Regulating Expression of the Transcription Factors PLZF and Egr2. Journal of Immunology, 2015, 195, 5272-5284.	0.8	8
76	Type II Natural Killer T Cells Contribute to Protection Against Systemic Methicillin-Resistant Staphylococcus aureus Infection. Frontiers in Immunology, 2020, 11, 610010.	4.8	8
77	Identification, expression, and crystallization of the proteaseâ€resistant conserved domain of synapsin I. Protein Science, 1997, 6, 2264-2267.	7.6	7
78	Type II natural killer T cells foster the antitumor activity of CpG-oligodeoxynucleotides. Oncolmmunology, 2014, 3, e28977.	4.6	5
79	Invariant Natural Killer T-Cells and Total CD1d Restricted Cells Differentially Influence Lipid Metabolism and Atherosclerosis in Low Density Receptor Deficient Mice. International Journal of Molecular Sciences, 2019, 20, 4566.	4.1	5
80	Role of Group 1 CD1-restricted T Cells in Host Defense and Inflammatory Diseases. Critical Reviews in Immunology, 2021, 41, 1-21.	0.5	3
81	CD1-Restricted T Cells in Inflammatory Skin Diseases. Journal of Investigative Dermatology, 2021, , .	0.7	2
82	Structure and dynamics of major histocompatibility class Ib molecule H2-M3 complexed with mitochondrial-derived peptides. Journal of Biomolecular Structure and Dynamics, 2022, 40, 10300-10312.	3.5	1
83	Abstract B21: CD1 d-expressing breast cancer cells promote iNKT-mediated antitumor immunity in a mouse model of breast cancer bone metastasis. , 2010, , .		0