

Ewa Kwiatkowska

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7108004/ewa-kwiatkowska-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

45
papers

279
citations

9
h-index

14
g-index

55
ext. papers

388
ext. citations

3.1
avg, IF

3.12
L-index

#	Paper	IF	Citations
45	Effectiveness of Different Algorithms and Cut-off Value in Preeclampsia First Trimester Screening.. <i>Journal of Pregnancy</i> , 2022 , 2022, 6414857	2.5	
44	The Most Promising Biomarkers of Allogeneic Kidney Transplant Rejection. <i>Journal of Immunology Research</i> , 2022 , 2022, 1-18	4.5	1
43	The role of uropathogenic Escherichia coli adhesive molecules in inflammatory response-comparative study on immunocompetent hosts and kidney recipients. <i>PLoS ONE</i> , 2022 , 17, e0268243	3.7	
42	Molecular Pathways of Cellular Senescence and Placental Aging in Late Fetal Growth Restriction and Stillbirth. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	3
41	Application of Telehealth in Prenatal Care during the COVID-19 Pandemic-A Cross-Sectional Survey of Polish Women. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	5
40	The Mechanism of Drug Nephrotoxicity and the Methods for Preventing Kidney Damage. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	17
39	Diagnosis of placental insufficiency independently of clinical presentations using sFlt-1/PLGF ratio, including SGA patients. <i>Pregnancy Hypertension</i> , 2021 , 25, 244-248	2.6	1
38	The Clinical Importance of IL-6, IL-8, and TNF- α in Patients with Ovarian Carcinoma and Benign Cystic Lesions. <i>Diagnostics</i> , 2021 , 11,	3.8	3
37	Podocytes-The Most Vulnerable Renal Cells in Preeclampsia. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	8
36	The sFlt-1/PLGF ratio values within the 85 brackets as compared to perinatal outcomes. <i>Journal of Perinatal Medicine</i> , 2019 , 47, 732-740	2.7	8
35	The role of disordered angiogenesis tissue markers (sflt-1, Plgf) in present day diagnosis of preeclampsia. <i>Ginekologia Polska</i> , 2019 , 90, 173-176	1	6
34	C/D Ratio in Long-Term Renal Function. <i>Transplantation Proceedings</i> , 2019 , 51, 3265-3270	1.1	5
33	Do the physiological aging of the placenta and the changes in angiogenesis marker sFlt-1 and PLGF concentrations predispose patients to late-onset preeclampsia?. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2019 , 32, 11-20	2	9
32	Plasma concentration of urokinase plasminogen activator receptor is a marker of kidney allograft function. <i>Irish Journal of Medical Science</i> , 2018 , 187, 1083-1087	1.9	1
31	sFlt-1/PLGF and Doppler ultrasound parameters in SGA pregnancies with confirmed neonatal birth weight below 10th percentile. <i>Pregnancy Hypertension</i> , 2018 , 14, 79-85	2.6	9
30	Association Between Plasma Concentration of Klotho Protein, Osteocalcin, Leptin, Adiponectin, and Bone Mineral Density in Patients with Chronic Kidney Disease. <i>Hormone and Metabolic Research</i> , 2018 , 50, 816-821	3.1	8
29	Urinary IL-8 is a marker of early and long-term graft function after renal transplantation. <i>Renal Failure</i> , 2017 , 39, 484-490	2.9	5

28	Maternal endothelial damage as a disorder shared by early preeclampsia, late preeclampsia and intrauterine growth restriction. <i>Journal of Perinatal Medicine</i> , 2017 , 45, 793-802	2.7	10
27	Activity of urine arylsulfatase A in brain-dead graft donors is a predictor of early and late graft function. <i>Postepy Higieny I Medycyny Doswiadczalnej</i> , 2017 , 71, 1-4	0.3	
26	Activity of urine arylsulfatase A in brain-dead graft donors is a predictor of early and late graft function. <i>Postepy Higieny I Medycyny Doswiadczalnej</i> , 2017 , 71, 1-4	0.3	1
25	Clinical and Biochemical Characteristics of Brain-Dead Donors as Predictors of Early- and Long-Term Renal Function After Transplant. <i>Experimental and Clinical Transplantation</i> , 2017 , 15, 387-393	0.8	1
24	Ischemic placental syndrome--prediction and new disease monitoring. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2016 , 29, 2033-9	2	3
23	Urinary Metalloproteinases-9 and -2 and Their Inhibitors TIMP-1 and TIMP-2 are Markers of Early and Long-Term Graft Function After Renal Transplantation. <i>Kidney and Blood Pressure Research</i> , 2016 , 41, 288-97	3.1	19
22	Development of a focal segmental glomerulosclerosis after pregnancy complicated by preeclampsia: case report and review of literature. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2016 , 29, 1566-9	2	8
21	A Common Profile of Disordered Angiogenic Factor Production and the Exacerbation of Inflammation in Early Preeclampsia, Late Preeclampsia, and Intrauterine Growth Restriction. <i>PLoS ONE</i> , 2016 , 11, e0165060	3.7	18
20	Kidney Allograft Telomere Length Is Not Associated with Sex, Recipient Comorbid Conditions, Post-Transplant Infections, or CMV Reactivation. <i>Annals of Transplantation</i> , 2016 , 21, 392-9	1.4	1
19	Using Doppler ultrasound of the uterine and umbilical arteries and disordered angiogenesis markers (sFlt-1/PLGF) in unified monitoring of ischemic placental syndrome patients. <i>Hypertension in Pregnancy</i> , 2016 , 35, 490-498	2	1
18	hTERT, BICD1 and chromosome 18 polymorphisms associated with telomere length affect kidney allograft function after transplantation. <i>Kidney and Blood Pressure Research</i> , 2015 , 40, 111-20	3.1	5
17	Effect of delayed graft function, acute rejection and chronic allograft dysfunction on kidney allograft telomere length in patients after transplantation: a prospective cohort study. <i>BMC Nephrology</i> , 2015 , 16, 23	2.7	15
16	IL2-IL21 gene cluster polymorphism is not associated with allograft function after kidney transplantation. <i>International Urology and Nephrology</i> , 2014 , 46, 2415-20	2.3	5
15	Urinary lysosomal enzyme excretion in pregnant women with hypertensive disorders. <i>Hypertension in Pregnancy</i> , 2014 , 33, 349-59	2	2
14	Gamma-glutamyl transpeptidase as the marker of kidney graft function. <i>Advances in Clinical and Experimental Medicine</i> , 2014 , 23, 947-52	1.8	4
13	N-acetyl-beta-glucosaminidase urine activity as a marker of early proximal tubule damage and a predictor of the long-term function of the transplanted kidneys. <i>Acta Biochimica Polonica</i> , 2014 , 61, 275-80	2	3
12	The impact of CTLA4 and PTPN22 genes polymorphisms on long-term renal allograft function and transplant outcomes. <i>Renal Failure</i> , 2013 , 35, 1223-7	2.9	9
11	Correlation between ICAM1 and VCAM1 gene polymorphisms and histopathological changes in kidney allograft biopsies. <i>Archives of Medical Science</i> , 2013 , 9, 276-82	2.9	9

10	Minimal-Change Disease Secondary to <i>Borrelia burgdorferi</i> Infection. <i>Case Reports in Nephrology</i> , 2012 , 2012, 294532	0.8	3
9	Successful pregnancy in the patient with Fanconi-Bickel syndrome undergoing daily hemodialysis. <i>American Journal of Medical Genetics, Part A</i> , 2011 , 155A, 2028-30	2.5	6
8	Urinary hepatocyte growth factor indicates ischemia/reperfusion injury after kidney transplantation 2010 , 120, 437-42		2
7	Influence of glucose in the dialysate on the activity of erythrocyte-glutathione-peroxidase and blood selenium concentration in hemodialyzed patients. <i>Archives of Medical Research</i> , 2007 , 38, 330-6	6.6	9
6	Effect of hemodialysis on the content of fatty acids in monolayers of erythrocyte membranes in patients with chronic renal failure. <i>Renal Failure</i> , 2007 , 29, 447-52	2.9	9
5	GNB3 C825T and ACE I/D polymorphisms on the sodium-proton exchanger and the prevalence of essential hypertension in males. <i>Archives of Medical Research</i> , 2006 , 37, 150-7	6.6	11
4	Erythrocyte antioxidant defense system in patients with chronic renal failure according to the hemodialysis conditions. <i>Archives of Medical Research</i> , 2006 , 37, 353-9	6.6	17
3	Does glucose present in the dialysate limit oxidative stress in patients undergoing regular hemodialysis?. <i>Blood Purification</i> , 2005 , 23, 219-25	3.1	6
2	Trace elements modify the activity of sodium transporting systems in erythrocyte membrane in patients with essential hypertension-preliminary study. <i>Nephrology Dialysis Transplantation</i> , 2005 , 20, 469-71	4.3	8
1	Do trace elements modify the activity of erythrocyte sodium-proton exchanger in hemodialyzed patients?. <i>Biological Trace Element Research</i> , 2005 , 104, 107-20	4.5	3