

# Charlotte Edwardson

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/710930/charlotte-edwardson-publications-by-year.pdf>

**Version:** 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

115  
papers

5,010  
citations

29  
h-index

69  
g-index

131  
ext. papers

6,313  
ext. citations

4  
avg, IF

5.56  
L-index

#	Paper	IF	Citations
115	ACCEPTANCE: protocol for a feasibility study of a multicomponent physical activity intervention following treatment for cervical cancer.. <i>BMJ Open</i> , <b>2022</b> , 12, e048203	3	
114	Ethnic differences in the relationship between step cadence and physical function in older adults.. <i>Journal of Sports Sciences</i> , <b>2022</b> , 1-8	3.6	
113	Relative protein intake and associations with markers of physical function in those with type 2 diabetes.. <i>Diabetic Medicine</i> , <b>2022</b> , e14851	3.5	1
112	Differences in Accelerometer-Measured Patterns of Physical Activity and Sleep/Rest Between Ethnic Groups and Age: An Analysis of UK Biobank. <i>Journal of Physical Activity and Health</i> , <b>2021</b> , 1-10	2.5	1
111	Management of fatigue with physical activity and behavioural change support in vasculitis: a feasibility study. <i>Rheumatology</i> , <b>2021</b> , 60, 4130-4140	3.9	4
110	Sit Less and Move More-A Multicomponent Intervention With and Without Height-Adjustable Workstations in Contact Center Call Agents: A Pilot Randomized Controlled Trial. <i>Journal of Occupational and Environmental Medicine</i> , <b>2021</b> , 63, 44-56	2	2
109	Wrist-worn accelerometers: recommending ~1.0 m as the minimum clinically important difference (MCID) in daily average acceleration for inactive adults. <i>British Journal of Sports Medicine</i> , <b>2021</b> , 55, 814-815	10.3	9
108	A randomised-controlled feasibility study of the REgulate your Sitting Time (RESIT) intervention for reducing sitting time in individuals with type 2 diabetes: study protocol. <i>Pilot and Feasibility Studies</i> , <b>2021</b> , 7, 76	1.9	1
107	Concurrent screen use and cross-sectional association with lifestyle behaviours and psychosocial health in adolescent females. <i>Acta Paediatrica, International Journal of Paediatrics</i> , <b>2021</b> , 110, 2164-2170	3.1	1
106	The impact of COVID-19 restrictions on accelerometer-assessed physical activity and sleep in individuals with type 2 diabetes. <i>Diabetic Medicine</i> , <b>2021</b> , 38, e14549	3.5	9
105	Stand Out in Class: Investigating the Potential Impact of a Sit-Stand Desk Intervention on Children's Sitting and Physical Activity during Class Time and after School. <i>International Journal of Environmental Research and Public Health</i> , <b>2021</b> , 18,	4.6	1
104	Increasing physical activity levels following treatment for cervical cancer: an intervention mapping approach. <i>Journal of Cancer Survivorship</i> , <b>2021</b> , 1	5.1	3
103	Device-assessed total and prolonged sitting time: associations with anxiety, depression, and health-related quality of life in adults. <i>Journal of Affective Disorders</i> , <b>2021</b> , 287, 107-114	6.6	5
102	Promoting physical activity in a multi-ethnic population at high risk of diabetes: the 48-month PROPELS randomised controlled trial. <i>BMC Medicine</i> , <b>2021</b> , 19, 130	11.4	1
101	Effect of exercise on sleep and bi-directional associations with accelerometer-assessed physical activity in men with obesity. <i>Applied Physiology, Nutrition and Metabolism</i> , <b>2021</b> , 46, 597-605	3	1
100	Improvements in Glycemic Control After Acute Moderate-Intensity Continuous or High-Intensity Interval Exercise Are Greater in South Asians Than White Europeans With Nondiabetic Hyperglycemia: A Randomized Crossover Study. <i>Diabetes Care</i> , <b>2021</b> , 44, 201-209	14.6	2
99	Association of Timing and Balance of Physical Activity and Rest/Sleep With Risk of COVID-19: A UK Biobank Study. <i>Mayo Clinic Proceedings</i> , <b>2021</b> , 96, 156-164	6.4	14

98	Device-measured physical activity and its association with physical function in adults with type 2 diabetes mellitus. <i>Diabetic Medicine</i> , <b>2021</b> , 38, e14393	3.5	3
97	Comparing 24 h physical activity profiles: Office workers, women with a history of gestational diabetes and people with chronic disease condition(s). <i>Journal of Sports Sciences</i> , <b>2021</b> , 39, 219-226	3.6	2
96	Development of an Interactive Lifestyle Programme for Adolescents at Risk of Developing Type 2 Diabetes: PRE-START. <i>Children</i> , <b>2021</b> , 8,	2.8	1
95	Sleep duration and sleep efficiency in UK long-distance heavy goods vehicle drivers. <i>Occupational and Environmental Medicine</i> , <b>2021</b> ,	2.1	2
94	Normative wrist-worn accelerometer values for self-paced walking and running: a walk in the park. <i>Journal of Sports Sciences</i> , <b>2021</b> , 1-8	3.6	0
93	Feature selection for unsupervised machine learning of accelerometer data physical activity clusters - A systematic review. <i>Gait and Posture</i> , <b>2021</b> , 90, 120-128	2.6	2
92	Behavioural interventions to promote physical activity in a multiethnic population at high risk of diabetes: PROPELS three-arm RCT.. <i>Health Technology Assessment</i> , <b>2021</b> , 25, 1-190	4.4	0
91	Reducing sitting at work: process evaluation of the SMARt Work (Stand More At Work) intervention. <i>Trials</i> , <b>2020</b> , 21, 403	2.8	4
90	A Cost and Cost-Benefit Analysis of the Stand More AT Work (SMARt Work) Intervention. <i>International Journal of Environmental Research and Public Health</i> , <b>2020</b> , 17,	4.6	7
89	Stand Out in Class: restructuring the classroom environment to reduce sitting time - findings from a pilot cluster randomised controlled trial. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , <b>2020</b> , 17, 55	8.4	11
88	Sedentary Time And Markers Of Physical Function In Those With Established Type 2 Diabetes. <i>Medicine and Science in Sports and Exercise</i> , <b>2020</b> , 52, 164-164	1.2	
87	SitStand desks to reduce sedentary behaviour in 9- to 10-year-olds: the Stand Out in Class pilot cluster RCT. <i>Public Health Research</i> , <b>2020</b> , 8, 1-126	1.7	5
86	Associations between physical activity and trimethylamine -oxide in those at risk of type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , <b>2020</b> , 8,	4.5	5
85	Equivalency of Sleep Estimates: Comparison of Three Research-Grade Accelerometers. <i>Journal for the Measurement of Physical Behaviour</i> , <b>2020</b> , 3, 294-303	2.3	3
84	Predictors of the Acute Postprandial Response to Breaking Up Prolonged Sitting. <i>Medicine and Science in Sports and Exercise</i> , <b>2020</b> , 52, 1385-1393	1.2	6
83	activPAL and ActiGraph Assessed Sedentary Behavior and Cardiometabolic Health Markers. <i>Medicine and Science in Sports and Exercise</i> , <b>2020</b> , 52, 391-397	1.2	11
82	Maturational timing, physical self-perceptions and physical activity in UK adolescent females: investigation of a mediated effects model. <i>Annals of Human Biology</i> , <b>2020</b> , 47, 384-390	1.7	1
81	Physical Activity for Bone Health: How Much and/or How Hard?. <i>Medicine and Science in Sports and Exercise</i> , <b>2020</b> , 52, 2331-2341	1.2	4

80	Joint associations of accelerometer measured physical activity and sedentary time with all-cause mortality: a harmonised meta-analysis in more than 44 000 middle-aged and older individuals. <i>British Journal of Sports Medicine</i> , <b>2020</b> , 54, 1499-1506	10.3	43
79	Physical behaviors and chronotype in people with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , <b>2020</b> , 8,	4.5	5
78	Prospectively Reallocating Sedentary Time: Associations with Cardiometabolic Health. <i>Medicine and Science in Sports and Exercise</i> , <b>2020</b> , 52, 844-850	1.2	4
77	Metabolic Effects of Breaking Prolonged Sitting With Standing or Light Walking in Older South Asians and White Europeans: A Randomized Acute Study. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , <b>2020</b> , 75, 139-146	6.4	28
76	Physical activity and lipidomics in a population at high risk of type 2 diabetes mellitus. <i>Journal of Sports Sciences</i> , <b>2020</b> , 38, 1150-1160	3.6	4
75	A data-driven, meaningful, easy to interpret, standardised accelerometer outcome variable for global surveillance. <i>Journal of Science and Medicine in Sport</i> , <b>2019</b> , 22, 1132-1138	4.4	20
74	Dose-response associations between accelerometry measured physical activity and sedentary time and all cause mortality: systematic review and harmonised meta-analysis. <i>BMJ, The</i> , <b>2019</b> , 366, l4570	5.9	416
73	Impact of Depression and Anxiety on Change to Physical Activity Following a Pragmatic Diabetes Prevention Program Within Primary Care: Pooled Analysis From Two Randomized Controlled Trials. <i>Diabetes Care</i> , <b>2019</b> , 42, 1847-1853	14.6	12
72	Process evaluation of the school-based Girls Active programme. <i>BMC Public Health</i> , <b>2019</b> , 19, 1187	4.1	8
71	Association of depression and anxiety with clinical, sociodemographic, lifestyle and environmental factors in South Asian and white European individuals at high risk of diabetes. <i>Diabetic Medicine</i> , <b>2019</b> , 36, 1158-1167	3.5	2
70	A multi-component intervention to sit less and move more in a contact centre setting: a feasibility study. <i>BMC Public Health</i> , <b>2019</b> , 19, 292	4.1	7
69	Differences in objectively measured physical activity and sedentary behaviour between white Europeans and south Asians recruited from primary care: cross-sectional analysis of the PROPELS trial. <i>BMC Public Health</i> , <b>2019</b> , 19, 95	4.1	15
68	Micro-costing and a cost-consequence analysis of the 'Girls Active' programme: A cluster randomised controlled trial. <i>PLoS ONE</i> , <b>2019</b> , 14, e0221276	3.7	3
67	Evaluation and refinement of the PRESTART tool for identifying 12-14 year olds at high lifetime risk of developing type 2 diabetes compared to a clinicians assessment of risk: a cross-sectional study. <i>BMC Endocrine Disorders</i> , <b>2019</b> , 19, 79	3.3	3
66	Towards a Portable Model to Discriminate Activity Clusters from Accelerometer Data. <i>Sensors</i> , <b>2019</b> , 19,	3.8	4
65	Reply to Mekary, R.A.; Ding, E.L. Isotemporal Substitution as the Gold Standard Model for Physical Activity Epidemiology: Why It Is the Most Appropriate for Activity Time Research. <i>Int. J. Environ. Res. Public Health</i> 2019, 16, 797. <i>International Journal of Environmental Research and Public Health</i> , <b>2019</b> , 16,	4.6	3
64	Providing a Basis for Harmonization of Accelerometer-Assessed Physical Activity Outcomes Across Epidemiological Datasets. <i>Journal for the Measurement of Physical Behaviour</i> , <b>2019</b> , 2, 131-142	2.3	16
63	Enhancing the value of accelerometer-assessed physical activity: meaningful visual comparisons of data-driven translational accelerometer metrics. <i>Sports Medicine - Open</i> , <b>2019</b> , 5, 47	6.1	20

62	A school-based intervention (Girls Active) to increase physical activity levels among 11- to 14-year-old girls: cluster RCT. <i>Public Health Research</i> , <b>2019</b> , 7, 1-162	1.7	9
61	Rationale and design of a cross-sectional study to investigate and describe the chronotype of patients with type 2 diabetes and the effect on glycaemic control: the CODEC study. <i>BMJ Open</i> , <b>2019</b> , 9, e027773	3	10
60	Activity Intensity, Volume, and Norms: Utility and Interpretation of Accelerometer Metrics. <i>Medicine and Science in Sports and Exercise</i> , <b>2019</b> , 51, 2410-2422	1.2	19
59	Structured lifestyle education for people with schizophrenia, schizoaffective disorder and first-episode psychosis (STEPWISE): randomised controlled trial. <i>British Journal of Psychiatry</i> , <b>2019</b> , 214, 63-73	5.4	46
58	The impact of neighbourhood walkability on the effectiveness of a structured education programme to increase objectively measured walking. <i>Journal of Public Health</i> , <b>2018</b> , 40, 82-89	3.5	
57	Beyond Cut Points: Accelerometer Metrics that Capture the Physical Activity Profile. <i>Medicine and Science in Sports and Exercise</i> , <b>2018</b> , 50, 1323-1332	1.2	57
56	Sedentary Behaviour, Diabetes, and the Metabolic Syndrome. <i>Springer Series on Epidemiology and Public Health</i> , <b>2018</b> , 193-214	0.4	
55	Sedentary Time and MRI-Derived Measures of Adiposity in Active Versus Inactive Individuals. <i>Obesity</i> , <b>2018</b> , 26, 29-36	8	11
54	Reallocating sitting time to standing or stepping through isothermal analysis: associations with markers of chronic low-grade inflammation. <i>Journal of Sports Sciences</i> , <b>2018</b> , 36, 1586-1593	3.6	9
53	Effectiveness of the 'Girls Active' school-based physical activity programme: A cluster randomised controlled trial. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , <b>2018</b> , 15, 40	8.4	29
52	Associations Between Sedentary Behaviors and Cognitive Function: Cross-Sectional and Prospective Findings From the UK Biobank. <i>American Journal of Epidemiology</i> , <b>2018</b> , 187, 441-454	3.8	31
51	Accelerometer-assessed Physical Activity in Epidemiology: Are Monitors Equivalent?. <i>Medicine and Science in Sports and Exercise</i> , <b>2018</b> , 50, 257-265	1.2	76
50	Stand Out in Class: restructuring the classroom environment to reduce sedentary behaviour in 9-10-year-olds Study protocol for a pilot cluster randomised controlled trial. <i>Pilot and Feasibility Studies</i> , <b>2018</b> , 4, 103	1.9	8
49	Stand More AT Work (SMArT Work): using the behaviour change wheel to develop an intervention to reduce sitting time in the workplace. <i>BMC Public Health</i> , <b>2018</b> , 18, 319	4.1	42
48	Efficacy of a Multicomponent Intervention to Reduce Workplace Sitting Time in Office Workers: A Cluster Randomized Controlled Trial. <i>Journal of Occupational and Environmental Medicine</i> , <b>2018</b> , 60, 787-795	2	19
47	Steps per Day Measured by Consumer Activity Trackers Worn at the Non-Dominant and Dominant Wrist Relative to a Waist-Worn Pedometer. <i>Journal for the Measurement of Physical Behaviour</i> , <b>2018</b> , 1, 2-8	2.3	3
46	Moderate increases in daily step count are associated with reduced IL6 and CRP in women with PCOS. <i>Endocrine Connections</i> , <b>2018</b> , 7, 1442-1447	3.5	3
45	Structured lifestyle education to support weight loss for people with schizophrenia, schizoaffective disorder and first episode psychosis: the STEPWISE RCT. <i>Health Technology Assessment</i> , <b>2018</b> , 22, 1-160	4.4	24

44	Structured education programme for women with polycystic ovary syndrome: a randomised controlled trial. <i>Endocrine Connections</i> , <b>2018</b> , 7, 26-35	3.5	6
43	Compliance of Adolescent Girls to Repeated Deployments of Wrist-Worn Accelerometers. <i>Medicine and Science in Sports and Exercise</i> , <b>2018</b> , 50, 1508-1517	1.2	15
42	Effectiveness of the Stand More AT (SMARt) Work intervention: cluster randomised controlled trial. <i>BMJ, The</i> , <b>2018</b> , 363, k3870	5.9	76
41	Associations of Physical Behaviours and Behavioural Reallocations with Markers of Metabolic Health: A Compositional Data Analysis. <i>International Journal of Environmental Research and Public Health</i> , <b>2018</b> , 15,	4.6	28
40	A three arm cluster randomised controlled trial to test the effectiveness and cost-effectiveness of the SMART Work & Life intervention for reducing daily sitting time in office workers: study protocol. <i>BMC Public Health</i> , <b>2018</b> , 18, 1120	4.1	11
39	The Impact of a Novel Structured Health Intervention for Truckers (SHIFT) on Physical Activity and Cardiometabolic Risk Factors. <i>Journal of Occupational and Environmental Medicine</i> , <b>2018</b> , 60, 368-376	2	9
38	Considerations when using the activPAL monitor in field-based research with adult populations. <i>Journal of Sport and Health Science</i> , <b>2017</b> , 6, 162-178	8.2	209
37	Associations of objectively measured moderate-to-vigorous-intensity physical activity and sedentary time with all-cause mortality in a population of adults at high risk of type 2 diabetes mellitus. <i>Preventive Medicine Reports</i> , <b>2017</b> , 5, 285-288	2.6	8
36	Associations of reallocating sitting time into standing or stepping with glucose, insulin and insulin sensitivity: a cross-sectional analysis of adults at risk of type 2 diabetes. <i>BMJ Open</i> , <b>2017</b> , 7, e014267	3	27
35	Reducing sedentary time in adults at risk of type 2 diabetes: process evaluation of the STAND (Sedentary Time AND Diabetes) RCT. <i>BMC Public Health</i> , <b>2017</b> , 17, 80	4.1	7
34	Associations of moderate-to-vigorous-intensity physical activity and body mass index with glycated haemoglobin within the general population: a cross-sectional analysis of the 2008 Health Survey for England. <i>BMJ Open</i> , <b>2017</b> , 7, e014456	3	6
33	The association between air pollution and type 2 diabetes in a large cross-sectional study in Leicester: The CHAMPIONS Study. <i>Environment International</i> , <b>2017</b> , 104, 41-47	12.9	23
32	Fitness Moderates Glycemic Responses to Sitting and Light Activity Breaks. <i>Medicine and Science in Sports and Exercise</i> , <b>2017</b> , 49, 2216-2222	1.2	23
31	Breaking up sedentary time with seated upper body activity can regulate metabolic health in obese high-risk adults: A randomized crossover trial. <i>Diabetes, Obesity and Metabolism</i> , <b>2017</b> , 19, 1732-1739	6.7	17
30	Associations of Physical Activity Intensities with Markers of Insulin Sensitivity. <i>Medicine and Science in Sports and Exercise</i> , <b>2017</b> , 49, 2451-2458	1.2	7
29	Study design and protocol for a mixed methods evaluation of an intervention to reduce and break up sitting time in primary school classrooms in the UK: The CLASS PAL (Physically Active Learning) Programme. <i>BMJ Open</i> , <b>2017</b> , 7, e019428	3	6
28	Walking Away from Type 2 diabetes: a cluster randomized controlled trial. <i>Diabetic Medicine</i> , <b>2017</b> , 34, 698-707	3.5	46
27	Change in Sedentary Time, Physical Activity, Bodyweight, and HbA1c in High-Risk Adults. <i>Medicine and Science in Sports and Exercise</i> , <b>2017</b> , 49, 1120-1125	1.2	8



26	STEPWISE - STRUCTURED lifestyle Education for People With SchizophrEnia: a study protocol for a randomised controlled trial. <i>Trials</i> , <b>2016</b> , 17, 475	2.8	10
25	Breaking Up Prolonged Sitting With Standing or Walking Attenuates the Postprandial Metabolic Response in Postmenopausal Women: A Randomized Acute Study. <i>Diabetes Care</i> , <b>2016</b> , 39, 130-8	14.6	171
24	Associations of mutually exclusive categories of physical activity and sedentary time with markers of cardiometabolic health in English adults: a cross-sectional analysis of the Health Survey for England. <i>BMC Public Health</i> , <b>2016</b> , 16, 25	4.1	64
23	Devices for Self-Monitoring Sedentary Time or Physical Activity: A Scoping Review. <i>Journal of Medical Internet Research</i> , <b>2016</b> , 18, e90	7.6	78
22	Intensity Thresholds on Raw Acceleration Data: Euclidean Norm Minus One (ENMO) and Mean Amplitude Deviation (MAD) Approaches. <i>PLoS ONE</i> , <b>2016</b> , 11, e0164045	3.7	55
21	Accuracy of Posture Allocation Algorithms for Thigh- and Waist-Worn Accelerometers. <i>Medicine and Science in Sports and Exercise</i> , <b>2016</b> , 48, 1085-90	1.2	54
20	Raw Accelerometer Data Analysis with GGIR R-package: Does Accelerometer Brand Matter?. <i>Medicine and Science in Sports and Exercise</i> , <b>2016</b> , 48, 1935-41	1.2	75
19	Identifying adults' valid waking wear time by automated estimation in activPAL data collected with a 24 h wear protocol. <i>Physiological Measurement</i> , <b>2016</b> , 37, 1653-1668	2.9	125
18	Is the number of fast-food outlets in the neighbourhood related to screen-detected type 2 diabetes mellitus and associated risk factors?. <i>Public Health Nutrition</i> , <b>2015</b> , 18, 1698-705	3.3	33
17	A cluster randomised controlled trial to investigate the effectiveness and cost effectiveness of the 'Girls Active' intervention: a study protocol. <i>BMC Public Health</i> , <b>2015</b> , 15, 526	4.1	25
16	PRomotion Of Physical activity through structured Education with differing Levels of ongoing Support for people at high risk of type 2 diabetes (PROPELS): study protocol for a randomized controlled trial. <i>Trials</i> , <b>2015</b> , 16, 289	2.8	16
15	Providing NHS staff with height-adjustable workstations and behaviour change strategies to reduce workplace sitting time: protocol for the Stand More AT (SMaRT) Work cluster randomised controlled trial. <i>BMC Public Health</i> , <b>2015</b> , 15, 1219	4.1	21
14	Energy expenditure during common sitting and standing tasks: examining the 1.5 MET definition of sedentary behaviour. <i>BMC Public Health</i> , <b>2015</b> , 15, 516	4.1	105
13	Objectively measured sedentary time and associations with insulin sensitivity: Importance of reallocating sedentary time to physical activity. <i>Preventive Medicine</i> , <b>2015</b> , 76, 79-83	4.3	46
12	Effect of the PPARG2 Pro12Ala Polymorphism on Associations of Physical Activity and Sedentary Time with Markers of Insulin Sensitivity in Those with an Elevated Risk of Type 2 Diabetes. <i>PLoS ONE</i> , <b>2015</b> , 10, e0124062	3.7	9
11	Differences in levels of physical activity between White and South Asian populations within a healthcare setting: impact of measurement type in a cross-sectional study. <i>BMJ Open</i> , <b>2015</b> , 5, e006181 <sup>3</sup>		23
10	A Randomised Controlled Trial to Reduce Sedentary Time in Young Adults at Risk of Type 2 Diabetes Mellitus: Project STAND (Sedentary Time ANd Diabetes). <i>PLoS ONE</i> , <b>2015</b> , 10, e0143398	3.7	47
9	The association between neighbourhood greenspace and type 2 diabetes in a large cross-sectional study. <i>BMJ Open</i> , <b>2014</b> , 4, e006076	3	61

8	Office workers' objectively measured sedentary behavior and physical activity during and outside working hours. <i>Journal of Occupational and Environmental Medicine</i> , <b>2014</b> , 56, 298-303	2	170
7	Associations of objectively measured sedentary behaviour and physical activity with markers of cardiometabolic health. <i>Diabetologia</i> , <b>2013</b> , 56, 1012-20	10.3	226
6	Sedentary time and markers of chronic low-grade inflammation in a high risk population. <i>PLoS ONE</i> , <b>2013</b> , 8, e78350	3.7	109
5	Walking away from type 2 diabetes: trial protocol of a cluster randomised controlled trial evaluating a structured education programme in those at high risk of developing type 2 diabetes. <i>BMC Family Practice</i> , <b>2012</b> , 13, 46	2.6	44
4	Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis. <i>Diabetologia</i> , <b>2012</b> , 55, 2895-905	10.3	1100
3	Association of sedentary behaviour with metabolic syndrome: a meta-analysis. <i>PLoS ONE</i> , <b>2012</b> , 7, e34916	16.7	307
2	Methods of Measurement in epidemiology: sedentary Behaviour. <i>International Journal of Epidemiology</i> , <b>2012</b> , 41, 1460-71	7.8	356
1	Comparability of Postural and Physical Activity Metrics from Different Accelerometer Brands Worn on the Thigh: Data Harmonization Possibilities. <i>Measurement in Physical Education and Exercise Science</i> , 1-12	1.9	5