

Anna Di Benedetto

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7110900/publications.pdf>

Version: 2024-02-01

35
papers

1,074
citations

516710

16
h-index

395702

33
g-index

35
all docs

35
docs citations

35
times ranked

2110
citing authors

#	ARTICLE	IF	CITATIONS
1	p53 and BLC2 Immunohistochemical Expression Across Molecular Subtypes in 1099 Early Breast Cancer Patients With Long-Term Follow-up: An Observational Study. <i>Clinical Breast Cancer</i> , 2020, 20, e761-e770.	2.4	4
2	Prognostic relevance of DNA damage and repair biomarkers in elderly patients with hormone-receptor-positive breast cancer treated with neoadjuvant hormone therapy: evidence from the real-world setting. <i>Therapeutic Advances in Medical Oncology</i> , 2019, 11, 175883591985319.	3.2	2
3	Observational study of coagulation activation in early breast cancer: development of a prognostic model based on data from the real world setting. <i>Journal of Translational Medicine</i> , 2018, 16, 129.	4.4	16
4	Serum miR-22 as potential non-invasive predictor of poor clinical outcome in newly diagnosed, uniformly treated patients with diffuse large B-cell lymphoma: an explorative pilot study. <i>Journal of Experimental and Clinical Cancer Research</i> , 2018, 37, 95.	8.6	25
5	hMENA isoforms impact NSCLC patient outcome through fibronectin/ β 1 integrin axis. <i>Oncogene</i> , 2018, 37, 5605-5617.	5.9	17
6	Body mass index modifies the relationship between γ -H2AX, a DNA damage biomarker, and pathological complete response in triple-negative breast cancer. <i>BMC Cancer</i> , 2017, 17, 101.	2.6	12
7	Expression of phosphorylated Hippo pathway kinases (MST1/2 and LATS1/2) in HER2-positive and triple-negative breast cancer patients treated with neoadjuvant therapy. <i>Cancer Biology and Therapy</i> , 2017, 18, 339-346.	3.4	22
8	Characterisation of male breast cancer: a descriptive biomarker study from a large patient series. <i>Scientific Reports</i> , 2017, 7, 45293.	3.3	50
9	Association between AXL, Hippo Transducers, and Survival Outcomes in Male Breast Cancer. <i>Journal of Cellular Physiology</i> , 2017, 232, 2246-2252.	4.1	9
10	A Case-Matched Gender Comparison Transcriptomic Screen Identifies eIF4E and eIF5 as Potential Prognostic Markers in Male Breast Cancer. <i>Clinical Cancer Research</i> , 2017, 23, 2575-2583.	7.0	16
11	Analysis of the ATR-Chk1 and ATM-Chk2 pathways in male breast cancer revealed the prognostic significance of ATR expression. <i>Scientific Reports</i> , 2017, 7, 8078.	3.3	14
12	Breast carcinomas with low amplified/equivocal HER2 by Ish: potential supporting role of multiplex ligation-dependent probe amplification. <i>Journal of Experimental and Clinical Cancer Research</i> , 2017, 36, 143.	8.6	4
13	Molecular genetic alterations in egfr CA-SSR-1 microsatellite and egfr copy number changes are associated with aggressiveness in thymoma. <i>Journal of Thoracic Disease</i> , 2016, 8, 386-395.	1.4	4
14	Neoadjuvant Sequential Docetaxel Followed by High-Dose Epirubicin in Combination With Cyclophosphamide Administered Concurrently With Trastuzumab. The DECT Trial. <i>Journal of Cellular Physiology</i> , 2016, 231, 2541-2547.	4.1	12
15	HMG-CoAR expression in male breast cancer: relationship with hormone receptors, Hippo transducers and survival outcomes. <i>Scientific Reports</i> , 2016, 6, 35121.	3.3	6
16	Topographic expression of the Hippo transducers TAZ and YAP in triple-negative breast cancer treated with neoadjuvant chemotherapy. <i>Journal of Experimental and Clinical Cancer Research</i> , 2016, 35, 62.	8.6	24
17	The Hippo transducers TAZ/YAP and their target CTGF in male breast cancer. <i>Oncotarget</i> , 2016, 7, 43188-43198.	1.8	35
18	Cdx2 Polymorphism Affects the Activities of Vitamin D Receptor in Human Breast Cancer Cell Lines and Human Breast Carcinomas. <i>PLoS ONE</i> , 2015, 10, e0124894.	2.5	21

#	ARTICLE	IF	CITATIONS
19	ATM kinase sustains HER2 tumorigenicity in breast cancer. <i>Nature Communications</i> , 2015, 6, 6886.	12.8	50
20	Androgen receptor and antiandrogen therapy in male breast cancer. <i>Cancer Letters</i> , 2015, 368, 20-25.	7.2	17
21	Predictive significance of DNA damage and repair biomarkers in triple-negative breast cancer patients treated with neoadjuvant chemotherapy: An exploratory analysis. <i>Oncotarget</i> , 2015, 6, 42773-42780.	1.8	14
22	Antiandrogen therapy in metastatic male breast cancer: results from an updated analysis in an expanded case series. <i>Breast Cancer Research and Treatment</i> , 2014, 148, 73-80.	2.5	24
23	The Hippo transducer TAZ as a biomarker of pathological complete response in HER2-positive breast cancer patients treated with trastuzumab-based neoadjuvant therapy. <i>Oncotarget</i> , 2014, 5, 9619-9625.	1.8	35
24	Epidermal growth factor receptor gene copy number may predict lapatinib sensitivity in HER2-positive metastatic breast cancer. <i>Expert Opinion on Pharmacotherapy</i> , 2013, 14, 699-706.	1.8	16
25	Overexpression of activated phospholipase C β 1 is a risk factor for distant metastases in T1 \leq T2, N0 breast cancer patients undergoing adjuvant chemotherapy. <i>International Journal of Cancer</i> , 2013, 132, 1022-1031.	5.1	41
26	miR-10b*, a master inhibitor of the cell cycle, is downregulated in human breast tumours. <i>EMBO Molecular Medicine</i> , 2012, 4, 1214-1229.	6.9	85
27	A comparative biomarker study of 514 matched cases of male and female breast cancer reveals gender-specific biological differences. <i>Breast Cancer Research and Treatment</i> , 2012, 133, 949-958.	2.5	119
28	HER2 Protein and Gene Variation between Primary and Metastatic Breast Cancer: Significance and Impact on Patient Care. <i>Clinical Cancer Research</i> , 2011, 17, 2055-2064.	7.0	92
29	Assessment of PTEN and PI3K Status in Primary Breast Cancer and Corresponding Metastases: Is It Worthwhile?. <i>Journal of Clinical Oncology</i> , 2011, 29, 2834-2835.	1.6	5
30	Increased expression of urokinase plasminogen activator and its cognate receptor in human seminomas. <i>BMC Cancer</i> , 2010, 10, 151.	2.6	14
31	Chromogenic in situ hybridization to detect EGFR gene copy number in cell blocks from fine-needle aspirates of non small cell lung carcinomas and lung metastases from colo-rectal cancer. <i>Journal of Experimental and Clinical Cancer Research</i> , 2010, 29, 125.	8.6	14
32	A divergent role for estrogen receptor-beta in node-positive and node-negative breast cancer classified according to molecular subtypes: an observational prospective study. <i>Breast Cancer Research</i> , 2008, 10, R74.	5.0	89
33	Bio-pathologic Characteristics Related to Chromosome 11 Aneusomy and Cyclin D1 Gene Status in Surgically Resected Stage I and II Breast Cancer: Identification of an Adverse Prognostic Profile. <i>American Journal of Surgical Pathology</i> , 2007, 31, 247-254.	3.7	5
34	High-mobility group A1 inhibits p53 by cytoplasmic relocalization of its proapoptotic activator HIPK2. <i>Journal of Clinical Investigation</i> , 2007, 117, 693-702.	8.2	88
35	The Cytoskeleton Regulatory Protein hMena (ENAH) Is Overexpressed in Human Benign Breast Lesions with High Risk of Transformation and Human Epidermal Growth Factor Receptor-2 α -Positive/Hormonal Receptor α -Negative Tumors. <i>Clinical Cancer Research</i> , 2006, 12, 1470-1478.	7.0	73