
Monika Migdalska-SÄ₩

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7111988/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Altered miRNA expression in pulmonary sarcoidosis. BMC Medical Genetics, 2016, 17, 2.	2.1	29
2	Evaluation of Selected MicroRNAs Expression in Remission Phase of Multiple Sclerosis and Their Potential Link to Cognition, Depression, and Disability. Journal of Molecular Neuroscience, 2017, 63, 275-282.	2.3	27
3	Expression level and methylation status of three tumor suppressor genes, DLEC1, ITGA9 and MLH1, in non-small cell lung cancer. Medical Oncology, 2016, 33, 75.	2.5	22
4	The Expression Levels of IL-4/IL-13/STAT6 Signaling Pathway Genes and SOCS3 Could Help to Differentiate the Histopathological Subtypes of Non-Small Cell Lung Carcinoma. Molecular Diagnosis and Therapy, 2018, 22, 621-629.	3.8	22
5	An assessment of the relationship between the expression of CCR7/CCL19 axis and selected regulatory miRNAs in non-small cell lung cancer. Molecular Biology Reports, 2019, 46, 5389-5396.	2.3	22
6	FHIT promoter methylation status, low protein and high mRNA levels in patients with non-small cell lung cancer. International Journal of Oncology, 2016, 49, 1175-1184.	3.3	18
7	Diagnostic value of DNA alteration: loss of heterozygosity or allelic imbalance—promising for molecular staging of prostate cancers. Medical Oncology, 2013, 30, 391.	2.5	15
8	Immunoexpression of TGF-β/Smad and VEGF-A proteins in serum and BAL fluid of sarcoidosis patients. BMC Immunology, 2015, 16, 58.	2.2	15
9	Quantitative analysis of mRNA expression levels and DNA methylation profiles of three neighboring genes: FUS1, NPRL2/G21 and RASSF1A in non-small cell lung cancer patients. Respiratory Research, 2015, 16, 76.	3.6	15
10	HPV16 E6*II gene expression in intraepithelial cervical lesions as an indicator of neoplastic grade: a pilot study. Medical Oncology, 2014, 31, 842.	2.5	13
11	Serum Extracellular Vesicle-Derived miRNAs in Patients with Non-Small Cell Lung Cancer—Search for Non-Invasive Diagnostic Biomarkers. Diagnostics, 2021, 11, 425.	2.6	13
12	Expression of STAT5, COX-2 and PIAS3 in Correlation with NSCLC Histhopathological Features. PLoS ONE, 2014, 9, e104265.	2.5	12
13	An Analysis of IL-10, IL-17A, IL-17RA, IL-23A and IL-23R Expression and Their Correlation with Clinical Course in Patients with Psoriasis. Journal of Clinical Medicine, 2021, 10, 5834.	2.4	11
14	Aberrant methylation as a main mechanism of TSGs silencing in PTC. Frontiers in Bioscience - Elite, 2011, E3, 137-157.	1.8	10
15	Significant frequency of allelic imbalance in 3p region covering RARÎ ² and MLH1 loci seems to be essential in molecular non-small cell lung cancer diagnosis. Medical Oncology, 2013, 30, 532.	2.5	10
16	Clinicopathological Significance of Overall Frequency of Allelic Loss (OFAL) in Lesions Derived from Thyroid Follicular Cell. Molecular Diagnosis and Therapy, 2019, 23, 369-382.	3.8	9
17	Assessment of the frequency of genetic alterations (LOH/MSI) in patients with intraepithelial cervical lesions with HPV infection: a pilot study. Medical Oncology, 2016, 33, 51.	2.5	8
18	Analysis of molecular markers as IL-12, IL-22 and IFN-Î ³ in correlation with a clinical course in patients with psoriasis. International Journal of Occupational Medicine and Environmental Health, 2020, 33, 635-647.	1.3	7

#	Article	IF	CITATIONS
19	Diagnostic value of PPARδand miRNA-17 expression levels in patients with non-small cell lung cancer. Scientific Reports, 2021, 11, 24136.	3.3	7
20	Investigation of Cervical Tumor Biopsies for Chromosomal Loss of Heterozygosity (LOH) and Microsatellite Instability (MSI) at the HLA II Locus in HIV-1/HPV Co-infected Women. Frontiers in Oncology, 2019, 9, 951.	2.8	6
21	Evaluation of the relationship between the IL-17A gene expression level and regulatory miRNA-9 in relation to tumor progression in patients with non-small cell lung cancer: a pilot study. Molecular Biology Reports, 2020, 47, 583-592.	2.3	6
22	Circulating miRNAs Related to Epithelial–Mesenchymal Transitions (EMT) as the New Molecular Markers in Endometriosis. Current Issues in Molecular Biology, 2021, 43, 900-916.	2.4	6
23	CTLA-4 polymorphisms (+49 A/G and -318 C/T) are important genetic determinants of AITD susceptibility and predisposition to high levels of thyroid autoantibodies in Polish children - preliminary study. Acta Biochimica Polonica, 2013, 60, 641-6.	0.5	5
24	Correlation between the Positive Effect of Vitamin D Supplementation and Physical Performance in Young Male Soccer Players. International Journal of Environmental Research and Public Health, 2022, 19, 5138.	2.6	5
25	MSI and LOH in the development and prognosis of follicular cell-derived thyroid tumours. Endokrynologia Polska, 2012, 63, 126-36.	1.0	4
26	Assessment of Wnt pathway selected gene expression levels in peripheral blood mononuclear cells (PBMCs) of postmenopausal patients with low bone mass. Bosnian Journal of Basic Medical Sciences, 2021, 21, 461-470.	1.0	2
27	Altered Cyclooxygenase-2 Expression in Pulmonary Sarcoidosis is not Related to Clinical Classifications. Inflammation, 2016, 39, 1302-1309.	3.8	1
28	Type of training has a significant influence on the GH/IGF-1 axis but not on regulating miRNAs. Biology of Sport, 2020, 37, 217-228.	3.2	1