

Tamara Djuric

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7112932/tamara-djuric-publications-by-year.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

43 papers	286 citations	10 h-index	14 g-index
45 ext. papers	341 ext. citations	3.1 avg, IF	2.71 L-index

#	Paper	IF	Citations
43	Tag Variants of LGALS-3 Containing Haplotype Block in Advanced Carotid Atherosclerosis. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2021 , 31, 106212	2.8	0
42	The association of glutathione S-transferase and deletions with myocardial infarction. <i>Free Radical Research</i> , 2021 , 55, 267-274	4	0
41	Association of PHACTR1 intronic variants with the first myocardial infarction and their effect on PHACTR1 mRNA expression in PBMCs. <i>Gene</i> , 2021 , 775, 145428	3.8	
40	The Effects of Juice Consumption on the mRNA Expression Profile in Peripheral Blood Mononuclear Cells in Subjects at Cardiovascular Risk. <i>Nutrients</i> , 2020 , 12,	6.7	1
39	Perimatrix of middle ear cholesteatoma: A granulation tissue with a specific transcriptomic signature. <i>Laryngoscope</i> , 2020 , 130, E220-E227	3.6	4
38	PHACTR1 haplotypes are associated with carotid plaque presence and affect PHACTR1 mRNA expression in carotid plaque tissue. <i>Gene</i> , 2019 , 710, 273-278	3.8	1
37	CDKN2B gene expression is affected by 9p21.3 rs10757278 in CAD patients, six months after the MI. <i>Clinical Biochemistry</i> , 2019 , 73, 70-76	3.5	3
36	The HACD4 haplotype as a risk factor for atherosclerosis in males. <i>Gene</i> , 2018 , 641, 35-40	3.8	1
35	Transcriptome-driven integrative exploration of functional state of ureter tissue affected by CAKUT. <i>Life Sciences</i> , 2018 , 212, 1-8	6.8	5
34	Left ventricular remodeling after the first myocardial infarction in association with LGALS-3 neighbouring variants rs2274273 and rs17128183 and its relative mRNA expression: a prospective study. <i>Molecular Biology Reports</i> , 2018 , 45, 2227-2236	2.8	5
33	Overview of MMP Biology and Gene Associations in Human Diseases 2017 ,		7
32	Fructose-rich diet induces gender-specific changes in expression of the renin-angiotensin system in rat heart and upregulates the ACE/AT1R axis in the male rat aorta. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2016 , 17, 1470320316642915	3	18
31	Angiotensin receptor type 1 polymorphism A1166C is associated with altered AT1R and miR-155 expression in carotid plaque tissue and development of hypoechoic carotid plaques. <i>Atherosclerosis</i> , 2016 , 248, 132-9	3.1	9
30	9p21 locus rs10757278 is associated with advanced carotid atherosclerosis in a gender-specific manner. <i>Experimental Biology and Medicine</i> , 2016 , 241, 1210-6	3.7	3
29	Transcriptome-wide based identification of miRs in congenital anomalies of the kidney and urinary tract (CAKUT) in children: the significant upregulation of tissue miR-144 expression. <i>Journal of Translational Medicine</i> , 2016 , 14, 193	8.5	6
28	Genetic Variants in the Vicinity of LGALS-3 Gene and LGALS-3 mRNA Expression in Advanced Carotid Atherosclerosis: An Exploratory Study. <i>Journal of Clinical Laboratory Analysis</i> , 2016 , 30, 1150-1157	3.7	8
27	Gender-Specific Association between Angiotensin II Type 2 Receptor -1332 A/G Gene Polymorphism and Advanced Carotid Atherosclerosis. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2016 , 25, 1622-1630	2.8	5

26	Effects of glutathione S-transferase T1 and M1 deletions on advanced carotid atherosclerosis, oxidative, lipid and inflammatory parameters. <i>Molecular Biology Reports</i> , 2014 , 41, 1157-64	2.8	6
25	The co-inertia approach in identification of specific microRNA in early and advanced atherosclerosis plaque. <i>Medical Hypotheses</i> , 2014 , 83, 11-5	3.8	22
24	The gender-specific association of CXCL16 A181V gene polymorphism with susceptibility to multiple sclerosis, and its effects on PBMC mRNA and plasma soluble CXCL16 levels: preliminary findings. <i>Journal of Neurology</i> , 2014 , 261, 1544-51	5.5	7
23	Apolipoprotein E gene polymorphisms as risk factors for carotid atherosclerosis. <i>Vojnosanitetski Pregled</i> , 2014 , 71, 362-7	0.1	5
22	MMP-1 and -3 haplotype is associated with congenital anomalies of the kidney and urinary tract. <i>Pediatric Nephrology</i> , 2014 , 29, 879-84	3.2	5
21	Basic use of DNA analyses in medicine. <i>Srce I Krvni Sudovi</i> , 2013 , 32, 104-109		
20	eNOS Glu298Asp polymorphism is associated with development of complicated plaques in patients from Serbia with advanced carotid atherosclerosis. <i>Archives of Biological Sciences</i> , 2013 , 65, 143-149	0.7	1
19	The association of V249I and T280M fractalkine receptor haplotypes with disease course of multiple sclerosis. <i>Journal of Neuroimmunology</i> , 2012 , 245, 87-92	3.5	13
18	Matrix metalloproteinase-1 promoter genotypes and haplotypes are associated with carotid plaque presence. <i>Clinical Biochemistry</i> , 2012 , 45, 1353-6	3.5	14
17	The sex-specific association of Met62Ile gene polymorphism in P-selectin glycoprotein ligand (PSGL-1) with carotid plaque presence: preliminary study. <i>Molecular Biology Reports</i> , 2012 , 39, 6479-85	2.8	3
16	The association of ACE I/D gene polymorphism with severe carotid atherosclerosis in patients undergoing carotid endarterectomy. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2012 , 13, 141-7	3	4
15	Association of MMP-8 promoter gene polymorphisms with carotid atherosclerosis: preliminary study. <i>Atherosclerosis</i> , 2011 , 219, 673-8	3.1	20
14	Interleukin 7 receptor alpha polymorphism rs6897932 and susceptibility to multiple sclerosis in the Western Balkans. <i>Multiple Sclerosis Journal</i> , 2010 , 16, 533-6	5	8
13	Plasma levels of matrix metalloproteinase-8 in patients with carotid atherosclerosis. <i>Journal of Clinical Laboratory Analysis</i> , 2010 , 24, 246-51	3	17
12	Pro12Ala gene polymorphism in the peroxisome proliferator-activated receptor gamma as a risk factor for the onset of type 2 diabetes mellitus in the Serbian population. <i>Archives of Biological Sciences</i> , 2010 , 62, 263-270	0.7	1
11	Lack of association between eNOS Glu298Asp gene polymorphism and carotid atherosclerosis in a Serbian population. <i>Clinical Chemistry and Laboratory Medicine</i> , 2009 , 47, 1573-5	5.9	1
10	Association of the MMP-3 5A/6A gene polymorphism with multiple sclerosis in patients from Serbia. <i>Journal of the Neurological Sciences</i> , 2008 , 267, 62-5	3.2	4
9	Association of MMP-3 5A/6A gene polymorphism with susceptibility to carotid atherosclerosis. <i>Clinical Biochemistry</i> , 2008 , 41, 1326-9	3.5	24

8	Matrix metalloproteinase-9 -1562 C/T gene polymorphism in Serbian patients with multiple sclerosis. <i>Journal of Neuroimmunology</i> , 2007 , 189, 147-50	3.5	21
7	X-linked angiotensin II type 2 receptor gene polymorphism -1332A/G in male patients with essential hypertension. <i>Clinica Chimica Acta</i> , 2007 , 386, 110-3	6.2	14
6	Apolipoprotein B gene polymorphisms in patients from Serbia with ischemic cerebrovascular disease. <i>Archives of Biological Sciences</i> , 2007 , 59, 303-309	0.7	2
5	Risk factors of atherosclerosis: A review of genetic epidemiology data from a Serbian population. <i>Experimental and Clinical Cardiology</i> , 2006 , 11, 78-82		1
4	Association of ACE I/D and MMP-3 5A/6A gene polymorphisms with hypertension in men from Serbia. <i>Archives of Biological Sciences</i> , 2006 , 58, 205-210	0.7	2
3	Endothelial NOS G894 T and MMP-3 5A/6A gene polymorphisms and hypertension in Serbian population. <i>Journal of Clinical Laboratory Analysis</i> , 2005 , 19, 241-6	3	11
2	Apolipoprotein(a) gene polymorphisms (TTTTA)n and G/A-914 affect Lp(a) levels in ischemic heart disease patients from Serbia. <i>Wiener Klinische Wochenschrift</i> , 2005 , 117, 406-11	2.3	3
1	Association of lipoprotein lipase gene Asn291Ser DNA polymorphism with plasma lipid levels and blood pressure levels in healthy population of Serbia. <i>Journal of Medical Biochemistry</i> , 2003 , 22, 237-242		1