

# Wonchoel Lee

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/7120328/wonchoel-lee-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

106  
papers

642  
citations

13  
h-index

20  
g-index

123  
ext. papers

756  
ext. citations

1.2  
avg, IF

4.14  
L-index

#	Paper	IF	Citations
106	sp. n. (Nematoda, Desmodorida) from South Korea with a taxonomic review of the genus.. <i>PeerJ</i> , <b>2022</b> , 10, e13010	3.1	
105	Four New Species of Zosimeidae (Copepoda: Harpacticoida) from the Southwestern Gulf of Mexico. <i>Diversity</i> , <b>2022</b> , 14, 198	2.5	1
104	On two new species of Nannopus Brady, 1880 (Copepoda: Harpacticoida: Nannopodidae) from intertidal mudflats of the Korean west coast (Yellow Sea). <i>Zootaxa</i> , <b>2021</b> , 5051, 529-549	0.5	1
103	A New Deep-Sea Enhydrosoma (Copepoda, Harpacticoida, Cletodidae) from the Northern Gulf of Mexico. <i>Taxonomy</i> , <b>2021</b> , 1, 23-36		0
102	The First Insight into the Patterns of Size and Shape Variation of a Microcerberid Isopod. <i>Water (Switzerland)</i> , <b>2021</b> , 13, 515	3	1
101	A Brief Report of Five Newly Recorded Korean Modern Benthic Foraminiferal Species. <i>Taxonomy</i> , <b>2021</b> , 1, 438-447		
100	Doolia, A New Genus of Nannopodidae (Crustacea: Copepoda: Harpacticoida) from off Jeju Island, Korea. <i>Diversity</i> , <b>2020</b> , 12, 3	2.5	0
99	Two New Marine Free-Living Nematodes from Jeju Island Together with a Review of the Genus Gammanema Cobb 1920 (Nematoda, Chromadorida, Selachinematidae). <i>Diversity</i> , <b>2020</b> , 12, 19	2.5	2
98	A New Species of Monstrillopsis Sars, 1921 (Copepoda: Monstrilloida) with an Unusually Reduced Urosome. <i>Diversity</i> , <b>2020</b> , 12, 9	2.5	1
97	Two species of Thoracostomopsidae (Nematoda: Enoplida) from Jeju Island, South Korea. <i>PeerJ</i> , <b>2020</b> , 8, e9037	3.1	2
96	A new species of the genus Huys, 2009 (Copepoda: Harpacticoida: Miraciidae) from a sublittoral zone of Hawaii. <i>PeerJ</i> , <b>2020</b> , 8, e8506	3.1	2
95	Description of two new Procerocia species (Ostracoda: Halocyprididae) from neritic waters off South Korea with an insight into the morphological and molecular diversity of the genus. <i>Zootaxa</i> , <b>2020</b> , 4896, zootaxa.4896.2.2	0.5	
94	Meiobenthic nematodes in the assessment of the relative impact of human activities on coastal marine ecosystem. <i>Environmental Monitoring and Assessment</i> , <b>2020</b> , 192, 81	3.1	6
93	Integrated Taxonomy for Halistemma Species from the Northwest Pacific Ocean. <i>Water (Switzerland)</i> , <b>2020</b> , 12, 3283	3	0
92	Six new Krithe from the Kuril-Kamchatka Trench, with the first insight into phylogeography of deep-sea ostracods. <i>Progress in Oceanography</i> , <b>2019</b> , 176, 102128	3.8	0
91	Arthropoda: Copepoda <b>2019</b> , 761-780		
90	A new species of the genus Thalassironus (Nematoda: Enoplida: Ironidae) from the coasts of South Korea. <i>Zootaxa</i> , <b>2019</b> , 4563, zootaxa.4563.3.6	0.5	3

89	New species of Jeon, Lee & Soh, 2018 (Crustacea, Copepoda, Monstrilloida) from Chuja Island, Korea. <i>ZooKeys</i> , <b>2019</b> , 33-51	1.2	2
88	Cytheroid ostracods (Crustacea) from South Korea, with description of a new species. <i>Zoosystema</i> , <b>2019</b> , 41, 419	0.7	0
87	Two new species of the similis-subgroup of Triconia B�tger-Schnack, 1999 (Copepoda, Oncaeidae) and a redescription of T. denticula Wi, Shin & Soh, 2011 from the northeastern equatorial Pacific. <i>Zoosystema</i> , <b>2019</b> , 41, 567	0.7	0
86	Bibliographic revision of Filipjev, 1927 (Nematoda: Thoracostomopsidae) with description of a new species from Jeju Island, South Korea. <i>PeerJ</i> , <b>2019</b> , 7, e8023	3.1	3
85	The chemical composition of a new "mica sandwich" foraminiferal species from the East Coast of Korea: sp. nov. <i>PeerJ</i> , <b>2019</b> , 7, e6642	3.1	1
84	Unraveling the intricate biodiversity of the benthic harpacticoid genus Nannopus (Copepoda, Harpacticoida, Nannopodidae) in Korean waters. <i>Molecular Phylogenetics and Evolution</i> , <b>2019</b> , 130, 366-379	4.1	8
83	A new genus and two new species of monstrilloid copepods (Copepoda: Monstrillidae): integrating morphological, molecular phylogenetic, and ecological evidence. <i>Journal of Crustacean Biology</i> , <b>2018</b> , 38, 45-65	0.8	10
82	Incongruence between morphological and molecular diversity in <i>Coxicerberus fukudai</i> (Ito, 1974) (Isopoda: Microcerberidea) from East Asia. <i>Journal of Crustacean Biology</i> , <b>2018</b> , 38, 315-328	0.8	2
81	First use of molecular evidence to match sexes in the Monstrilloida (Crustacea: Copepoda), and taxonomic implications of the newly recognized and described, partly -like females of Lee, Kim & Chang, 2016. <i>PeerJ</i> , <b>2018</b> , 6, e4938	3.1	5
80	A new minute ectosymbiotic harpacticoid copepod living on the sea cucumber in the East/Japan Sea. <i>PeerJ</i> , <b>2018</b> , 6, e4979	3.1	7
79	Instant taxonomy: choosing adequate characters for species delimitation and description through congruence between molecular data and quantitative shape analysis. <i>Invertebrate Systematics</i> , <b>2018</b> , 32, 551	1.2	10
78	Two new species of the genus <i>Mesocletodes</i> (Copepoda, Harpacticoida, Argestidae) from the Pacific Ocean, San Diego Trough. <i>Proceedings of the Biological Society of Washington</i> , <b>2017</b> , 130, 34-51	0.2	1
77	Effect of increased CO in seawater on survival rate of different developmental stages of the harpacticoid copepod. <i>Animal Cells and Systems</i> , <b>2017</b> , 21, 217-222	2.3	6
76	Five new species of the genus <i>Nannopus</i> (Copepoda: Harpacticoida: Nannopodidae) from intertidal mudflats of the Korean West Coast (Yellow Sea). <i>Zootaxa</i> , <b>2017</b> , 4360, 1-66	0.5	3
75	Two species of the conifera-subgroup of <i>Triconia</i> (Copepoda, Oncaeidae) from the northeastern equatorial Pacific, with a description of the unknown male of T. hirsuta. <i>Zootaxa</i> , <b>2017</b> , 4286, 347	0.5	1
74	Two new species of the genus <i>Mesocletodes</i> (Copepoda, Harpacticoida, Argestidae) from the Pacific Ocean, San Diego Trough. <i>Proceedings of the Biological Society of Washington</i> , <b>2017</b> , 130, 34-51	0.2	
73	Redescription of Two Species of <i>Triconia</i> (Copepoda, Cyclopoida, Oncaeidae) Based on Their First Records in the Tropical Pacific. <i>Hangug Hwangyeong Saengmul Haghoeji</i> , <b>2017</b> , 35, 64-82	0.3	1
72	First records of <i>Hemicyclops tanakai</i> Itoh and Nishida, 2002 and <i>Tisbe ensifer</i> Fischer, 1860 (Crustacea, Copepoda) in Korea. <i>Journal of Species Research</i> , <b>2016</b> , 5, 289-299		1

71	A new species of the genus <i>Nannopus</i> (Copepoda, Harpacticoida, Nannopodidae) from the mudflat of Ganghwa Island, Korea. <i>Proceedings of the Biological Society of Washington</i> , <b>2016</b> , 129, 212-233	0.2	4
70	Revision of <i>Cervonema</i> Wieser, 1954 and <i>Laimella</i> Cobb, 1920 (Nematoda: ĄComesomatidae) with descriptions of two species from East Sea, Korea. <i>Zootaxa</i> , <b>2016</b> , 4098, 333-57	0.5	4
69	Modern benthic foraminiferal diversity of Jeju Island and initial insights into the total foraminiferal diversity of Korea. <i>Marine Biodiversity</i> , <b>2016</b> , 46, 337-354	1.4	13
68	Harpacticoida (Copepoda) of the northern East Sea (the Sea of Japan) and the southern Sea of Okhotsk: diversity, taxocenes, and biogeographical aspects. <i>Journal of Natural History</i> , <b>2015</b> , 49, 2869-2890	0.5	5
67	Concordance between molecular and morphology-based phylogenies of Korean Enhydrosoma (Copepoda: Harpacticoida: Cletodidae) highlights important synapomorphies and homoplasies in this genus globally. <i>Zootaxa</i> , <b>2015</b> , 3990, 451-96	0.5	8
66	A new species of the genus <i>Marylynnia</i> (Nematoda: Chromadorida: Cyatholaimidae) from Gwangyang Bay, Korea. <i>Proceedings of the Biological Society of Washington</i> , <b>2015</b> , 128, 227-238	0.2	1
65	Diversity of freshwater Cladoceran species (Crustacea: Branchiopoda) in South Korea. <i>Journal of Ecology and Environment</i> , <b>2015</b> , 38, 361-366	2	3
64	A new species of the genus <i>Normanella</i> (Copepoda: Harpacticoida) from Gwangyang Bay, Korea. <i>Proceedings of the Biological Society of Washington</i> , <b>2014</b> , 127, 175-187	0.2	2
63	Introduction to the International Association of Meiobenthologists (IAM). <i>Proceedings of the Biological Society of Washington</i> , <b>2014</b> , 127, 3-4	0.2	0
62	A new species of <i>Enhydrosoma</i> (Copepoda: Harpacticoida: Cletodidae) from Korea, with redescription of <i>E. intermedia</i> and establishment of a new genus. <i>Proceedings of the Biological Society of Washington</i> , <b>2014</b> , 127, 248-283	0.2	7
61	The genus <i>Paradoxostoma</i> (Ostracoda: Podocopida: Paradoxostomatidae) from South Korea, with description of one new species. <i>Proceedings of the Biological Society of Washington</i> , <b>2014</b> , 127, 188-208	0.2	1
60	A new species of the genus <i>Tigriopus</i> (Copepoda: Harpacticoida: Harpacticidae) from Antarctica. <i>Proceedings of the Biological Society of Washington</i> , <b>2014</b> , 127, 138-154	0.2	15
59	Checklist of the freshwater Cladocera (Crustacea: Branchiopoda) of South Korea. <i>Proceedings of the Biological Society of Washington</i> , <b>2014</b> , 127, 216-228	0.2	18
58	Two new species of free-living marine nematodes (Nematoda: Oncholaimida: Enchelidiidae) from Maemul Island, Korea. <i>Zootaxa</i> , <b>2014</b> , 3785, 419-37	0.5	3
57	Two new species of the genus <i>Wellsopsyllus</i> (Copepoda; Harpacticoida; ĄParamesochridae) from the Yellow Sea. <i>Zootaxa</i> , <b>2014</b> , 3895, 346-66	0.5	0
56	Seasonal variation of the neustonic zooplankton community in southern waters of Korea. <i>Ocean Science Journal</i> , <b>2014</b> , 49, 167-181	1.1	3
55	A new genus (Copepoda, Harpacticoida, Laophontidae) from Jeju Island of Korea. <i>ZooKeys</i> , <b>2014</b> , 1-20	1.2	3
54	A new species of <i>Harpacticella</i> Sars, 1908 (Copepoda, Harpacticoida), from a tidal pool on Jeju Island, Korea. <i>ZooKeys</i> , <b>2014</b> , 13-30	1.2	1

53	Morphological and molecular affinities of two East Asian species of Stenhelia (Crustacea, Copepoda, Harpacticoida). <i>ZooKeys</i> , <b>2014</b> , 105-43	1.2	13
52	Study of morphological deformity of <i>Tigriopus japonicus</i> l. by mid-ultraviolet radiation (UVB). <i>Proceedings of the Biological Society of Washington</i> , <b>2014</b> , 127, 87-98	0.2	2
51	A Study on Meiofauna Community in the Subtidal Sediment outside of the Saemangeum Seadike in the West Coast of Korea. <i>Ocean and Polar Research</i> , <b>2014</b> , 36, 209-223		4
50	Seven new species of <i>Hatschekia</i> Poche, 1902 (Copepoda: Siphonostomatoida: Hatschekiidae) parasitic on fishes of New Caledonia, and a redescription of <i>H. cadenati</i> Nunes-Ruivo, 1954. <i>Zoosystema</i> , <b>2013</b> , 35, 377-413	0.7	2
49	Three new species of the genus <i>Paramesochra</i> T. Scott, 1892 (Copepoda: Harpacticoida: Paramesochridae) from Yellow Sea, Korea with a redescription of <i>Paramesochra similis</i> Kunz, 1936. <i>Journal of Natural History</i> , <b>2013</b> , 47, 769-803	0.5	3
48	On the ostracod genus <i>Ilyocypris</i> , with description of one new species from Korea and the first report of males of <i>I. bradyi</i> (Crustacea: Ostracoda: Podocopida). <i>Proceedings of the Biological Society of Washington</i> , <b>2013</b> , 126, 39-71	0.2	8
47	Two new Korean earthworms (Annelida, Oligochaeta, Megadrilacea, Megascolecidae). <i>ZooKeys</i> , <b>2013</b> , 35-44	1.2	2
46	Endemism of subterranean <i>Diacyclops</i> in Korea and Japan, with descriptions of seven new species of the languidoides-group and redescrptions of <i>D. brevifurcus</i> Ishida, 2006 and <i>D. suoensis</i> Ito, 1954 (Crustacea, Copepoda, Cyclopoida). <i>ZooKeys</i> , <b>2013</b> , 1-76	1.2	20
45	Torrenticolid water mites from Korea and the Russian Far East. <i>ZooKeys</i> , <b>2013</b> , 21-48	1.2	6
44	A new species of the dentipes-subgroup of <i>Triconia</i> and a redescription of <i>T. giesbrechti</i> and <i>T. elongata</i> (Copepoda: Cyclopoida: Oncaeidae) from the tropical Pacific and the Korea Strait. <i>Journal of Natural History</i> , <b>2013</b> , 47, 1707-1743	0.5	5
43	A new species of the genus <i>Pleuroxus</i> Baird (Cladocera: Anomopoda: Chydoridae) from Jeju Island, South Korea. <i>Zootaxa</i> , <b>2013</b> , 3666, 31-40	0.5	3
42	Redefinition of the parastenocaridid genus <i>Proserpinicaris</i> (Copepoda: Harpacticoida), with description of three new species from Korea. <i>Journal of Natural History</i> , <b>2012</b> , 46, 1573-1613	0.5	25
41	Cladocera (Crustacea: Branchiopoda) of the south-east of the Korean Peninsula, with twenty new records for Korea. <i>Zootaxa</i> , <b>2012</b> , 3368, 50	0.5	35
40	A review of candonid ostracods (Crustacea: Ostracoda: Podocopida) from East Asia, with descriptions of five new species from South Korea. <i>Zootaxa</i> , <b>2012</b> , 3368, 7	0.5	14
39	Two new species of <i>Apodopsyllus</i> (Copepoda, Harpacticoida) from Jeju Island, Korea. <i>Zootaxa</i> , <b>2012</b> , 3368, 128	0.5	3
38	Biodiversity of invertebrates in Korea (cover). <i>Zootaxa</i> , <b>2012</b> , 3368, 1	0.5	2
37	New species of free-living marine <i>Sabatierinae</i> (Nematoda: Monhysterida: Comesomatidae) from. <i>Zootaxa</i> , <b>2012</b> , 3368, 263	0.5	10
36	A new species of the genus <i>Cerviniopsis</i> from Sagami Bay, Japan and reinstatement of the genus <i>Neocervinia</i> , with a report on the male of <i>Neocervinia itoi</i> Lee & Yoo, 1998 (Copepoda: Harpacticoida: Aegisthidae). <i>Zootaxa</i> , <b>2012</b> , 3575, 27	0.5	2

35	Calypsophontodes gen. nov. (Copepoda: Harpacticoida: Ancorabolidae), a New Genus of Laophontodinae, Including Remarks on Ancorabolid Phylogeny. <i>Journal of Crustacean Biology</i> , <b>2012</b> , 32, 263-280	0.8	6
34	The study on highly expressed proteins as a function of an elevated ultraviolet radiation in the copepod, <i>Tigriopus japonicus</i> . <i>Ocean Science Journal</i> , <b>2012</b> , 47, 75-82	1.1	2
33	Two new candonid species from South Korea (Ostracoda, Podocopida). <i>Crustaceana</i> , <b>2012</b> , 85, 1633-1656.	0.4	6
32	A new species of the genus <i>Ilyocryptus</i> Sars, 1862 (Cladocera: Anomopoda: Ilyocryptidae) from the East Asian Palearctic. <i>Zootaxa</i> , <b>2012</b> , 3475, 36	0.5	4
31	A new species of <i>Parastenocaris</i> from Korea, with a redescription of the closely related <i>P. biwae</i> from Japan (Copepoda: Harpacticoida: Parastenocarididae). <i>Journal of Species Research</i> , <b>2012</b> , 1, 4-34		23
30	First record of the genus <i>Sinamphiascus</i> (Copepoda: Harpacticoida) from Korean waters. <i>Journal of Species Research</i> , <b>2012</b> , 1, 44-55		2
29	The first record of <i>Orthopsyllus</i> species (Copepoda: Harpacticoida: Orthopsyllidae) from Korean waters. <i>Journal of Species Research</i> , <b>2012</b> , 1, 56-67		3
28	First record of <i>Ishizakiella supralittoralis</i> (Ostracoda, Cytheroidea, Leptocytheridae) from South Korea with a key to species of the genus. <i>Journal of Species Research</i> , <b>2012</b> , 1, 68-77		2
27	A new cyclopoid copepod from Korean subterranean waters reveals an interesting connection with the Central Asian fauna (Crustacea: Copepoda: Cyclopoida). <i>Journal of Species Research</i> , <b>2012</b> , 1, 156-174		5
26	A new species of the genus <i>Heterolaophonte</i> Lang, 1948 (Copepoda: Harpacticoida: Laophontidae) from Maxwell Bay, King George Island, Antarctica. <i>Proceedings of the Biological Society of Washington</i> , <b>2011</b> , 124, 326-340	0.2	1
25	A new species of <i>Remanea</i> Klie, 1929 (Copepoda: Harpacticoida: Paramesochridae) with a redescription of the type species. <i>Journal of Natural History</i> , <b>2011</b> , 45, 2939-2964	0.5	5
24	A new species of <i>Orthopsyllus</i> Brady & Robinson, 1873 (Copepoda: Harpacticoida: Orthopsyllidae) from Maxwell Bay, King George Island, Antarctica. <i>Proceedings of the Biological Society of Washington</i> , <b>2011</b> , 124, 84-98	0.2	1
23	A new species of <i>Sentiropsis</i> (Copepoda: Harpacticoida: Pseudotachidiidae) from the upper sublittoral zone off Hyeopjae beach, Jeju Island, Korea, and a key to genera of the subfamily Danielsseniinae. <i>Proceedings of the Biological Society of Washington</i> , <b>2011</b> , 124, 179-197	0.2	1
22	The Community Structure of Meiofauna in Marian Cove, King George Island, Antarctica. <i>Ocean and Polar Research</i> , <b>2011</b> , 33, 265-280		6
21	A new species of the genus <i>Tegastes</i> (Copepoda: Harpacticoida: Tegastidae) from hydrothermal vents in the Okinawa Trough. <i>Zoological Science</i> , <b>2010</b> , 27, 678-88	0.8	6
20	A new species of the genus <i>Goniopsyllus</i> Brady (Copepoda, Harpacticoida, Clytemnestridae) from Korean waters. <i>Proceedings of the Biological Society of Washington</i> , <b>2010</b> , 123, 121-136	0.2	
19	A new species of <i>Bestiolina</i> (Crustacea: Copepoda: Calanoida) from the Yellow Sea, with notes on the zoogeography of the genus. <i>Proceedings of the Biological Society of Washington</i> , <b>2010</b> , 123, 32-46	0.2	7
18	A new species of the genus <i>Paramesochra</i> (Copepoda: Harpacticoida) from Korean waters. <i>Proceedings of the Biological Society of Washington</i> , <b>2010</b> , 123, 47-61	0.2	1



17	A new species of the genus <i>Bryocamptus</i> (Copepoda: Harpacticoida: Canthocamptidae) from Korea. <i>Proceedings of the Biological Society of Washington</i> , <b>2010</b> , 123, 204-219	0.2	2
16	Sex Ratio and Gut Contents of the Deep-Sea Harpacticoid <i>Neocervinia Itoi</i> and Other Cerviniids: A Possibility of Reduced Foraging among Males. <i>Journal of Crustacean Biology</i> , <b>2009</b> , 29, 183-191	0.8	8
15	Detecting points as developmental delay based on the life-history development and urosome deformity of the harpacticoid copepod, <i>Tigriopus japonicus sensu lato</i> , following exposure to benzo(a)pyrene. <i>Chemosphere</i> , <b>2009</b> , 76, 1435-9	8.4	16
14	A new species of the genus <i>Amphiascoides</i> (Copepoda: Harpacticoida) from Korean waters. <i>Proceedings of the Biological Society of Washington</i> , <b>2007</b> , 120, 279-292	0.2	4
13	A new species of the genus <i>Ambanus</i> (Arachnida: Araneae: Amaurobiidae) from Korea. <i>Proceedings of the Biological Society of Washington</i> , <b>2007</b> , 120, 327-336	0.2	9
12	Spatial and bathymetric trends in Harpacticoida (Copepoda) community structure in the Northern Gulf of Mexico deep-sea. <i>Journal of Experimental Marine Biology and Ecology</i> , <b>2006</b> , 330, 327-341	2.1	57
11	The complete mitochondrial genome of the intertidal copepod <i>Tigriopus</i> sp. (Copepoda, Harpacticidae) from Korea and phylogenetic considerations. <i>Journal of Experimental Marine Biology and Ecology</i> , <b>2006</b> , 333, 251-262	2.1	75
10	Seasonal and spatial variations of zooplankton in the central and southeastern Bering Sea during the mid-1990s. <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , <b>2006</b> , 53, 795-803	2.5	6
9	Two new species of the genus <i>Heteropsyllus</i> (Crustacea, Copepoda, Harpacticoida) from Jeju Island, Korea and Devon, England. <i>Journal of Natural History</i> , <b>2006</b> , 40, 1719-1745	0.5	4
8	A new species of the genus <i>Rhizothrix</i> (Copepoda: Harpacticoida: Rhizothricidae) from Korean waters. <i>Proceedings of the Biological Society of Washington</i> , <b>2005</b> , 118, 692-705	0.2	5
7	cDNA cloning of translationally controlled tumor protein/histamine releasing factor (TCTP/HRF) from the intertidal harpacticoid copepod <i>Tigriopus japonicus</i> . <i>DNA Sequence</i> , <b>2004</b> , 15, 159-63		7
6	The intertidal harpacticoid copepod <i>Tigriopus japonicus</i> (Crustacea: Copepoda) beta-actin gene: cloning, sequence and intraspecies variation. <i>DNA Sequence</i> , <b>2003</b> , 14, 279-84		13
5	A marine harpacticoid, <i>Quinquelaophonte koreana</i> sp. nov. from a sandy beach in Korea (Crustacea: Copepoda). <i>Zoological Science</i> , <b>2003</b> , 20, 657-68	0.8	5
4	A new genus of groundwater Ameiridae (Copepoda, Harpacticoida) from boreholes in Western Australia and the artificial status of <i>Stygonitocrella</i> Petkovski, 1976. <i>Bulletin of the Natural History Museum Zoology Series</i> , <b>2002</b> , 68,		3
3	New Aegisthidae (Copepoda: Harpacticoida) from western Pacific cold seeps and hydrothermal vents. <i>Zoological Journal of the Linnean Society</i> , <b>2000</b> , 129, 1-71	2.4	8
2	Proposal of <i>Marbefia</i> , gen. n. and <i>Inermiphonte</i> , gen. n., including updated keys to the species of <i>Pseudonychocamptus</i> Lang, 1944 and <i>Paralaophonte</i> Lang, 1948 (Copepoda, Harpacticoida, Laophontidae). <i>ZooKeys</i> , <b>23</b> , 1-38	1.2	3
1	Integrative Description of Cryptic <i>Tigriopus</i> Species From Korea Using MALDI-TOF MS and DNA Barcoding. <i>Frontiers in Marine Science</i> , <b>8</b> ,	4.5	2