
Olli Eskola

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7121025/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Monoacylglycerol lipase inhibitor JZL184 reduces neuroinflammatory response in APdE9 mice and in adult mouse glial cells. Journal of Neuroinflammation, 2015, 12, 81.	7.2	59
2	In Vivo PET Imaging Demonstrates Diminished Microglial Activation After Fingolimod Treatment in an Animal Model of Multiple Sclerosis. Journal of Nuclear Medicine, 2015, 56, 305-310.	5.0	57
3	Automated synthesis and purification of [18F]bromofluoromethane at high specific radioactivity. Applied Radiation and Isotopes, 2001, 54, 927-933.	1.5	56
4	Neuroinflammation Appears Early on PET Imaging and Then Plateaus in a Mouse Model of Alzheimer Disease. Journal of Nuclear Medicine, 2018, 59, 509-515.	5.0	40
5	In vivo PET imaging of beta-amyloid deposition in mouse models of Alzheimer's disease with a high specific activity PET imaging agent [18F]flutemetamol. EJNMMI Research, 2014, 4, 37.	2.5	22
6	Tracer Level Electrophilic Synthesis and Pharmacokinetics of the Hypoxia Tracer [18F]EF5. Molecular Imaging and Biology, 2012, 14, 205-212.	2.6	17
7	Novel electrophilic synthesis of 6-[18F]fluorodopamine and comprehensive biological evaluation. European Journal of Nuclear Medicine and Molecular Imaging, 2012, 39, 800-810.	6.4	13
8	Influence of transport line material on the molar activity of cyclotron produced [18F]fluoride. Nuclear Medicine and Biology, 2018, 64-65, 8-15.	0.6	10
9	Effect of genotype and age on cerebral [18F]FDG uptake varies between transgenic APPSwe-PS1dE9 and Tg2576 mouse models of Alzheimer's disease. Scientific Reports, 2019, 9, 5700.	3.3	8
10	Dimethyl fumarate decreases short-term but not long-term inflammation in a focal EAE model of neuroinflammation. EJNMMI Research, 2022, 12, 6.	2.5	7
11	Exercise training alters lipoprotein particles independent of brown adipose tissue metabolic activity. Obesity Science and Practice, 2019, 5, 258-272.	1.9	5
12	Cessation of anti-VLA-4 therapy in a focal rat model of multiple sclerosis causes an increase in neuroinflammation. EJNMMI Research, 2019, 9, 38.	2.5	4