

Sibylle Loibl

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/712130/publications.pdf>

Version: 2024-02-01

271
papers

45,952
citations

5248

83
h-index

2071

204
g-index

275
all docs

275
docs citations

275
times ranked

35170
citing authors

#	ARTICLE	IF	CITATIONS
1	Pathological complete response and long-term clinical benefit in breast cancer: the CTNeoBC pooled analysis. <i>Lancet, The</i> , 2014, 384, 164-172.	6.3	3,224
2	Personalizing the treatment of women with early breast cancer: highlights of the St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2013. <i>Annals of Oncology</i> , 2013, 24, 2206-2223.	0.6	2,805
3	Definition and Impact of Pathologic Complete Response on Prognosis After Neoadjuvant Chemotherapy in Various Intrinsic Breast Cancer Subtypes. <i>Journal of Clinical Oncology</i> , 2012, 30, 1796-1804.	0.8	2,062
4	Trastuzumab Emtansine for Residual Invasive HER2-Positive Breast Cancer. <i>New England Journal of Medicine</i> , 2019, 380, 617-628.	13.9	1,610
5	Alpelisib for <i>PIK3CA</i> -Mutated, Hormone Receptor-Positive Advanced Breast Cancer. <i>New England Journal of Medicine</i> , 2019, 380, 1929-1940.	13.9	1,582
6	Tumor-Associated Lymphocytes As an Independent Predictor of Response to Neoadjuvant Chemotherapy in Breast Cancer. <i>Journal of Clinical Oncology</i> , 2010, 28, 105-113.	0.8	1,438
7	Tumour-infiltrating lymphocytes and prognosis in different subtypes of breast cancer: a pooled analysis of 3771 patients treated with neoadjuvant therapy. <i>Lancet Oncology, The</i> , 2018, 19, 40-50.	5.1	1,327
8	Palbociclib in Hormone-Receptor-Positive Advanced Breast Cancer. <i>New England Journal of Medicine</i> , 2015, 373, 209-219.	13.9	1,239
9	Association analysis identifies 65 new breast cancer risk loci. <i>Nature</i> , 2017, 551, 92-94.	13.7	1,099
10	Tumor-Infiltrating Lymphocytes and Response to Neoadjuvant Chemotherapy With or Without Carboplatin in Human Epidermal Growth Factor Receptor 2-Positive and Triple-Negative Primary Breast Cancers. <i>Journal of Clinical Oncology</i> , 2015, 33, 983-991.	0.8	863
11	Neoadjuvant carboplatin in patients with triple-negative and HER2-positive early breast cancer (GeparSixto; GBG 66): a randomised phase 2 trial. <i>Lancet Oncology, The</i> , 2014, 15, 747-756.	5.1	810
12	Overall Survival with Palbociclib and Fulvestrant in Advanced Breast Cancer. <i>New England Journal of Medicine</i> , 2018, 379, 1926-1936.	13.9	805
13	Tucatinib, Trastuzumab, and Capecitabine for HER2-Positive Metastatic Breast Cancer. <i>New England Journal of Medicine</i> , 2020, 382, 597-609.	13.9	789
14	Adjuvant Olaparib for Patients with <i>BRCA1</i> - or <i>BRCA2</i> -Mutated Breast Cancer. <i>New England Journal of Medicine</i> , 2021, 384, 2394-2405.	13.9	764
15	Breast cancer. <i>Lancet, The</i> , 2021, 397, 1750-1769.	6.3	731
16	Long-term outcomes for neoadjuvant versus adjuvant chemotherapy in early breast cancer: meta-analysis of individual patient data from ten randomised trials. <i>Lancet Oncology, The</i> , 2018, 19, 27-39.	5.1	717
17	Polygenic Risk Scores for Prediction of Breast Cancer and Breast Cancer Subtypes. <i>American Journal of Human Genetics</i> , 2019, 104, 21-34.	2.6	711
18	Trastuzumab Beyond Progression in Human Epidermal Growth Factor Receptor 2-Positive Advanced Breast Cancer: A German Breast Group 26/Breast International Group 03-05 Study. <i>Journal of Clinical Oncology</i> , 2009, 27, 1999-2006.	0.8	685

#	ARTICLE	IF	CITATIONS
19	HER2-positive breast cancer. <i>Lancet, The</i> , 2017, 389, 2415-2429.	6.3	655
20	Sacituzumab Govitecan in Metastatic Triple-Negative Breast Cancer. <i>New England Journal of Medicine</i> , 2021, 384, 1529-1541.	13.9	601
21	Multicenter Phase II Study of Lapatinib in Patients with Brain Metastases from HER2-Positive Breast Cancer. <i>Clinical Cancer Research</i> , 2009, 15, 1452-1459.	3.2	592
22	Recommendations From an International Expert Panel on the Use of Neoadjuvant (Primary) Systemic Treatment of Operable Breast Cancer: An Update. <i>Journal of Clinical Oncology</i> , 2006, 24, 1940-1949.	0.8	579
23	Plasma <i>ESR1</i> Mutations and the Treatment of Estrogen Receptor-Positive Advanced Breast Cancer. <i>Journal of Clinical Oncology</i> , 2016, 34, 2961-2968.	0.8	573
24	Assessing Tumor-Infiltrating Lymphocytes in Solid Tumors: A Practical Review for Pathologists and Proposal for a Standardized Method from the International Immunology Biomarkers Working Group: Part 2: TILs in Melanoma, Gastrointestinal Tract Carcinomas, Non-Small Cell Lung Carcinoma and Mesothelioma, Endometrial and Ovarian Carcinomas, Squamous Cell Carcinoma of the Head and Neck, Genitourinary Carcinomas, and Primary Brain Tumors. <i>Advances in Anatomic Pathology</i> , 2017, 24, 311-335.	2.4	530
25	Addition of the PARP inhibitor veliparib plus carboplatin or carboplatin alone to standard neoadjuvant chemotherapy in triple-negative breast cancer (BrighTNess): a randomised, phase 3 trial. <i>Lancet Oncology, The</i> , 2018, 19, 497-509.	5.1	530
26	Management of elderly patients with breast cancer: updated recommendations of the International Society of Geriatric Oncology (SIOG) and European Society of Breast Cancer Specialists (EUSOMA). <i>Lancet Oncology, The</i> , 2012, 13, e148-e160.	5.1	505
27	Neoadjuvant Treatment With Trastuzumab in HER2-Positive Breast Cancer: Results From the GeparQuattro Study. <i>Journal of Clinical Oncology</i> , 2010, 28, 2024-2031.	0.8	487
28	Neoadjuvant Chemotherapy and Bevacizumab for HER2-Negative Breast Cancer. <i>New England Journal of Medicine</i> , 2012, 366, 299-309.	13.9	473
29	Assessing Tumor-Infiltrating Lymphocytes in Solid Tumors: A Practical Review for Pathologists and Proposal for a Standardized Method From the International Immunology Biomarkers Working Group: Part 1: Assessing the Host Immune Response, TILs in Invasive Breast Carcinoma and Ductal Carcinoma In Situ, Metastatic Tumor Deposits and Areas for Further Research. <i>Advances in Anatomic Pathology</i> , 2017, 24, 235-251.	2.4	469
30	Detection and HER2 Expression of Circulating Tumor Cells: Prospective Monitoring in Breast Cancer Patients Treated in the Neoadjuvant GeparQuattro Trial. <i>Clinical Cancer Research</i> , 2010, 16, 2634-2645.	3.2	463
31	Pathologic Complete Response After Neoadjuvant Chemotherapy Plus Trastuzumab Predicts Favorable Survival in Human Epidermal Growth Factor Receptor 2-Overexpressing Breast Cancer: Results From the TECHNO Trial of the AGO and GBG Study Groups. <i>Journal of Clinical Oncology</i> , 2011, 29, 3351-3357.	0.8	456
32	Novel Theranostic Opportunities Offered by Characterization of Altered Membrane Lipid Metabolism in Breast Cancer Progression. <i>Cancer Research</i> , 2011, 71, 3236-3245.	0.4	444
33	Lapatinib versus trastuzumab in combination with neoadjuvant anthracycline-taxane-based chemotherapy (GeparQuinto, GBG 44): a randomised phase 3 trial. <i>Lancet Oncology, The</i> , 2012, 13, 135-144.	5.1	425
34	The Genetic Landscape and Clonal Evolution of Breast Cancer Resistance to Palbociclib plus Fulvestrant in the PALOMA-3 Trial. <i>Cancer Discovery</i> , 2018, 8, 1390-1403.	7.7	397
35	Neoadjuvant Chemotherapy, Endocrine Therapy, and Targeted Therapy for Breast Cancer: ASCO Guideline. <i>Journal of Clinical Oncology</i> , 2021, 39, 1485-1505.	0.8	395
36	Surgery in Recurrent Ovarian Cancer: The Arbeitsgemeinschaft Gynaekologische Onkologie (AGO) DESKTOP OVAR Trial. <i>Annals of Surgical Oncology</i> , 2006, 13, 1702-1710.	0.7	367

#	ARTICLE	IF	CITATIONS
37	Customizing local and systemic therapies for women with early breast cancer: the St. Gallen International Consensus Guidelines for treatment of early breast cancer 2021. <i>Annals of Oncology</i> , 2021, 32, 1216-1235.	0.6	354
38	Intracranial Efficacy and Survival With Tucatinib Plus Trastuzumab and Capecitabine for Previously Treated HER2-Positive Breast Cancer With Brain Metastases in the HER2CLIMB Trial. <i>Journal of Clinical Oncology</i> , 2020, 38, 2610-2619.	0.8	331
39	Nab-paclitaxel versus solvent-based paclitaxel in neoadjuvant chemotherapy for early breast cancer (GeparSeptoâ€”GBC 69): a randomised, phase 3 trial. <i>Lancet Oncology</i> , The, 2016, 17, 345-356.	5.1	316
40	Response-Guided Neoadjuvant Chemotherapy for Breast Cancer. <i>Journal of Clinical Oncology</i> , 2013, 31, 3623-3630.	0.8	302
41	Germline Mutation Status, Pathological Complete Response, and Disease-Free Survival in Triple-Negative Breast Cancer. <i>JAMA Oncology</i> , 2017, 3, 1378.	3.4	300
42	Identification of ten variants associated with risk of estrogen-receptor-negative breast cancer. <i>Nature Genetics</i> , 2017, 49, 1767-1778.	9.4	289
43	Neoadjuvant Vinorelbine-Capecitabine Versus Docetaxel-Doxorubicin-Cyclophosphamide in Early Nonresponsive Breast Cancer: Phase III Randomized GeparTrio Trial. <i>Journal of the National Cancer Institute</i> , 2008, 100, 542-551.	3.0	268
44	Gonadotropin-Releasing Hormone Agonists During Chemotherapy for Preservation of Ovarian Function and Fertility in Premenopausal Patients With Early Breast Cancer: A Systematic Review and Meta-Analysis of Individual Patientâ€”Level Data. <i>Journal of Clinical Oncology</i> , 2018, 36, 1981-1990.	0.8	268
45	Cyclin E1 Expression and Palbociclib Efficacy in Previously Treated Hormone Receptorâ€”Positive Metastatic Breast Cancer. <i>Journal of Clinical Oncology</i> , 2019, 37, 1169-1178.	0.8	266
46	Effect of Luteinizing Hormoneâ€”Releasing Hormone Agonist on Ovarian Function After Modern Adjuvant Breast Cancer Chemotherapy: The GBC 37 ZORO Study. <i>Journal of Clinical Oncology</i> , 2011, 29, 2334-2341.	0.8	263
47	Prognosis of Women With Primary Breast Cancer Diagnosed During Pregnancy: Results From an International Collaborative Study. <i>Journal of Clinical Oncology</i> , 2013, 31, 2532-2539.	0.8	261
48	Increasing the dose intensity of chemotherapy by more frequent administration or sequential scheduling: a patient-level meta-analysis of 37â€”298 women with early breast cancer in 26 randomised trials. <i>Lancet</i> , The, 2019, 393, 1440-1452.	6.3	260
49	Effect of neoadjuvant anthracyclineâ€”taxane-based chemotherapy in different biological breast cancer phenotypes: overall results from the GeparTrio study. <i>Breast Cancer Research and Treatment</i> , 2010, 124, 133-140.	1.1	252
50	Clinical and molecular characteristics of HER2-low-positive breast cancer: pooled analysis of individual patient data from four prospective, neoadjuvant clinical trials. <i>Lancet Oncology</i> , The, 2021, 22, 1151-1161.	5.1	248
51	Intensified Neoadjuvant Chemotherapy in Early-Responding Breast Cancer: Phase III Randomized GeparTrio Study. <i>Journal of the National Cancer Institute</i> , 2008, 100, 552-562.	3.0	231
52	<i>PIK3CA</i> Mutations Are Associated With Lower Rates of Pathologic Complete Response to Antiâ€”Human Epidermal Growth Factor Receptor 2 (HER2) Therapy in Primary HER2-Overexpressing Breast Cancer. <i>Journal of Clinical Oncology</i> , 2014, 32, 3212-3220.	0.8	231
53	Standardized evaluation of tumor-infiltrating lymphocytes in breast cancer: results of the ring studies of the international immuno-oncology biomarker working group. <i>Modern Pathology</i> , 2016, 29, 1155-1164.	2.9	230
54	Impact of treatment characteristics on response of different breast cancer phenotypes: pooled analysis of the German neo-adjuvant chemotherapy trials. <i>Breast Cancer Research and Treatment</i> , 2011, 125, 145-156.	1.1	228

#	ARTICLE	IF	CITATIONS
55	Treatment of breast cancer during pregnancy: an observational study. <i>Lancet Oncology</i> , The, 2012, 13, 887-896.	5.1	224
56	Breast cancer in pregnancy. <i>Lancet</i> , The, 2012, 379, 570-579.	6.3	216
57	Capecitabine in Addition to Anthracycline- and Taxane-Based Neoadjuvant Treatment in Patients With Primary Breast Cancer: Phase III GeparQuattro Study. <i>Journal of Clinical Oncology</i> , 2010, 28, 2015-2023.	0.8	194
58	Prospective Validation of Immunological Infiltrate for Prediction of Response to Neoadjuvant Chemotherapy in HER2-Negative Breast Cancer – A Substudy of the Neoadjuvant GeparQuinto Trial. <i>PLoS ONE</i> , 2013, 8, e79775.	1.1	187
59	A transcriptome-wide association study of 229,000 women identifies new candidate susceptibility genes for breast cancer. <i>Nature Genetics</i> , 2018, 50, 968-978.	9.4	184
60	Androgen receptor expression in primary breast cancer and its predictive and prognostic value in patients treated with neoadjuvant chemotherapy. <i>Breast Cancer Research and Treatment</i> , 2011, 130, 477-487.	1.1	180
61	Breast Cancer Diagnosed During Pregnancy. <i>JAMA Oncology</i> , 2015, 1, 1145.	3.4	169
62	Palbociclib with adjuvant endocrine therapy in early breast cancer (PALLAS): interim analysis of a multicentre, open-label, randomised, phase 3 study. <i>Lancet Oncology</i> , The, 2021, 22, 212-222.	5.1	169
63	Advances in the treatment of advanced oestrogen-receptor-positive breast cancer. <i>Lancet</i> , The, 2017, 389, 2403-2414.	6.3	168
64	Tumor-Infiltrating Lymphocytes: A Predictive and Prognostic Biomarker in Neoadjuvant-Treated HER2-Positive Breast Cancer. <i>Clinical Cancer Research</i> , 2016, 22, 5747-5754.	3.2	158
65	Palbociclib for Residual High-Risk Invasive HR-Positive and HER2-Negative Early Breast Cancer – The Penelope-B Trial. <i>Journal of Clinical Oncology</i> , 2021, 39, 1518-1530.	0.8	153
66	Downregulation of human polo-like kinase activity by antisense oligonucleotides induces growth inhibition in cancer cells. <i>Oncogene</i> , 2002, 21, 3162-3171.	2.6	150
67	Trastuzumab for early-stage, HER2-positive breast cancer: a meta-analysis of 13 864 women in seven randomised trials. <i>Lancet Oncology</i> , The, 2021, 22, 1139-1150.	5.1	147
68	Second international consensus guidelines for breast cancer in young women (BCY2). <i>Breast</i> , 2016, 26, 87-99.	0.9	142
69	Ki67 Measured after Neoadjuvant Chemotherapy for Primary Breast Cancer. <i>Clinical Cancer Research</i> , 2013, 19, 4521-4531.	3.2	137
70	First international consensus guidelines for breast cancer in young women (BCY1). <i>Breast</i> , 2014, 23, 209-220.	0.9	135
71	Moving From Poly (ADP-Ribose) Polymerase Inhibition to Targeting DNA Repair and DNA Damage Response in Cancer Therapy. <i>Journal of Clinical Oncology</i> , 2019, 37, 2257-2269.	0.8	135
72	Use of anastrozole for breast cancer prevention (IBIS-II): long-term results of a randomised controlled trial. <i>Lancet</i> , The, 2020, 395, 117-122.	6.3	128

#	ARTICLE	IF	CITATIONS
73	Silencing of the HER2/neu Gene by siRNA Inhibits Proliferation and Induces Apoptosis in HER2/neu-Overexpressing Breast Cancer Cells. <i>Neoplasia</i> , 2004, 6, 786-795.	2.3	113
74	The tale of TILs in breast cancer: A report from The International Immuno-Oncology Biomarker Working Group. <i>Npj Breast Cancer</i> , 2021, 7, 150.	2.3	112
75	Changes in serum levels of miR-21, miR-210, and miR-373 in HER2-positive breast cancer patients undergoing neoadjuvant therapy: a translational research project within the Geparquinto trial. <i>Breast Cancer Research and Treatment</i> , 2014, 147, 61-68.	1.1	108
76	Palbociclib Combined with Fulvestrant in Premenopausal Women with Advanced Breast Cancer and Prior Progression on Endocrine Therapy: PALOMA-3 Results. <i>Oncologist</i> , 2017, 22, 1028-1038.	1.9	108
77	13th St. Gallen International Breast Cancer Conference 2013: Primary Therapy of Early Breast Cancer Evidence, Controversies, Consensus - Opinion of a German Team of Experts (Zurich 2013). <i>Breast Care</i> , 2013, 8, 221-229.	0.8	98
78	NAB-Paclitaxel Improves Disease-Free Survival in Early Breast Cancer: GBG 69 "GeparSepto". <i>Journal of Clinical Oncology</i> , 2019, 37, 2226-2234.	0.8	95
79	Response and prognosis after neoadjuvant chemotherapy in 1,051 patients with infiltrating lobular breast carcinoma. <i>Breast Cancer Research and Treatment</i> , 2014, 144, 153-162.	1.1	92
80	Impact of body mass index on neoadjuvant treatment outcome: a pooled analysis of eight prospective neoadjuvant breast cancer trials. <i>Breast Cancer Research and Treatment</i> , 2015, 150, 127-139.	1.1	92
81	Classical pathology and mutational load of breast cancer " integration of two worlds. <i>Journal of Pathology: Clinical Research</i> , 2015, 1, 225-238.	1.3	91
82	Extended adjuvant intermittent letrozole versus continuous letrozole in postmenopausal women with breast cancer (SOLE): a multicentre, open-label, randomised, phase 3 trial. <i>Lancet Oncology</i> , The, 2018, 19, 127-138.	5.1	91
83	Evaluating the impact of Relative Total Dose Intensity (RTDI) on patients' short and long-term outcome in taxane- and anthracycline-based chemotherapy of metastatic breast cancer- a pooled analysis. <i>BMC Cancer</i> , 2011, 11, 131.	1.1	87
84	BRCA1/2 Mutations and Bevacizumab in the Neoadjuvant Treatment of Breast Cancer: Response and Prognosis Results in Patients With Triple-Negative Breast Cancer From the GeparQuinto Study. <i>Journal of Clinical Oncology</i> , 2018, 36, 2281-2287.	0.8	86
85	Outcome after neoadjuvant chemotherapy in young breast cancer patients: a pooled analysis of individual patient data from eight prospectively randomized controlled trials. <i>Breast Cancer Research and Treatment</i> , 2015, 152, 377-387.	1.1	85
86	Standardized Ki67 Diagnostics Using Automated Scoring "Clinical Validation in the GeparTrio Breast Cancer Study. <i>Clinical Cancer Research</i> , 2015, 21, 3651-3657.	3.2	85
87	Prognostic Impact of Circulating Tumor Cells for Breast Cancer Patients Treated in the Neoadjuvant "Geparquattro" Trial. <i>Clinical Cancer Research</i> , 2017, 23, 5384-5393.	3.2	85
88	Intense dose-dense epirubicin, paclitaxel, cyclophosphamide "versus weekly paclitaxel, liposomal doxorubicin (plus carboplatin in triple-negative breast cancer) for neoadjuvant treatment of high-risk early breast cancer (GeparOcto "GBG 84): A randomised phase III trial. <i>European Journal of Cancer</i> , 2019, 106, 181-192.	1.3	84
89	Surgical Procedures After Neoadjuvant Chemotherapy in Operable Breast Cancer: Results of the GEPARDUO Trial. <i>Annals of Surgical Oncology</i> , 2006, 13, 1434-1442.	0.7	83
90	Diagnosis of pathological complete response to neoadjuvant chemotherapy in breast cancer by minimal invasive biopsy techniques. <i>British Journal of Cancer</i> , 2015, 113, 1565-1570.	2.9	83

#	ARTICLE	IF	CITATIONS
91	Associations of obesity and circulating insulin and glucose with breast cancer risk: a Mendelian randomization analysis. <i>International Journal of Epidemiology</i> , 2019, 48, 795-806.	0.9	81
92	Breast Conservation After Neoadjuvant Chemotherapy for Triple-Negative Breast Cancer. <i>JAMA Surgery</i> , 2020, 155, e195410.	2.2	81
93	Cytoplasmic Poly(Adenosine Diphosphate-Ribose) Polymerase Expression Is Predictive and Prognostic in Patients With Breast Cancer Treated With Neoadjuvant Chemotherapy. <i>Journal of Clinical Oncology</i> , 2011, 29, 2150-2157.	0.8	79
94	Effect of Tailored Dose-Dense Chemotherapy vs Standard 3-Weekly Adjuvant Chemotherapy on Recurrence-Free Survival Among Women With High-Risk Early Breast Cancer. <i>JAMA - Journal of the American Medical Association</i> , 2016, 316, 1888.	3.8	79
95	Genomic and Transcriptomic Analyses of Breast Cancer Primaries and Matched Metastases in AURORA, the Breast International Group (BIG) Molecular Screening Initiative. <i>Cancer Discovery</i> , 2021, 11, 2796-2811.	7.7	79
96	Impact of Multifocal or Multicentric Disease on Surgery and Locoregional, Distant and Overall Survival of 6,134 Breast Cancer Patients Treated With Neoadjuvant Chemotherapy. <i>Annals of Surgical Oncology</i> , 2015, 22, 1118-1127.	0.7	77
97	Neoadjuvant chemotherapy with paclitaxel and everolimus in breast cancer patients with non-responsive tumours to epirubicin/cyclophosphamide (EC) ± bevacizumab Results of the randomised GeparQuinto study (GBC 44). <i>European Journal of Cancer</i> , 2013, 49, 2284-2293.	1.3	75
98	The BCY3/BCC 2017 survey on physicians' knowledge, attitudes and practice towards fertility and pregnancy-related issues in young breast cancer patients. <i>Breast</i> , 2018, 42, 41-49.	0.9	75
99	AGO Recommendations for the Diagnosis and Treatment of Patients with Early Breast Cancer: Update 2019. <i>Breast Care</i> , 2019, 14, 224-245.	0.8	72
100	Intestinal microbiota influences clinical outcome and side effects of early breast cancer treatment. <i>Cell Death and Differentiation</i> , 2021, 28, 2778-2796.	5.0	72
101	Expression of endothelial and inducible nitric oxide synthase in benign and malignant lesions of the breast and measurement of nitric oxide using electron paramagnetic resonance spectroscopy. <i>Cancer</i> , 2002, 95, 1191-1198.	2.0	70
102	Interdisciplinary Screening, Diagnosis, Therapy and Follow-up of Breast Cancer. Guideline of the DGGG and the DKG (S3-Level, AWMF Registry Number 032/045OL, December 2017) Part 2 with Recommendations for the Therapy of Primary, Recurrent and Advanced Breast Cancer. <i>Geburtshilfe Und Frauenheilkunde</i> , 2018, 78, 1056-1088.	0.8	69
103	NO Signaling Confers Cytoprotectivity through the Survivin Network in Ovarian Carcinomas. <i>Cancer Research</i> , 2008, 68, 5159-5166.	0.4	68
104	Integration of Metabolomics and Expression of Glycerol-3-phosphate Acyltransferase (GPAM) in Breast Cancer Link to Patient Survival, Hormone Receptor Status, and Metabolic Profiling. <i>Journal of Proteome Research</i> , 2012, 11, 850-860.	1.8	68
105	Poor Outcome in Estrogen Receptor-Positive Breast Cancers Predicted by Loss of Plexin B1. <i>Clinical Cancer Research</i> , 2007, 13, 1115-1122.	3.2	66
106	Overall Survival with Palbociclib and Fulvestrant in Women with HR+/HER2- ABC: Updated Exploratory Analyses of PALOMA-3, a Double-blind, Phase III Randomized Study. <i>Clinical Cancer Research</i> , 2022, 28, 3433-3442.	3.2	65
107	Tyrosine kinase inhibitors for brain metastases in HER2-positive breast cancer. <i>Cancer Treatment Reviews</i> , 2018, 67, 71-77.	3.4	64
108	Pregnancy occurring during or following adjuvant trastuzumab in patients enrolled in the HERA trial (BIG 01-01). <i>Breast Cancer Research and Treatment</i> , 2012, 133, 387-391.	1.1	61

#	ARTICLE	IF	CITATIONS
109	Neoadjuvant Chemotherapy Shows Similar Response in Patients With Inflammatory or Locally Advanced Breast Cancer When Compared With Operable Breast Cancer: A Secondary Analysis of the GeparTrio Trial Data. <i>Journal of Clinical Oncology</i> , 2010, 28, 83-91.	0.8	59
110	RANK expression as a prognostic and predictive marker in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2014, 145, 307-315.	1.1	59
111	Mutational profiles in triple-negative breast cancer defined by ultradeep multigene sequencing show high rates of PI3K pathway alterations and clinically relevant entity subgroup specific differences. <i>Oncotarget</i> , 2014, 5, 9952-9965.	0.8	58
112	Using ultrasound and palpation for predicting axillary lymph node status following neoadjuvant chemotherapy – Results from the multi-center SENTINA trial. <i>Breast</i> , 2017, 31, 202-207.	0.9	57
113	The definition of pregnancy-associated breast cancer is outdated and should no longer be used. <i>Lancet Oncology</i> , The, 2021, 22, 753-754.	5.1	57
114	Comparison of BEAMing and Droplet Digital PCR for Circulating Tumor DNA Analysis. <i>Clinical Chemistry</i> , 2019, 65, 1405-1413.	1.5	53
115	Genome-wide association study of germline variants and breast cancer-specific mortality. <i>British Journal of Cancer</i> , 2019, 120, 647-657.	2.9	52
116	Taxanes in the treatment of breast cancer: Have we better defined their role in older patients? A position paper from a SIOG Task Force. <i>Cancer Treatment Reviews</i> , 2016, 43, 19-26.	3.4	51
117	AGO Recommendations for the Diagnosis and Treatment of Patients with Early Breast Cancer: Update 2021. <i>Breast Care</i> , 2021, 16, 214-227.	0.8	51
118	Role of TP53 mutations in triple negative and HER2-positive breast cancer treated with neoadjuvant anthracycline/taxane-based chemotherapy. <i>Oncotarget</i> , 2016, 7, 67686-67698.	0.8	50
119	Responsiveness of adjacent ductal carcinoma in situ and changes in HER2 status after neoadjuvant chemotherapy/trastuzumab treatment in early breast cancer – results from the GeparQuattro study (GBG 40). <i>Breast Cancer Research and Treatment</i> , 2012, 132, 863-870.	1.1	49
120	Breast conservation and axillary management after primary systemic therapy in patients with early-stage breast cancer: the Lucerne toolbox. <i>Lancet Oncology</i> , The, 2021, 22, e18-e28.	5.1	49
121	Monitoring serum HER2 levels during neoadjuvant trastuzumab treatment within the GeparQuattro trial. <i>Breast Cancer Research and Treatment</i> , 2010, 123, 437-445.	1.1	47
122	A randomized phase 2 study comparing EC or CMF versus nab-paclitaxel plus capecitabine as adjuvant chemotherapy for nonfrail elderly patients with moderate to high-risk early breast cancer (ICE II – GBG) Tj ETQq0 0.0 rgBT / Overlock 10	0.0	0
123	AGO Recommendations for the Diagnosis and Treatment of Patients with Locally Advanced and Metastatic Breast Cancer: Update 2020. <i>Breast Care</i> , 2020, 15, 294-309.	0.8	47
124	Utility of the CPS+EG staging system in hormone receptor-positive, human epidermal growth factor receptor 2-negative breast cancer treated with neoadjuvant chemotherapy. <i>European Journal of Cancer</i> , 2016, 53, 65-74.	1.3	46
125	14th St. Gallen International Breast Cancer Conference 2015: Evidence, Controversies, Consensus - Primary Therapy of Early Breast Cancer: Opinions Expressed by German Experts. <i>Breast Care</i> , 2015, 10, 211-219.	0.8	43
126	Loss of ARID1A Activates ANXA1, which Serves as a Predictive Biomarker for Trastuzumab Resistance. <i>Clinical Cancer Research</i> , 2016, 22, 5238-5248.	3.2	43

#	ARTICLE	IF	CITATIONS
127	Survival Analysis After Neoadjuvant Chemotherapy With Trastuzumab or Lapatinib in Patients With Human Epidermal Growth Factor Receptor 2-Positive Breast Cancer in the GeparQuinto (G5) Study (GBG 44). <i>Journal of Clinical Oncology</i> , 2018, 36, 1308-1316.	0.8	43
128	Association of Germline Variant Status With Therapy Response in High-risk Early-Stage Breast Cancer. <i>JAMA Oncology</i> , 2020, 6, 744.	3.4	42
129	Integrated Analysis of PTEN and p4EBP1 Protein Expression as Predictors for pCR in HER2-Positive Breast Cancer. <i>Clinical Cancer Research</i> , 2016, 22, 2675-2683.	3.2	41
130	Therapy response and prognosis of patients with early breast cancer with low positivity for hormone receptors – An analysis of 2765 patients from neoadjuvant clinical trials. <i>European Journal of Cancer</i> , 2021, 148, 159-170.	1.3	41
131	Monounsaturated fatty acids in serum triacylglycerols are associated with response to neoadjuvant chemotherapy in breast cancer patients. <i>International Journal of Cancer</i> , 2014, 134, 1725-1733.	2.3	40
132	Lucitanib for the Treatment of HR+/HER2- Metastatic Breast Cancer: Results from the Multicohort Phase II FINESSE Study. <i>Clinical Cancer Research</i> , 2020, 26, 354-363.	3.2	40
133	Dual Blockade with Afatinib and Trastuzumab as Neoadjuvant Treatment for Patients with Locally Advanced or Operable Breast Cancer Receiving Taxane-Containing Anthracycline Chemotherapy – DAFNE (GBG-70). <i>Clinical Cancer Research</i> , 2015, 21, 2924-2931.	3.2	38
134	Repurposing anticancer drugs for COVID-19-induced inflammation, immune dysfunction, and coagulopathy. <i>British Journal of Cancer</i> , 2020, 123, 694-697.	2.9	37
135	Association of Immunophenotype With Pathologic Complete Response to Neoadjuvant Chemotherapy for Triple-Negative Breast Cancer. <i>JAMA Oncology</i> , 2021, 7, 603.	3.4	37
136	Genomic correlates of response to adjuvant trastuzumab (H) and pertuzumab (P) in HER2+ breast cancer (BC): Biomarker analysis of the APHINITY trial. <i>Journal of Clinical Oncology</i> , 2019, 37, 1012-1012.	0.8	35
137	Phase II multicenter, uncontrolled trial of sorafenib in patients with metastatic breast cancer. <i>Anti-Cancer Drugs</i> , 2009, 20, 616-24.	0.7	34
138	Tumor-Infiltrating Lymphocytes: A Promising Biomarker in Breast Cancer. <i>Breast Care</i> , 2016, 11, 96-100.	0.8	33
139	Outcome after neoadjuvant chemotherapy in estrogen receptor-positive and progesterone receptor-negative breast cancer patients: a pooled analysis of individual patient data from ten prospectively randomized controlled neoadjuvant trials. <i>Breast Cancer Research and Treatment</i> , 2018, 167, 59-71.	1.1	32
140	AGO Recommendations for the Diagnosis and Treatment of Patients with Locally Advanced and Metastatic Breast Cancer: Update 2019. <i>Breast Care</i> , 2019, 14, 247-255.	0.8	32
141	Mutational Diversity and Therapy Response in Breast Cancer: A Sequencing Analysis in the Neoadjuvant GeparSepto Trial. <i>Clinical Cancer Research</i> , 2019, 25, 3986-3995.	3.2	32
142	Bevacizumab Treatment for Advanced Breast Cancer. <i>Oncologist</i> , 2011, 16, 1684-1697.	1.9	31
143	Genetic variants in VEGF pathway genes in neoadjuvant breast cancer patients receiving bevacizumab: Results from the randomized phase III GeparQuinto study. <i>International Journal of Cancer</i> , 2015, 137, 2981-2988.	2.3	31
144	Outcome after neoadjuvant chemotherapy in elderly breast cancer patients - a pooled analysis of individual patient data from eight prospectively randomized controlled trials. <i>Oncotarget</i> , 2018, 9, 15168-15179.	0.8	29

#	ARTICLE	IF	CITATIONS
145	Post-Mastectomy Radiotherapy After Neoadjuvant Chemotherapy in Breast Cancer: A Pooled Retrospective Analysis of Three Prospective Randomized Trials. <i>Annals of Surgical Oncology</i> , 2019, 26, 3892-3901.	0.7	29
146	Knowledge, attitudes and practice of physicians towards fertility and pregnancy-related issues in youngBRCA-mutated breast cancer patients. <i>Reproductive BioMedicine Online</i> , 2019, 38, 835-844.	1.1	29
147	PIK3CA H1047R Mutation Associated with a Lower Pathological Complete Response Rate in Triple-Negative Breast Cancer Patients Treated with Anthracycline-Taxane-Based Neoadjuvant Chemotherapy. <i>Cancer Research and Treatment</i> , 2020, 52, 689-696.	1.3	29
148	Everolimus as treatment for breast cancer patients with bone metastases only: results of the phase II RADAR study. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013, 139, 2047-2056.	1.2	28
149	Diagnosis and Therapy of Triple-Negative Breast Cancer (TNBC) – Recommendations for Daily Routine Practice. <i>Geburtshilfe Und Frauenheilkunde</i> , 2019, 79, 605-617.	0.8	28
150	Side Effects of Standard Adjuvant and Neoadjuvant Chemotherapy Regimens According to Age Groups in Primary Breast Cancer. <i>Breast Care</i> , 2013, 8, 60-66.	0.8	27
151	Tamoxifen treatment for male breast cancer and risk of thromboembolism: prospective cohort analysis. <i>British Journal of Cancer</i> , 2019, 120, 301-305.	2.9	27
152	Immune-related Gene Expression Predicts Response to Neoadjuvant Chemotherapy but not Additional Benefit from PD-L1 Inhibition in Women with Early Triple-negative Breast Cancer. <i>Clinical Cancer Research</i> , 2021, 27, 2584-2591.	3.2	27
153	Clinical relevance and concordance of HER2 status in local and central testing—an analysis of 1581 HER2-positive breast carcinomas over 12 years. <i>Modern Pathology</i> , 2018, 31, 607-615.	2.9	25
154	Breast cancer, placenta and pregnancy. <i>European Journal of Cancer</i> , 2019, 115, 68-78.	1.3	24
155	Efficacy of Endocrine Therapy for the Treatment of Breast Cancer in Men. <i>JAMA Oncology</i> , 2021, 7, 565.	3.4	24
156	Effect of Celecoxib vs Placebo as Adjuvant Therapy on Disease-Free Survival Among Patients With Breast Cancer. <i>JAMA Oncology</i> , 2021, 7, 1291.	3.4	24
157	Comparison of pegfilgrastim on day 2 vs. day 4 as primary prophylaxis of intense dose-dense chemotherapy in patients with node-positive primary breast cancer within the prospective, multi-center GAIN study: (GBG 33). <i>Supportive Care in Cancer</i> , 2011, 19, 1789-1795.	1.0	23
158	Surgical Outcome after Neoadjuvant Chemotherapy and Bevacizumab: Results from the GeparQuinto Study (GBG 44). <i>Annals of Surgical Oncology</i> , 2014, 21, 2517-2524.	0.7	23
159	Cardiotoxicity and Cardiovascular Biomarkers in Patients With Breast Cancer: Data From the GeparOcto-GBG 84 Trial. <i>Journal of the American Heart Association</i> , 2020, 9, e018143.	1.6	23
160	Patient-Reported Outcomes in Patients With PIK3CA-Mutated Hormone Receptor-Positive, Human Epidermal Growth Factor Receptor-Negative Advanced Breast Cancer From SOLAR-1. <i>Journal of Clinical Oncology</i> , 2021, 39, 2005-2015.	0.8	23
161	loncopy: a novel method for calling copy number alterations in amplicon sequencing data including significance assessment. <i>Oncotarget</i> , 2016, 7, 13236-13247.	0.8	23
162	New Therapeutic Options for Breast Cancer during Pregnancy. <i>Breast Care</i> , 2008, 3, 4-4.	0.8	22

#	ARTICLE	IF	CITATIONS
163	Capecitabine plus paclitaxel versus epirubicin plus paclitaxel as first-line treatment for metastatic breast cancer: efficacy and safety results of a randomized, phase III trial by the AGO Breast Cancer Study Group. <i>Breast Cancer Research and Treatment</i> , 2013, 139, 779-787.	1.1	21
164	Tumor-infiltrating lymphocytes in breast cancer. <i>Oncoimmunology</i> , 2014, 3, e27926.	2.1	21
165	Human leucocyte antigen class I in hormone receptor-positive, HER2-negative breast cancer: association with response and survival after neoadjuvant chemotherapy. <i>Breast Cancer Research</i> , 2019, 21, 142.	2.2	21
166	Efficacy and safety of nab-paclitaxel 125Âmg/m ² and nab-paclitaxel 150Âmg/m ² compared to paclitaxel in early high-risk breast cancer. Results from the neoadjuvant randomized GeparSepto study (GBG 69). <i>Breast Cancer Research and Treatment</i> , 2017, 163, 495-506.	1.1	20
167	AGO Recommendations for the Diagnosis and Treatment of Patients with Locally Advanced and Metastatic Breast Cancer: Update 2021. <i>Breast Care</i> , 2021, 16, 228-235.	0.8	20
168	Anti-epidermal growth factor receptor-antibody therapy for treatment of breast cancer. <i>International Journal of Cancer</i> , 2002, 101, 390-394.	2.3	19
169	Update on neoadjuvant/preoperative therapy of breast cancer. <i>Current Opinion in Obstetrics and Gynecology</i> , 2013, 25, 66-73.	0.9	19
170	Predictive Factors for Response to Neoadjuvant Therapy in Breast Cancer. <i>Oncology Research and Treatment</i> , 2014, 37, 5-5.	0.8	19
171	Phase III study on efficacy of taxanes plus bevacizumab with or without capecitabine as first-line chemotherapy in metastatic breast cancer. <i>Breast Cancer Research and Treatment</i> , 2015, 149, 141-149.	1.1	19
172	Endocrine Therapy in Premenopausal Hormone Receptor Positive/Human Epidermal Growth Receptor 2 Negative Metastatic Breast Cancer: Between Guidelines and Literature. <i>Oncologist</i> , 2018, 23, 974-981.	1.9	19
173	Optimal Systemic Treatment for Early Triple-Negative Breast Cancer. <i>Breast Care</i> , 2020, 15, 217-226.	0.8	19
174	AGO Recommendations for the Surgical Therapy of the Axilla After Neoadjuvant Chemotherapy: 2021 Update. <i>Geburtshilfe Und Frauenheilkunde</i> , 2021, 81, 1112-1120.	0.8	17
175	Prospective, Multicenter, Randomized Phase III Trial Evaluating the Impact of Lymphoscintigraphy as Part of Sentinel Node Biopsy in Early Breast Cancer: SenSzi (GBG80) Trial. <i>Journal of Clinical Oncology</i> , 2019, 37, 1490-1498.	0.8	16
176	Predictive Markers of Response to Neoadjuvant Durvalumab with Nab-Paclitaxel and Dose-Dense Doxorubicin/Cyclophosphamide in Basal-Like Triple-Negative Breast Cancer. <i>Clinical Cancer Research</i> , 2022, 28, 2587-2597.	3.2	16
177	Re-Challenging Taxanes in Recurrent Breast Cancer in Patients Treated with (Neo-)Adjuvant Taxane-Based Therapy. <i>Breast Care</i> , 2011, 6, 279-283.	0.8	14
178	Reporting of myelotoxicity associated with emerging regimens for the treatment of selected solid tumors. <i>Critical Reviews in Oncology/Hematology</i> , 2012, 81, 136-150.	2.0	14
179	The Importance of Supportive Care in Breast Cancer Patients. <i>Breast Care</i> , 2014, 9, 230-231.	0.8	14
180	Neoadjuvant treatment of breast cancer. <i>Current Opinion in Obstetrics and Gynecology</i> , 2015, 27, 85-91.	0.9	14

#	ARTICLE	IF	CITATIONS
181	Efficacy and safety of tailored and dose-dense adjuvant chemotherapy and trastuzumab for resected HER2-positive breast cancer: Results from the phase 3 PANTHER trial. <i>Cancer</i> , 2020, 126, 1175-1182.	2.0	14
182	Patient-reported outcomes from KATHERINE: A phase 3 study of adjuvant trastuzumab emtansine versus trastuzumab in patients with residual invasive disease after neoadjuvant therapy for human epidermal growth factor receptor 2-positive breast cancer. <i>Cancer</i> , 2020, 126, 3132-3139.	2.0	14
183	Chemotherapy-induced ovarian failure in young women with early breast cancer: Prospective analysis of four randomised neoadjuvant/adjuvant breast cancer trials. <i>European Journal of Cancer</i> , 2021, 152, 193-203.	1.3	14
184	Integrative proteomic and gene expression analysis identify potential biomarkers for adjuvant trastuzumab resistance: analysis from the Fin-her phase III randomized trial. <i>Oncotarget</i> , 2015, 6, 30306-30316.	0.8	14
185	Use of goserelin in the treatment of breast cancer. <i>Expert Review of Anticancer Therapy</i> , 2005, 5, 591-604.	1.1	13
186	Paclitaxel in patients with advanced solid tumors and hepatic dysfunction: a pilot study. <i>Expert Opinion on Drug Safety</i> , 2010, 9, 515-523.	1.0	13
187	Predictive value of sphingosine kinase 1 expression in neoadjuvant treatment of breast cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013, 139, 1681-1689.	1.2	13
188	Randomised, open-label, phase II study comparing the efficacy and the safety of cabazitaxel versus weekly paclitaxel given as neoadjuvant treatment in patients with operable triple-negative or luminal B/HER2-negative breast cancer (GENEVIEVE). <i>European Journal of Cancer</i> , 2017, 84, 1-8.	1.3	13
189	Reactive stroma and trastuzumab resistance in HER2-positive early breast cancer. <i>International Journal of Cancer</i> , 2020, 147, 266-276.	2.3	13
190	Effects of capecitabine as part of neo-/adjuvant chemotherapy – A meta-analysis of individual breast cancer patient data from 13 randomised trials including 15,993 patients. <i>European Journal of Cancer</i> , 2022, 166, 185-201.	1.3	13
191	Outcome of breast cancer patients treated with chemotherapy during pregnancy compared with non-pregnant controls. <i>European Journal of Cancer</i> , 2022, 170, 54-63.	1.3	13
192	Oxaliplatin and 5-fluorouracil for heavily pretreated metastatic breast cancer: a preliminary phase II study. <i>Anti-Cancer Drugs</i> , 2003, 14, 549-553.	0.7	12
193	Prediction of Response to Neoadjuvant Chemotherapy: New Biomarker Approaches and Concepts. <i>Breast Care</i> , 2011, 6, 265-272.	0.8	12
194	Adverse Event Management of Oral Mucositis in Patients with Breast Cancer. <i>Breast Care</i> , 2014, 9, 232-237.	0.8	12
195	Evaluation of soluble carbonic anhydrase IX as predictive marker for efficacy of bevacizumab: A biomarker analysis from the geparquinto phase III neoadjuvant breast cancer trial. <i>International Journal of Cancer</i> , 2019, 145, 857-868.	2.3	12
196	Androgen receptor expression and response to chemotherapy in breast cancer patients treated in the neoadjuvant TECHNO and PREPARE trial. <i>British Journal of Cancer</i> , 2019, 121, 1009-1015.	2.9	12
197	Gene Expression-Based Prediction of Neoadjuvant Chemotherapy Response in Early Breast Cancer: Results of the Prospective Multicenter EXPRESSION Trial. <i>Clinical Cancer Research</i> , 2021, 27, 2148-2158.	3.2	12
198	Survival analysis of the randomised phase III GeparOcto trial comparing neoadjuvant chemotherapy of intense dose-dense epirubicin, paclitaxel, cyclophosphamide versus weekly paclitaxel, liposomal doxorubicin (plus carboplatin in triple-negative breast cancer) for patients with high-risk early breast cancer. <i>European Journal of Cancer</i> , 2022, 160, 100-111.	1.3	12

#	ARTICLE	IF	CITATIONS
199	Six-year absolute invasive disease-free survival benefit of adding adjuvant pertuzumab to trastuzumab and chemotherapy for patients with early HER2-positive breast cancer: A Subpopulation Treatment Effect Pattern Plot (STEPP) analysis of the APHINITY (BIG 4-11) trial. <i>European Journal of Cancer</i> , 2022, 166, 219-228.	1.3	12
200	Being Pregnant and Diagnosed with Breast Cancer. <i>Breast Care</i> , 2012, 7, 204-209.	0.8	11
201	Risk Assessment after Neoadjuvant Chemotherapy in Luminal Breast Cancer Using a Clinicomolecular Predictor. <i>Clinical Cancer Research</i> , 2018, 24, 3358-3365.	3.2	11
202	Dual Blockade of HER-2 Provides a Greater Magnitude of Benefit in Patients With Hormone-Negative Versus Hormone-Positive Breast Cancer. <i>Clinical Breast Cancer</i> , 2016, 16, 444-455.	1.1	10
203	Clinical and analytical validation of Ki-67 in 9069 patients from IBCSG VIII, BIG1-98 and GeparTrio trial: systematic modulation of interobserver variance in a comprehensive in silico ring trial. <i>Breast Cancer Research and Treatment</i> , 2019, 176, 557-568.	1.1	10
204	Neutropenic complications in the PANTHER phase III study of adjuvant tailored dose-dense chemotherapy in early breast cancer. <i>Acta Oncologica</i> , 2020, 59, 75-81.	0.8	10
205	Phase III postneoadjuvant study evaluating sacituzumab govitecan, an antibody drug conjugate in primary HER2-negative breast cancer patients with high relapse risk after standard neoadjuvant treatment: SASCIA. <i>Journal of Clinical Oncology</i> , 2021, 39, TPS602-TPS602.	0.8	10
206	Integrating CDK4/6 inhibitors in the treatment of patients with early breast cancer. <i>Breast</i> , 2022, 62, S70-S79.	0.9	10
207	Breakthroughs in research and treatment of early breast cancer: an overview of the last three decades. <i>Archives of Gynecology and Obstetrics</i> , 2013, 288, 1203-1212.	0.8	9
208	Sorafenib in the Treatment of Early Breast Cancer: Results of the Neoadjuvant Phase II Study - SOFIA*. <i>Breast Care</i> , 2014, 9, 169-174.	0.8	9
209	Neoadjuvant chemotherapy for early breast cancer. <i>Lancet Oncology</i> , The, 2018, 19, e129.	5.1	9
210	A Small Hypoxia Signature Predicted pCR Response to Bevacizumab in the Neoadjuvant GeparQuinto Breast Cancer Trial. <i>Clinical Cancer Research</i> , 2020, 26, 1896-1904.	3.2	9
211	A plain language summary of the ASCENT study: Sacituzumab Govitecan for metastatic triple-negative breast cancer. <i>Future Oncology</i> , 2021, 17, 3911-3924.	1.1	9
212	Bendamustine in Metastatic Breast Cancer: An Old Drug in New Design. <i>Breast Care</i> , 2008, 3, 333-339.	0.8	8
213	Model-based optimization of G-CSF treatment during cytotoxic chemotherapy. <i>Journal of Cancer Research and Clinical Oncology</i> , 2018, 144, 343-358.	1.2	8
214	MGMT promoter methylation in triple negative breast cancer of the GeparSixto trial. <i>PLoS ONE</i> , 2020, 15, e0238021.	1.1	8
215	HER2-low-positive breast cancer from four neoadjuvant clinical trials – Authors' reply. <i>Lancet Oncology</i> , The, 2021, 22, e427.	5.1	8
216	Integrating Immunotherapy Into the Treatment Landscape for Patients With Triple-Negative Breast Cancer. <i>American Society of Clinical Oncology Educational Book / ASCO American Society of Clinical Oncology Meeting</i> , 2022, , 47-59.	1.8	8

#	ARTICLE	IF	CITATIONS
217	pCR rates in patients with bilateral breast cancer after neoadjuvant anthracycline-taxane based-chemotherapy – A retrospective pooled analysis of individual patients data of four German neoadjuvant trials. <i>Breast</i> , 2017, 32, 73-78.	0.9	7
218	The rise of oncology biosimilars: from process to promise. <i>Future Oncology</i> , 2019, 15, 3255-3265.	1.1	7
219	CDK4/6 inhibitors in breast cancer: one more step towards reduced mortality. <i>Lancet Oncology</i> , The, 2020, 21, 191-192.	5.1	7
220	Breast cancer diagnosed in the post-weaning period is indicative for a poor outcome. <i>European Journal of Cancer</i> , 2021, 155, 13-24.	1.3	7
221	The GISS Trial: a Phase II Prevention Trial of Screening Plus Goserelin, Ibandronate, versus Screening Alone in Premenopausal Women at Increased Risk of Breast Cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2011, 20, 2141-2149.	1.1	6
222	A clinical calculator to predict disease outcomes in women with hormone receptor-positive advanced breast cancer treated with first-line endocrine therapy. <i>Breast Cancer Research and Treatment</i> , 2021, 189, 15-23.	1.1	6
223	AGO Algorithms for the Treatment of Breast Cancer: Update 2021. <i>Geburtshilfe Und Frauenheilkunde</i> , 2021, 81, 1101-1111.	0.8	6
224	<i>mdm2</i> gene amplification is associated with luminal breast cancer progression in humanized <i>PDX</i> mice and a worse outcome of estrogen receptor positive disease. <i>International Journal of Cancer</i> , 2022, 150, 1357-1372.	2.3	6
225	Phase I dose finding study evaluating the combination of bendamustine with weekly paclitaxel in patients with pre-treated metastatic breast cancer: RiTa trial. <i>Cancer Chemotherapy and Pharmacology</i> , 2009, 63, 953-958.	1.1	5
226	Early assessment with magnetic resonance imaging for prediction of pathologic response to neoadjuvant chemotherapy in triple-negative breast cancer: Results from the phase III BrighTNess trial. <i>European Journal of Surgical Oncology</i> , 2020, 46, 223-228.	0.5	5
227	Genomic Aberrations and Late Recurrence in Postmenopausal Women with Hormone Receptor-positive Early Breast Cancer: Results from the SOLE Trial. <i>Clinical Cancer Research</i> , 2021, 27, 504-512.	3.2	5
228	Treatment of Patients with Early Breast Cancer: Evidence, Controversies, Consensus. <i>Geburtshilfe Und Frauenheilkunde</i> , 2021, 81, 637-653.	0.8	5
229	Immunophenotype and proliferation to predict for response to neoadjuvant chemotherapy in TNBC: Results from BrighTNess phase III study.. <i>Journal of Clinical Oncology</i> , 2019, 37, 510-510.	0.8	5
230	EGF-R and Her2/neu overexpressing tumors: independent collectives for treatment of breast cancer by specific monoclonal antibody-therapy. <i>Journal of Cancer Research and Clinical Oncology</i> , 2003, 129, 250-251.	1.2	4
231	Darbepoetin Alfa as Primary Prophylaxis of Anemia in Patients with Breast Cancer Treated Preoperatively with Docetaxel/Doxorubicin/Cyclophosphamide. <i>Supportive Cancer Therapy</i> , 2006, 3, 103-109.	0.3	4
232	Neoadjuvant Clinical Trials for the Treatment of Primary Breast Cancer: The Experience of the German Study Groups. <i>Current Oncology Reports</i> , 2012, 14, 27-34.	1.8	4
233	Evolution of adjuvant chemotherapy for breast cancer. <i>Lancet</i> , The, 2015, 385, 1812-1814.	6.3	4
234	Phase III randomised trial comparing intense dose-dense chemotherapy to tailored dose-dense chemotherapy in high-risk early breast cancer (GAIN-2). <i>European Journal of Cancer</i> , 2021, 156, 138-148.	1.3	4

#	ARTICLE	IF	CITATIONS
235	Response-based molecular subtypingâ€”emergence of the third generation of breast cancer subtypes. <i>Cancer Cell</i> , 2022, 40, 592-594.	7.7	4
236	Reply to A. Morabito et al and G. Valabrega et al. <i>Journal of Clinical Oncology</i> , 2009, 27, e124-e125.	0.8	3
237	What Can Be Learned From Trials Running Short of Patients or Events?. <i>Journal of Clinical Oncology</i> , 2012, 30, 901-903.	0.8	3
238	Present Status of Adjuvant Chemotherapy for Elderly Breast Cancer Patients. <i>Breast Care</i> , 2012, 7, 439-444.	0.8	3
239	Validation of a Nomogram Predicting Non-Sentinel Lymph Node Metastases among Patients with Breast Cancer after Primary Systemic Therapy - a transSENTINA Substudy. <i>Breast Care</i> , 2018, 13, 440-446.	0.8	3
240	A randomized phase II study to determine the efficacy and tolerability of two doses of eribulin plus lapatinib in trastuzumab-pretreated patients with HER-2-positive metastatic breast cancer (E-VITA). <i>Anti-Cancer Drugs</i> , 2019, 30, 394-401.	0.7	3
241	Residual Axillary Burden After Neoadjuvant Chemotherapy (NACT) in Early Breast Cancer in Patients with a priori Clinically Occult Nodal Metastases â€” a transSENTINA Analysis. <i>Geburtshilfe Und Frauenheilkunde</i> , 2020, 80, 1229-1236.	0.8	3
242	Neo-/adjuvant phase III trial to compare intense dose-dense (idd) treatment with EnPC to tailored dose-dense (dt) therapy with dtEC-dtD for patients with high-risk early breast cancer: Results on pathological complete response (pCR) for patients treated within the neoadjuvant setting.. <i>Journal of Clinical Oncology</i> , 2018, 36, 568-568.	0.8	3
243	Impact of Age on Clinical Outcomes and Efficacy of Adjuvant Dual Anti-HER2 Targeted Therapy. <i>Journal of the National Cancer Institute</i> , 2022, 114, 1117-1126.	3.0	3
244	Reply to L. Del Mastro and Z. Blumenfeld. <i>Journal of Clinical Oncology</i> , 2011, 29, 3341-3342.	0.8	2
245	Multicenter Phase II Study with Weekly Bendamustine and Paclitaxel as First- or Later-Line Therapy in Patients with Metastatic Breast Cancer: RiTa II Trial. <i>Breast Care</i> , 2011, 6, 457-461.	0.8	2
246	Targeting the Immune System in Breast Cancer: Hype or Hope?: TILs and Newer Immune-Based Therapies Being Evaluated for HER2+ and TNBC. <i>Current Breast Cancer Reports</i> , 2015, 7, 203-209.	0.5	2
247	Prognostic impact of HER3 based on protein and mRNA expression in high-grade serous ovarian carcinoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2017, 470, 143-151.	1.4	2
248	Impact of Nuclear Oestrogen Receptor Beta Expression in Breast Cancer Patients Undergoing Neoadjuvant Chemotherapy. <i>Geburtshilfe Und Frauenheilkunde</i> , 2019, 79, 1110-1117.	0.8	2
249	Autologous Lipotransfer - Daily Therapeutic Practice in Breast Cancer: An Intergroup Analysis Encompassing NOGGO, WSG, GBC, AWO Gyn and DGPRÄ„C. <i>Breast Care</i> , 2019, 14, 165-169.	0.8	2
250	Subgroup of post-neoadjuvant luminal-B tumors assessed by HTG in PENELOPE-B investigating palbociclib in high risk HER2-/HR+ breast cancer with residual disease.. <i>Journal of Clinical Oncology</i> , 2021, 39, 519-519.	0.8	2
251	Reply to Y. Kawamura et al. <i>Journal of Clinical Oncology</i> , 2021, 39, 3090-3091.	0.8	2
252	Data describing the poor outcome associated with a breast cancer diagnosis in the post-weaning period. <i>Data in Brief</i> , 2021, 38, 107354.	0.5	2

#	ARTICLE	IF	CITATIONS
253	Prevention Trials in Women at Moderate Risk of Breast Cancer. <i>Breast Care</i> , 2006, 1, 298-303.	0.8	1
254	Reply to F. Bellati et al. <i>Journal of Clinical Oncology</i> , 2010, 28, e473-e473.	0.8	1
255	Opinions on the ASCO 2011 Annual Meeting. <i>Breast Care</i> , 2011, 6, 315-319.	0.8	1
256	Concerning Dediu M, Zielinski A: A Proposal to Redefine Pathologic Complete Remission as Endpoint following Neoadjuvant Chemotherapy in Early Breast Cancer. <i>Breast Care</i> 2019; Doi 10.1159/000500620. <i>Breast Care</i> , 2020, 15, 96-101.	0.8	1
257	Reporting the Analytical Method Is Essential to Assessing Studies in Which Biomarkers Are a Major Study Objective—Reply. <i>JAMA Oncology</i> , 2021, 7, 1403.	3.4	1
258	The Dual Role of Tumor-Infiltrating Lymphocytes in Human Epidermal Growth Factor Receptor 2—Positive Primary Breast Cancer. <i>JAMA Oncology</i> , 2015, 1, 455.	3.4	1
259	Cyclin dependent kinase 4/6 inhibitors in early breast cancer: what is the role of Ki-67?. <i>Lancet Oncology</i> , The, 2022, 23, 325-328.	5.1	1
260	Bericht vom 5. GBG-Jahrestreffen 07.–08.03.2008. <i>Breast Care</i> , 2008, 3, 139-143.	0.8	0
261	Bericht vom 6. GBG-Jahrestreffen 19.–20.02.2009. <i>Breast Care</i> , 2009, 4, 130-134.	0.8	0
262	Can We Keep the “PROMISE”™?. <i>Breast Care</i> , 2011, 6, 467-470.	0.8	0
263	News from the San Antonio Breast Cancer Symposium 2010. <i>Breast Care</i> , 2011, 6, 56-61.	0.8	0
264	Incorporating Agents that Target HER2 in the Neoadjuvant Setting. <i>Current Breast Cancer Reports</i> , 2011, 3, 190-196.	0.5	0
265	Breast Cancer in Young Women - News from the BCY3 Consensus Conference. <i>Breast Care</i> , 2016, 11, 432-435.	0.8	0
266	Efficacy + safety of palbociclib (P) in patients (pts) ≥50 y with hormone receptor-positive (HR+)/human epidermal growth factor receptor 2-negative (HER2-negative) advanced breast cancer (ABC): subgroup analysis of 2 randomized phase 3 studies. <i>Breast</i> , 2018, 41, S11-S12.	0.9	0
267	Breast Cancer in Young and Premenopausal Women. <i>Annals of Oncology</i> , 2019, 30, vi67.	0.6	0
268	Post-Neoadjuvant Therapy. <i>Breast Care</i> , 2019, 14, 409-413.	0.8	0
269	A PLAIN-LANGUAGE SUMMARY OF THE SOLAR-1 TRIAL: STUDYING ALPELISIB WITH FULVESTRANT IN PATIENTS WITH HR+, HER2-negative ADVANCED BREAST CANCER WHO HAD PREVIOUSLY RECEIVED AN AROMATASE INHIBITOR. <i>Breast</i> , 2019, 48, S48.	0.9	0
270	Corrigendum to “Efficacy + safety of palbociclib (P) in patients (pts) ≥50 y with hormone receptor-positive (HR+)/human epidermal growth factor receptor 2-negative (HER2-negative) advanced breast cancer (ABC): Subgroup analysis of 2 randomized phase 3 studies” [Breast 41S1 (2018) S11-S12]. <i>Breast</i> , 2020, 49, 131.	0.9	0

#	ARTICLE	IF	CITATIONS
271	THE ALPELISIB (ALP) EXPERIENCE IN THE SOLAR-1 AND BYLIEVE STUDIES: PERSPECTIVES FOR PRACTITIONERS CARING FOR PATIENTS (PTS) WITH HORMONE RECEPTOR-POSITIVE (HR+), HUMAN EPIDERMAL GROWTH FACTOR RECEPTOR 2-NEGATIVE (HER2 ⁺) ADVANCED BREAST CANCER (ABC). <i>Breast</i> , 2021, 59, S49.	0.9	0