Gianni Mazzoni

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7146266/publications.pdf Version: 2024-02-01



CIANNI ΜΑΖΖΟΝΙ

#	Article	IF	CITATIONS
1	Promotion and maintenance of physically active lifestyle in older outpatients 2 years after acute coronary syndrome. Aging Clinical and Experimental Research, 2022, 34, 1065-1072.	2.9	3
2	Maintenance of Exercise Habit and Its Impact on Cardiovascular Risk Factors in Previously Sedentary People: A 7-Year Follow-Up. Journal of Aging and Physical Activity, 2022, , 1-9.	1.0	0
3	Guided walking reduces blood pressure in hypertensive sedentary subjects including those with resistant hypertension. Journal of Human Hypertension, 2021, 35, 226-231.	2.2	6
4	Prognostic comparison of the FRIEND and Wasserman/Hansen peak VO2 equations applied to a submaximal walking test in outpatients with cardiovascular disease. European Journal of Preventive Cardiology, 2021, 28, 287-292.	1.8	12
5	The impact of periprocedural myocardial infarction on mortality in older adults with non-ST-segment elevation acute coronary syndrome. Journal of Cardiovascular Medicine, 2021, Publish Ahead of Print, 546-552.	1.5	1
6	Contrast Associated Acute Kidney Injury and Mortality in Older Adults with Acute Coronary Syndrome: A Pooled Analysis of the FRASER and HULK Studies. Journal of Clinical Medicine, 2021, 10, 2151.	2.4	8
7	Relationship between exercise capacity and grip strength in a cohort of older cardiac outpatients. Journal of Sports Medicine and Physical Fitness, 2021, , .	0.7	0
8	Physical fitness changes induced by thermal aquatic standardized exercise in chronic venous disease patients. Phlebology, 2021, , 026835552110519.	1.2	1
9	Peak Oxygen Uptake Estimation From A Moderate 1-KM Treadmill Walk in Women With Cardiovascular Disease. Journal of Cardiopulmonary Rehabilitation and Prevention, 2021, 41, 432-434.	2.1	4
10	Reply to commentary on: Prediction of VO2peak by an endurance test and prognostic value of the FRIEND equation. European Journal of Preventive Cardiology, 2020, 27, 2054-2056.	1.8	0
11	Improving walking speed reduces hospitalization costs in outpatients with cardiovascular disease. An analysis based on a multistrata non-parametric test. BMC Health Services Research, 2020, 20, 1048.	2.2	4
12	Impact of Improvement in Walking Speed on Hospitalization and Mortality in Females with Cardiovascular Disease. Journal of Clinical Medicine, 2020, 9, 1755.	2.4	6
13	Exercise intervention improves quality of life in older adults after myocardial infarction: randomised clinical trial. Heart, 2020, 106, 1658-1664.	2.9	37
14	Guided walking is more effective than suggested walking in reducing the blood pressure of hypertensive sedentary subjects and in modifying their lifestyle. Sport Sciences for Health, 2020, 16, 375-381.	1.3	1
15	A moderate 200-m walk test estimates peak oxygen uptake in elderly outpatients with cardiovascular disease. Journal of Sports Medicine and Physical Fitness, 2020, 60, 786-793.	0.7	6
16	Moderate walking speed predicts hospitalisation in hypertensive patients with cardiovascular disease. European Journal of Preventive Cardiology, 2018, 25, 1558-1560.	1.8	2
17	Determining the best percent-predicted equation for estimated VO 2 peak by a 1-km moderate perceptually-regulated treadmill walk to predict mortality in outpatients with cardiovascular disease. Journal of Science and Medicine in Sport, 2018, 21, 307-311.	1.3	13
18	Reduction in risk factors for cardiovascular diseases and long-lasting walking habit in sedentary male and female subjects following 1Âyear of guided walking. Sport Sciences for Health, 2018, 14, 121-126.	1.3	8

GIANNI MAZZONI

#	Article	IF	CITATIONS
19	500-meter and 1000-meter moderate walks equally assess cardiorespiratory fitness in male outpatients with cardiovascular diseases. Journal of Sports Medicine and Physical Fitness, 2018, 58, 1312-1317.	0.7	13
20	Physical activity intervention for elderly patients with reduced physical performance after acute coronary syndrome (HULK study): rationale and design of a randomized clinical trial. BMC Cardiovascular Disorders, 2018, 18, 98.	1.7	22
21	Peak oxygen uptake estimation from a moderate 500-m treadmill walk in older women with cardiovascular disease. Sport Sciences for Health, 2018, 14, 655-661.	1.3	1
22	A moderate 500-m treadmill walk for estimating peak oxygen uptake in men with NYHA class I-II heart failure and reduced left ventricular ejection fraction. BMC Cardiovascular Disorders, 2018, 18, 67.	1.7	12
23	A moderate 1-km treadmill walk predicts mortality in men with mid-range left ventricular dysfunction. European Journal of Preventive Cardiology, 2017, 24, 1670-1672.	1.8	12
24	Outdoor Reproducibility of a 1-km Treadmill Walking Test to Predict Peak Oxygen Uptake in Cardiac Patients. Journal of Cardiopulmonary Rehabilitation and Prevention, 2017, 37, 347-349.	2.1	19
25	Oxygen Uptake Attenuation at Ventilatory Threshold in Men With Coronary Artery Disease. Journal of Cardiopulmonary Rehabilitation and Prevention, 2016, 36, 258-262.	2.1	2
26	Improved walking speed is associated with lower hospitalisation rates in patients in an exercise-based secondary prevention programme. Heart, 2016, 102, 1902-1908.	2.9	36
27	Association between VO2 peak estimated by a 1-km treadmill walk and mortality. A 10-year follow-up study in patients with cardiovascular disease. International Journal of Cardiology, 2014, 173, 248-252.	1.7	33
28	Body image and weight perceptions in relation to actual measurements by means of a new index and level of physical activity in Italian university students. Journal of Translational Medicine, 2014, 12, 42.	4.4	80
29	Treadmill walking speed and survival prediction in men with cardiovascular disease: a 10-year follow-up study. BMJ Open, 2013, 3, e003446.	1.9	31
30	Peak Oxygen Uptake Prediction From a Moderate, Perceptually Regulated, 1-km Treadmill Walk in Male Cardiac Patients. Journal of Cardiopulmonary Rehabilitation and Prevention, 2012, 32, 262-269.	2.1	40
31	Incremental Exercise Using Progressive Versus Constant Pedaling Rates. Journal of Cardiopulmonary Rehabilitation and Prevention, 2011, 31, 303-307.	2.1	1
32	Home-centred physical fitness programme in morbidly obese individuals: a randomized controlled trial. Clinical Rehabilitation, 2008, 22, 940-950.	2.2	17
33	Identification of a JOURNAL/jscr/04.03/00124278-200807000-00013/OV0312_4/v/2021-02-09T093601Z/r/image-png o2 Deflection Point Coinciding With the Heart Rate Deflection Point and Ventilatory Threshold in Cycling. Journal of Strength and Conditioning Research. 2008. 22. 1116-1123.	¹ 2.1	17
34	Influence of low-density lipoprotein (LDL) receptor-related protein and ABO blood group genotypes on factor XI levels. Thrombosis and Haemostasis, 2008, 99, 789-790.	3.4	3
35	On the Methodology of the Conconi Test. International Journal of Sports Medicine, 2005, 26, 397-398.	1.7	1
36	An Incremental Test to Identify the Pain Threshold Speed in Patients With Intermittent Claudication Circulation Journal, 2002, 66, 1124-1127.	1.6	10

GIANNI MAZZONI

#	Article	IF	CITATIONS
37	Increased frequency of the homozygous II ACE genotype in Italian Olympic endurance athletes. European Journal of Human Genetics, 2002, 10, 576-577.	2.8	46
38	The power output/heart rate relationship in cycling: test standardization and repeatability. Medicine and Science in Sports and Exercise, 1999, 31, 1478.	0.4	27
39	The Conconi Test: Methodology After 12 Years of Application. International Journal of Sports Medicine, 1996, 17, 509-519.	1.7	113
40	Reproducibility of the Conconi Test: Test Repeatability and Observer Variations. International Journal of Sports Medicine, 1996, 17, 520-524.	1.7	21
41	Hematological Indices of Erythropoietin Administration in Athletes. International Journal of Sports Medicine, 1993, 14, 307-311.	1.7	56
42	Platelet count, mean platelet volume, and platelet dimensional width in professional cyclists during races. Thrombosis Research, 1991, 62, 791-792.	1.7	12