

Vincent Walter Bloks

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/714638/publications.pdf>

Version: 2024-02-01

94
papers

7,479
citations

94381

37
h-index

53190

85
g-index

96
all docs

96
docs citations

96
times ranked

10593
citing authors

#	ARTICLE	IF	CITATIONS
1	Transfer of Intestinal Microbiota From Lean Donors Increases Insulin Sensitivity in Individuals With Metabolic Syndrome. <i>Gastroenterology</i> , 2012, 143, 913-916.e7.	0.6	2,287
2	Stimulation of Lipogenesis by Pharmacological Activation of the Liver X Receptor Leads to Production of Large, Triglyceride-rich Very Low Density Lipoprotein Particles. <i>Journal of Biological Chemistry</i> , 2002, 277, 34182-34190.	1.6	420
3	Beyond intestinal soapâ€”bile acids in metabolic control. <i>Nature Reviews Endocrinology</i> , 2014, 10, 488-498.	4.3	354
4	24(S)-Hydroxycholesterol Participates in a Liver X Receptor-controlled Pathway in Astrocytes That Regulates Apolipoprotein E-mediated Cholesterol Efflux. <i>Journal of Biological Chemistry</i> , 2006, 281, 12799-12808.	1.6	204
5	In Utero Undernutrition in Male Mice Programs Liver Lipid Metabolism in the Second-Generation Offspring Involving Altered Lxra DNA Methylation. <i>Cell Metabolism</i> , 2014, 19, 941-951.	7.2	178
6	Regulation of Bile Acid Synthesis by the Nuclear Receptor Rev-erb β . <i>Gastroenterology</i> , 2008, 135, 689-698.e5.	0.6	175
7	Increased Hepatobiliary and Fecal Cholesterol Excretion upon Activation of the Liver X Receptor Is Independent of ABCA1. <i>Journal of Biological Chemistry</i> , 2002, 277, 33870-33877.	1.6	174
8	Improved glycemic control with colessevelam treatment in patients with type 2 diabetes is not directly associated with changes in bile acid metabolism. <i>Hepatology</i> , 2010, 52, 1455-1464.	3.6	163
9	Impaired secretion of very low density lipoprotein-triglycerides by apolipoprotein E- deficient mouse hepatocytes.. <i>Journal of Clinical Investigation</i> , 1997, 100, 2915-2922.	3.9	154
10	Peroxisome proliferator-activated receptor alpha (PPARalpha)-mediated regulation of multidrug resistance 2 (Mdr2) expression and function in mice. <i>Biochemical Journal</i> , 2003, 369, 539-547.	1.7	150
11	Malnutrition-associated liver steatosis and ATP depletion is caused by peroxisomal and mitochondrial dysfunction. <i>Journal of Hepatology</i> , 2016, 65, 1198-1208.	1.8	133
12	The liver X-receptor gene promoter is hypermethylated in a mouse model of prenatal protein restriction. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2010, 298, R275-R282.	0.9	131
13	Sitosterolemia in ABC-Transporter G5-deficient mice is aggravated on activation of the liver-X receptor. <i>Gastroenterology</i> , 2004, 126, 290-300.	0.6	130
14	Hepatobiliary cholesterol transport is not impaired in Abca1-null mice lacking HDL. <i>Journal of Clinical Investigation</i> , 2001, 108, 843-850.	3.9	127
15	Transintestinal Cholesterol Transport Is Active in Mice and Humans and Controls Ezetimibe-Induced Fecal Neutral Sterol Excretion. <i>Cell Metabolism</i> , 2016, 24, 783-794.	7.2	119
16	Liver X receptor activation restores memory in aged AD mice without reducing amyloid. <i>Neurobiology of Aging</i> , 2011, 32, 1262-1272.	1.5	118
17	Apolipoprotein E Participates in the Regulation of Very Low Density Lipoprotein-Triglyceride Secretion by the Liver. <i>Journal of Biological Chemistry</i> , 1999, 274, 35711-35718.	1.6	115
18	Intestinal Farnesoid X Receptor Controls Transintestinal Cholesterol Excretion in Mice. <i>Gastroenterology</i> , 2017, 152, 1126-1138.e6.	0.6	109

#	ARTICLE	IF	CITATIONS
19	Statins increase hepatic cholesterol synthesis and stimulate fecal cholesterol elimination in mice. <i>Journal of Lipid Research</i> , 2016, 57, 1455-1464.	2.0	102
20	Hypoxia and Complement-and-Coagulation Pathways in the Deceased Organ Donor as the Major Target for Intervention to Improve Renal Allograft Outcome. <i>Transplantation</i> , 2015, 99, 1293-1300.	0.5	99
21	A human-like bile acid pool induced by deletion of hepatic Cyp2c70 modulates effects of FXR activation in mice. <i>Journal of Lipid Research</i> , 2020, 61, 291-305.	2.0	93
22	Biliary fibrosis associated with altered bile composition in a mouse model of erythropoietic protoporphyria. <i>Gastroenterology</i> , 1999, 117, 696-705.	0.6	91
23	Reduction of Cholesterol Absorption by Dietary Plant Sterols and Stanols in Mice Is Independent of the Abcg5/8 Transporter. <i>Journal of Nutrition</i> , 2006, 136, 2135-2140.	1.3	80
24	Maternal western diet primes non-alcoholic fatty liver disease in adult mouse offspring. <i>Acta Physiologica</i> , 2014, 210, 215-227.	1.8	80
25	Induction of hepatic ABC transporter expression is part of the PPAR α -mediated fasting response in the mouse. <i>Gastroenterology</i> , 2003, 124, 160-171.	0.6	79
26	Differential effects of streptozotocin-induced diabetes on expression of hepatic ABC-transporters in rats. <i>Gastroenterology</i> , 2002, 122, 1842-1852.	0.6	67
27	3-hydroxy-3-methylglutaryl-coenzyme a reductase inhibitors (statins) induce hepatic expression of the phospholipid translocase mdr2 in rats. <i>Gastroenterology</i> , 1999, 117, 678-687.	0.6	61
28	Down-regulation of hepatic and intestinal Abcg5 and Abcg8 expression associated with altered sterol fluxes in rats with streptozotocin-induced diabetes. <i>Diabetologia</i> , 2004, 47, 104-112.	2.9	61
29	A novel approach to monitor glucose metabolism using stable isotopically labelled glucose in longitudinal studies in mice. <i>Laboratory Animals</i> , 2013, 47, 79-88.	0.5	57
30	New insights in the multiple roles of bile acids and their signaling pathways in metabolic control. <i>Current Opinion in Lipidology</i> , 2018, 29, 194-202.	1.2	57
31	Genetic and Microbial Associations to Plasma and Fecal Bile Acids in Obesity Relate to Plasma Lipids and Liver Fat Content. <i>Cell Reports</i> , 2020, 33, 108212.	2.9	55
32	An Increased Flux through the Glucose 6-Phosphate Pool in Enterocytes Delays Glucose Absorption in Fxr $^{-/-}$ Mice. <i>Journal of Biological Chemistry</i> , 2009, 284, 10315-10323.	1.6	51
33	Liver receptor homolog 1 is critical for adequate up-regulation of Cyp7a1 gene transcription and bile salt synthesis during bile salt sequestration. <i>Hepatology</i> , 2011, 53, 2075-2085.	3.6	48
34	Abcg5/Abcg8-independent pathways contribute to hepatobiliary cholesterol secretion in mice. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 291, G414-G423.	1.6	47
35	Characterization of gut microbial structural variations as determinants of human bile acid metabolism. <i>Cell Host and Microbe</i> , 2021, 29, 1802-1814.e5.	5.1	43
36	Plant Sterols Cause Macrothrombocytopenia in a Mouse Model of Sitosterolemia. <i>Journal of Biological Chemistry</i> , 2008, 283, 6281-6287.	1.6	40

#	ARTICLE	IF	CITATIONS
37	Essential fatty acid deficiency in mice is associated with hepatic steatosis and secretion of large VLDL particles. <i>American Journal of Physiology - Renal Physiology</i> , 2005, 288, G1150-G1158.	1.6	38
38	Cerebral Accumulation of Dietary Derivable Plant Sterols does not Interfere with Memory and Anxiety Related Behavior in <i>Abcg5^{-/-}/Abg8^{-/-}</i> Mice. <i>Plant Foods for Human Nutrition</i> , 2011, 66, 149-156.	1.4	38
39	Clinical symptoms of right ventricular failure in experimental chronic pressure load are associated with progressive diastolic dysfunction. <i>Journal of Molecular and Cellular Cardiology</i> , 2015, 79, 244-253.	0.9	38
40	Hyperlipidemia and atherosclerosis associated with liver disease in ferrochelatase-deficient mice. <i>Journal of Lipid Research</i> , 2001, 42, 41-50.	2.0	37
41	Gene expression profiling in livers of mice after acute inhibition of β -oxidation. <i>Genomics</i> , 2007, 90, 680-689.	1.3	36
42	Disturbed hepatic carbohydrate management during high metabolic demand in medium-chain acyl-CoA dehydrogenase (MCAD)-deficient mice. <i>Hepatology</i> , 2008, 47, 1894-1904.	3.6	36
43	Cholesterol feeding strongly reduces hepatic VLDL-triglyceride production in mice lacking the liver X receptor β . <i>Journal of Lipid Research</i> , 2007, 48, 337-347.	2.0	35
44	Cross-talk between liver and intestine in control of cholesterol and energy homeostasis. <i>Molecular Aspects of Medicine</i> , 2014, 37, 77-88.	2.7	34
45	Hepatic Carbohydrate Response Element Binding Protein Activation Limits Nonalcoholic Fatty Liver Disease Development in a Mouse Model for Glycogen Storage Disease Type 1a. <i>Hepatology</i> , 2020, 72, 1638-1653.	3.6	34
46	<i>Lxrβ</i> Deficiency Hampers the Hepatic Adaptive Response to Fasting in Mice. <i>Journal of Biological Chemistry</i> , 2008, 283, 25437-25445.	1.6	33
47	Alterations in Brain Cholesterol Metabolism in the APPSLxPS1mut mouse, a Model for Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2010, 19, 117-127.	1.2	32
48	Intestinal PPAR β protects against diet-induced obesity, insulin resistance and dyslipidemia. <i>Scientific Reports</i> , 2017, 7, 846.	1.6	32
49	Cholangiopathy and Biliary Fibrosis in <i>Cyp2c70</i> -Deficient Mice Are Fully Reversed by Ursodeoxycholic Acid. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2021, 11, 1045-1069.	2.3	31
50	Hepatic Farnesoid X-Receptor Isoforms β 2 and β 4 Differentially Modulate Bile Salt and Lipoprotein Metabolism in Mice. <i>PLoS ONE</i> , 2014, 9, e115028.	1.1	30
51	Hepatic lipid accumulation, altered very low density lipoprotein formation and apolipoprotein E deposition in apolipoprotein E3-Leiden transgenic mice. <i>Journal of Hepatology</i> , 2000, 33, 189-198.	1.8	28
52	The hepatocyte IKK:NF- κ B axis promotes liver steatosis by stimulating de novo lipogenesis and cholesterol synthesis. <i>Molecular Metabolism</i> , 2021, 54, 101349.	3.0	28
53	Epigenome-wide association study of incident type 2 diabetes: a meta-analysis of five prospective European cohorts. <i>Diabetologia</i> , 2022, 65, 763-776.	2.9	28
54	Fat malabsorption in essential fatty acid-deficient mice is not due to impaired bile formation. <i>American Journal of Physiology - Renal Physiology</i> , 2002, 283, G900-G908.	1.6	25

#	ARTICLE	IF	CITATIONS
55	The effects of bariatric surgery on clinical profile, DNA methylation, and ageing in severely obese patients. <i>Clinical Epigenetics</i> , 2020, 12, 14.	1.8	23
56	MdrP-glycoproteins are not essential for biliary excretion of the hydrophobic heme precursor protoporphyrin in a griseofulvin-induced mouse model of erythropoietic protoporphyria. <i>Hepatology</i> , 2002, 35, 299-306.	3.6	21
57	Rosuvastatin Reduces Plasma Lipids by Inhibiting VLDL Production and Enhancing Hepatobiliary Lipid Excretion in ApoE*3-Leiden Mice. <i>Journal of Cardiovascular Pharmacology</i> , 2005, 45, 53-60.	0.8	21
58	Secretory phospholipase A2 increases SR-BI-mediated selective uptake from HDL but not biliary cholesterol secretion. <i>Journal of Lipid Research</i> , 2008, 49, 563-571.	2.0	21
59	Chronic Prednisolone Treatment Reduces Hepatic Insulin Sensitivity while Perturbing the Fed-to-Fasting Transition in Mice. <i>Endocrinology</i> , 2010, 151, 2171-2178.	1.4	21
60	Glucose-6-Phosphate Regulates Hepatic Bile Acid Synthesis in Mice. <i>Hepatology</i> , 2019, 70, 2171-2184.	3.6	21
61	An epigenome-wide association study identifies multiple DNA methylation markers of exposure to endocrine disruptors. <i>Environment International</i> , 2020, 144, 106016.	4.8	21
62	Bile acid sequestration normalizes plasma cholesterol and reduces atherosclerosis in hypercholesterolemic mice. No additional effect of physical activity. <i>Atherosclerosis</i> , 2013, 228, 117-123.	0.4	19
63	Male apoE*3-Leiden.CETP mice on high-fat high-cholesterol diet exhibit a biphasic dyslipidemic response, mimicking the changes in plasma lipids observed through life in men. <i>Physiological Reports</i> , 2017, 5, e13376.	0.7	19
64	Epigenetic programming at the <i>Mogat1</i> locus may link neonatal overnutrition with long-term hepatic steatosis and insulin resistance. <i>FASEB Journal</i> , 2018, 32, 6025-6037.	0.2	19
65	FXR overexpression alters adipose tissue architecture in mice and limits its storage capacity leading to metabolic derangements. <i>Journal of Lipid Research</i> , 2019, 60, 1547-1561.	2.0	19
66	Short-term protein restriction at advanced age stimulates FGF21 signalling, energy expenditure and browning of white adipose tissue. <i>FEBS Journal</i> , 2021, 288, 2257-2277.	2.2	18
67	The phosphatidylethanolamine N-methyltransferase pathway is quantitatively not essential for biliary phosphatidylcholine secretion. <i>Journal of Lipid Research</i> , 2007, 48, 2058-2064.	2.0	16
68	HSPA6 is an ulcerative colitis susceptibility factor that is induced by cigarette smoke and protects intestinal epithelial cells by stabilizing anti-apoptotic Bcl-XL. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2016, 1862, 788-796.	1.8	16
69	Potential of Intestine-Selective FXR Modulation for Treatment of Metabolic Disease. <i>Handbook of Experimental Pharmacology</i> , 2019, 256, 207-234.	0.9	16
70	Sex-Dependent Programming of Glucose and Fatty Acid Metabolism in Mouse Offspring by Maternal Protein Restriction. <i>Gender Medicine</i> , 2012, 9, 166-179.e13.	1.4	15
71	Hypertrophy induced KIF5B controls mitochondrial localization and function in neonatal rat cardiomyocytes. <i>Journal of Molecular and Cellular Cardiology</i> , 2016, 97, 70-81.	0.9	15
72	Postnatal Treatment With Dexamethasone Perturbs Hepatic and Cardiac Energy Metabolism and Is Associated With a Sustained Atherogenic Plasma Lipid Profile in Suckling Rats. <i>Pediatric Research</i> , 2007, 61, 165-170.	1.1	13

#	ARTICLE	IF	CITATIONS
73	Neonatal dexamethasone administration causes progressive renal damage due to induction of an early inflammatory response. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 294, F768-F776.	1.3	12
74	Potential of therapeutic bile acids in the treatment of neonatal Hyperbilirubinemia. <i>Scientific Reports</i> , 2021, 11, 11107.	1.6	12
75	Chronic Prednisolone Treatment Aggravates Hyperglycemia in Mice Fed a High-Fat Diet but Does Not Worsen Dietary Fat-Induced Insulin Resistance. <i>Endocrinology</i> , 2012, 153, 3713-3723.	1.4	11
76	The role of transhepatic bile salt flux in the control of hepatic secretion of triacylglycerol-rich lipoproteins in vivo in rodents. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2002, 1573, 9-20.	1.1	9
77	Metabolic responses to long-term pharmacological inhibition of CB1-receptor activity in mice in relation to dietary fat composition. <i>International Journal of Obesity</i> , 2010, 34, 374-384.	1.6	9
78	Low production of 12 α -hydroxylated bile acids prevents hepatic steatosis in Cyp2c70 $^{-/-}$ mice by reducing fat absorption. <i>Journal of Lipid Research</i> , 2021, 62, 100134.	2.0	9
79	An early-life diet containing large phospholipid-coated lipid globules programmes later-life postabsorptive lipid trafficking in high-fat diet- but not in low-fat diet-fed mice. <i>British Journal of Nutrition</i> , 2021, 125, 961-971.	1.2	8
80	Milk cholesterol concentration in mice is not affected by high cholesterol diet- or genetically-induced hypercholesterolaemia. <i>Scientific Reports</i> , 2018, 8, 8824.	1.6	7
81	Mice with a deficiency in Peroxisomal Membrane Protein 4 (PXMP4) display mild changes in hepatic lipid metabolism. <i>Scientific Reports</i> , 2022, 12, 2512.	1.6	7
82	Spontaneous liver disease in wild-type C57BL/6J OlaHsd mice fed semisynthetic diet. <i>PLoS ONE</i> , 2020, 15, e0232069.	1.1	6
83	Two time-point assessment of bile acid kinetics in humans using stable isotopes. <i>Isotopes in Environmental and Health Studies</i> , 2010, 46, 325-336.	0.5	3
84	Resistance to diet-induced adiposity in cannabinoid receptor-1 deficient mice is not due to impaired adipocyte function. <i>Nutrition and Metabolism</i> , 2011, 8, 93.	1.3	3
85	Transcriptome analysis suggests a compensatory role of the cofactors coenzyme A and NAD ⁺ in medium-chain acyl-CoA dehydrogenase knockout mice. <i>Scientific Reports</i> , 2019, 9, 14539.	1.6	3
86	Sitosterolemia in ABCG5-Null mice is aggravated upon activation of the liver X-receptor. <i>Gastroenterology</i> , 2003, 124, A727.	0.6	0
87	Reply to: "Impaired expression of multidrug resistance-associated protein 2 and liver damage in erythropoietic protoporphyria". <i>Hepatology</i> , 2016, 63, 1743-1744.	3.6	0
88	Regulation of sterol transport in mouse enterocytes by the Liver-X-Receptor LXR. , 0, 2004, .		0
89	Chronic Prednisolone Treatment Reduces Hepatic Insulin Sensitivity while Perturbing the Fed-to-Fasting Transition in Mice. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010, 95, 1477-1477.	1.8	0
90	Response to Spontaneous Cholemia in C57BL/6 Mice Predisposes to Liver Cancer in NASH. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2022, 13, 1590.	2.3	0

#	ARTICLE	IF	CITATIONS
91	Spontaneous liver disease in wild-type C57BL/6J0laHsd mice fed semisynthetic diet. , 2020, 15, e0232069.		0
92	Spontaneous liver disease in wild-type C57BL/6J0laHsd mice fed semisynthetic diet. , 2020, 15, e0232069.		0
93	Spontaneous liver disease in wild-type C57BL/6J0laHsd mice fed semisynthetic diet. , 2020, 15, e0232069.		0
94	Spontaneous liver disease in wild-type C57BL/6J0laHsd mice fed semisynthetic diet. , 2020, 15, e0232069.		0