

Vincent Walter Bloks

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/714638/vincent-walter-bloks-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

91
papers

5,901
citations

34
h-index

76
g-index

96
ext. papers

6,754
ext. citations

6.8
avg, IF

5.04
L-index

#	Paper	IF	Citations
91	Response to Spontaneous Cholemia in C57BL/6 Mice Predisposes to Liver Cancer in NASH.. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2022 ,	7.9	
90	Epigenome-wide association study of incident type 2 diabetes: a meta-analysis of five prospective European cohorts.. <i>Diabetologia</i> , 2022 , 65, 763	10.3	2
89	Mice with a deficiency in Peroxisomal Membrane Protein 4 (PXMP4) display mild changes in hepatic lipid metabolism.. <i>Scientific Reports</i> , 2022 , 12, 2512	4.9	1
88	Characterization of gut microbial structural variations as determinants of human bile acid metabolism. <i>Cell Host and Microbe</i> , 2021 , 29, 1802-1814.e5	23.4	6
87	Low production of 12 β hydroxylated bile acids prevents hepatic steatosis in Cyp2c70 mice by reducing fat absorption. <i>Journal of Lipid Research</i> , 2021 , 100134	6.3	2
86	The hepatocyte IKK:NF- κ B axis promotes liver steatosis by stimulating de novo lipogenesis and cholesterol synthesis. <i>Molecular Metabolism</i> , 2021 , 54, 101349	8.8	6
85	Potential of therapeutic bile acids in the treatment of neonatal Hyperbilirubinemia. <i>Scientific Reports</i> , 2021 , 11, 11107	4.9	2
84	An early-life diet containing large phospholipid-coated lipid globules programmes later-life postabsorptive lipid trafficking in high-fat diet- but not in low-fat diet-fed mice. <i>British Journal of Nutrition</i> , 2021 , 125, 961-971	3.6	2
83	Short-term protein restriction at advanced age stimulates FGF21 signalling, energy expenditure and browning of white adipose tissue. <i>FEBS Journal</i> , 2021 , 288, 2257-2277	5.7	6
82	Cholangiopathy and Biliary Fibrosis in Cyp2c70-Deficient Mice Are Fully Reversed by Ursodeoxycholic Acid. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2021 , 11, 1045-1069	7.9	11
81	Hepatic Carbohydrate Response Element Binding Protein Activation Limits Nonalcoholic Fatty Liver Disease Development in a Mouse Model for Glycogen Storage Disease Type 1a. <i>Hepatology</i> , 2020 , 72, 1638-1653	11.2	11
80	The effects of bariatric surgery on clinical profile, DNA methylation, and ageing in severely obese patients. <i>Clinical Epigenetics</i> , 2020 , 12, 14	7.7	7
79	Genetic and Microbial Associations to Plasma and Fecal Bile Acids in Obesity Relate to Plasma Lipids and Liver Fat Content. <i>Cell Reports</i> , 2020 , 33, 108212	10.6	22
78	An epigenome-wide association study identifies multiple DNA methylation markers of exposure to endocrine disruptors. <i>Environment International</i> , 2020 , 144, 106016	12.9	6
77	Spontaneous liver disease in wild-type C57BL/6JOLA ^{Hsd} mice fed semisynthetic diet. <i>PLoS ONE</i> , 2020 , 15, e0232069	3.7	2
76	A human-like bile acid pool induced by deletion of hepatic modulates effects of FXR activation in mice. <i>Journal of Lipid Research</i> , 2020 , 61, 291-305	6.3	52
75	Spontaneous liver disease in wild-type C57BL/6JOLA ^{Hsd} mice fed semisynthetic diet 2020 , 15, e0232069		

74	Spontaneous liver disease in wild-type C57BL/6JOlaHsd mice fed semisynthetic diet 2020 , 15, e0232069		
73	Spontaneous liver disease in wild-type C57BL/6JOlaHsd mice fed semisynthetic diet 2020 , 15, e0232069		
72	Spontaneous liver disease in wild-type C57BL/6JOlaHsd mice fed semisynthetic diet 2020 , 15, e0232069		
71	Potential of Intestine-Selective FXR Modulation for Treatment of Metabolic Disease. <i>Handbook of Experimental Pharmacology</i> , 2019 , 256, 207-234	3.2	11
70	Glucose-6-Phosphate Regulates Hepatic Bile Acid Synthesis in Mice. <i>Hepatology</i> , 2019 , 70, 2171-2184	11.2	10
69	FXR overexpression alters adipose tissue architecture in mice and limits its storage capacity leading to metabolic derangements. <i>Journal of Lipid Research</i> , 2019 , 60, 1547-1561	6.3	8
68	Transcriptome analysis suggests a compensatory role of the cofactors coenzyme A and NAD in medium-chain acyl-CoA dehydrogenase knockout mice. <i>Scientific Reports</i> , 2019 , 9, 14539	4.9	2
67	New insights in the multiple roles of bile acids and their signaling pathways in metabolic control. <i>Current Opinion in Lipidology</i> , 2018 , 29, 194-202	4.4	37
66	Epigenetic programming at the Mogat1 locus may link neonatal overnutrition with long-term hepatic steatosis and insulin resistance. <i>FASEB Journal</i> , 2018 , 32, fj201700717RR	0.9	16
65	Milk cholesterol concentration in mice is not affected by high cholesterol diet- or genetically-induced hypercholesterolaemia. <i>Scientific Reports</i> , 2018 , 8, 8824	4.9	4
64	Intestinal Farnesoid X Receptor Controls Transintestinal Cholesterol Excretion in Mice. <i>Gastroenterology</i> , 2017 , 152, 1126-1138.e6	13.3	89
63	Intestinal PPAR α protects against diet-induced obesity, insulin resistance and dyslipidemia. <i>Scientific Reports</i> , 2017 , 7, 846	4.9	21
62	Male apoE*3-Leiden.CETP mice on high-fat high-cholesterol diet exhibit a biphasic dyslipidemic response, mimicking the changes in plasma lipids observed through life in men. <i>Physiological Reports</i> , 2017 , 5, e13376	2.6	12
61	Transintestinal Cholesterol Transport Is Active in Mice and Humans and Controls Ezetimibe-Induced Fecal Neutral Sterol Excretion. <i>Cell Metabolism</i> , 2016 , 24, 783-794	24.6	90
60	Statins increase hepatic cholesterol synthesis and stimulate fecal cholesterol elimination in mice. <i>Journal of Lipid Research</i> , 2016 , 57, 1455-64	6.3	74
59	Hypertrophy induced KIF5B controls mitochondrial localization and function in neonatal rat cardiomyocytes. <i>Journal of Molecular and Cellular Cardiology</i> , 2016 , 97, 70-81	5.8	11
58	Malnutrition-associated liver steatosis and ATP depletion is caused by peroxisomal and mitochondrial dysfunction. <i>Journal of Hepatology</i> , 2016 , 65, 1198-1208	13.4	78
57	Reply to: "Impaired expression of multidrug resistance-associated protein 2 and liver damage in erythropoietic protoporphyria". <i>Hepatology</i> , 2016 , 63, 1743-4	11.2	

56	HSPA6 is an ulcerative colitis susceptibility factor that is induced by cigarette smoke and protects intestinal epithelial cells by stabilizing anti-apoptotic Bcl-XL. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2016 , 1862, 788-796	6.9	12
55	Clinical symptoms of right ventricular failure in experimental chronic pressure load are associated with progressive diastolic dysfunction. <i>Journal of Molecular and Cellular Cardiology</i> , 2015 , 79, 244-53	5.8	30
54	Hypoxia and Complement-and-Coagulation Pathways in the Deceased Organ Donor as the Major Target for Intervention to Improve Renal Allograft Outcome. <i>Transplantation</i> , 2015 , 99, 1293-300	1.8	64
53	Cross-talk between liver and intestine in control of cholesterol and energy homeostasis. <i>Molecular Aspects of Medicine</i> , 2014 , 37, 77-88	16.7	18
52	In utero undernutrition in male mice programs liver lipid metabolism in the second-generation offspring involving altered Lxra DNA methylation. <i>Cell Metabolism</i> , 2014 , 19, 941-51	24.6	137
51	Beyond intestinal soap--bile acids in metabolic control. <i>Nature Reviews Endocrinology</i> , 2014 , 10, 488-98	15.2	280
50	Maternal western diet primes non-alcoholic fatty liver disease in adult mouse offspring. <i>Acta Physiologica</i> , 2014 , 210, 215-27	5.6	61
49	Hepatic farnesoid X-receptor isoforms α and β differentially modulate bile salt and lipoprotein metabolism in mice. <i>PLoS ONE</i> , 2014 , 9, e115028	3.7	23
48	A novel approach to monitor glucose metabolism using stable isotopically labelled glucose in longitudinal studies in mice. <i>Laboratory Animals</i> , 2013 , 47, 79-88	2.6	42
47	Bile acid sequestration normalizes plasma cholesterol and reduces atherosclerosis in hypercholesterolemic mice. No additional effect of physical activity. <i>Atherosclerosis</i> , 2013 , 228, 117-23	3.1	15
46	Transfer of intestinal microbiota from lean donors increases insulin sensitivity in individuals with metabolic syndrome. <i>Gastroenterology</i> , 2012 , 143, 913-6.e7	13.3	1766
45	Sex-dependent programming of glucose and fatty acid metabolism in mouse offspring by maternal protein restriction. <i>Gender Medicine</i> , 2012 , 9, 166-179.e13		13
44	Chronic prednisolone treatment aggravates hyperglycemia in mice fed a high-fat diet but does not worsen dietary fat-induced insulin resistance. <i>Endocrinology</i> , 2012 , 153, 3713-23	4.8	10
43	Liver X receptor activation restores memory in aged AD mice without reducing amyloid. <i>Neurobiology of Aging</i> , 2011 , 32, 1262-72	5.6	101
42	Resistance to diet-induced adiposity in cannabinoid receptor-1 deficient mice is not due to impaired adipocyte function. <i>Nutrition and Metabolism</i> , 2011 , 8, 93	4.6	3
41	Cerebral accumulation of dietary derivable plant sterols does not interfere with memory and anxiety related behavior in <i>Abcg5</i> ^{-/-} mice. <i>Plant Foods for Human Nutrition</i> , 2011 , 66, 149-56	3.9	26
40	Liver receptor homolog-1 is critical for adequate up-regulation of <i>Cyp7a1</i> gene transcription and bile salt synthesis during bile salt sequestration. <i>Hepatology</i> , 2011 , 53, 2075-85	11.2	41
39	Metabolic responses to long-term pharmacological inhibition of CB1-receptor activity in mice in relation to dietary fat composition. <i>International Journal of Obesity</i> , 2010 , 34, 374-84	5.5	8

38	Chronic prednisolone treatment reduces hepatic insulin sensitivity while perturbing the fed-to-fasting transition in mice. <i>Endocrinology</i> , 2010 , 151, 2171-8	4.8	20
37	Two time-point assessment of bile acid kinetics in humans using stable isotopes. <i>Isotopes in Environmental and Health Studies</i> , 2010 , 46, 325-36	1.5	3
36	The liver X-receptor gene promoter is hypermethylated in a mouse model of prenatal protein restriction. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2010 , 298, R275-82	3.2	116
35	Alterations in brain cholesterol metabolism in the APPSLxPS1mut mouse, a model for Alzheimer's disease. <i>Journal of Alzheimers Disease</i> , 2010 , 19, 117-27	4.3	29
34	Improved glycemic control with colesevelam treatment in patients with type 2 diabetes is not directly associated with changes in bile acid metabolism. <i>Hepatology</i> , 2010 , 52, 1455-64	11.2	140
33	An increased flux through the glucose 6-phosphate pool in enterocytes delays glucose absorption in Fxr ^{-/-} mice. <i>Journal of Biological Chemistry</i> , 2009 , 284, 10315-23	5.4	45
32	Regulation of bile acid synthesis by the nuclear receptor Rev-erb α . <i>Gastroenterology</i> , 2008 , 135, 689-98	4.3	156
31	Plant sterols cause macrothrombocytopenia in a mouse model of sitosterolemia. <i>Journal of Biological Chemistry</i> , 2008 , 283, 6281-7	5.4	34
30	Secretory phospholipase A2 increases SR-BI-mediated selective uptake from HDL but not biliary cholesterol secretion. <i>Journal of Lipid Research</i> , 2008 , 49, 563-71	6.3	20
29	Neonatal dexamethasone administration causes progressive renal damage due to induction of an early inflammatory response. <i>American Journal of Physiology - Renal Physiology</i> , 2008 , 294, F768-76	4.3	9
28	Lxr α deficiency hampers the hepatic adaptive response to fasting in mice. <i>Journal of Biological Chemistry</i> , 2008 , 283, 25437-25445	5.4	28
27	Disturbed hepatic carbohydrate management during high metabolic demand in medium-chain acyl-CoA dehydrogenase (MCAD)-deficient mice. <i>Hepatology</i> , 2008 , 47, 1894-904	11.2	34
26	Cholesterol feeding strongly reduces hepatic VLDL-triglyceride production in mice lacking the liver X receptor α . <i>Journal of Lipid Research</i> , 2007 , 48, 337-47	6.3	32
25	The phosphatidylethanolamine N-methyltransferase pathway is quantitatively not essential for biliary phosphatidylcholine secretion. <i>Journal of Lipid Research</i> , 2007 , 48, 2058-64	6.3	14
24	Postnatal treatment with dexamethasone perturbs hepatic and cardiac energy metabolism and is associated with a sustained atherogenic plasma lipid profile in suckling rats. <i>Pediatric Research</i> , 2007 , 61, 165-70	3.2	11
23	Gene expression profiling in livers of mice after acute inhibition of beta-oxidation. <i>Genomics</i> , 2007 , 90, 680-9	4.3	31
22	Abcg5/Abcg8-independent pathways contribute to hepatobiliary cholesterol secretion in mice. <i>American Journal of Physiology - Renal Physiology</i> , 2006 , 291, G414-23	5.1	44
21	24(S)-hydroxycholesterol participates in a liver X receptor-controlled pathway in astrocytes that regulates apolipoprotein E-mediated cholesterol efflux. <i>Journal of Biological Chemistry</i> , 2006 , 281, 12799-808	5.4	166

20	Reduction of cholesterol absorption by dietary plant sterols and stanols in mice is independent of the Abcg5/8 transporter. <i>Journal of Nutrition</i> , 2006 , 136, 2135-40	4.1	67
19	Rosuvastatin reduces plasma lipids by inhibiting VLDL production and enhancing hepatobiliary lipid excretion in ApoE*3-leiden mice. <i>Journal of Cardiovascular Pharmacology</i> , 2005 , 45, 53-60	3.1	16
18	Essential fatty acid deficiency in mice is associated with hepatic steatosis and secretion of large VLDL particles. <i>American Journal of Physiology - Renal Physiology</i> , 2005 , 288, G1150-8	5.1	35
17	Down-regulation of hepatic and intestinal Abcg5 and Abcg8 expression associated with altered sterol fluxes in rats with streptozotocin-induced diabetes. <i>Diabetologia</i> , 2004 , 47, 104-12	10.3	56
16	Sitosterolemia in ABC-transporter G5-deficient mice is aggravated on activation of the liver-X receptor. <i>Gastroenterology</i> , 2004 , 126, 290-300	13.3	124
15	Peroxisome proliferator-activated receptor alpha (PPARalpha)-mediated regulation of multidrug resistance 2 (Mdr2) expression and function in mice. <i>Biochemical Journal</i> , 2003 , 369, 539-47	3.8	141
14	Induction of hepatic ABC transporter expression is part of the PPARalpha-mediated fasting response in the mouse. <i>Gastroenterology</i> , 2003 , 124, 160-71	13.3	66
13	Mdr P-glycoproteins are not essential for biliary excretion of the hydrophobic heme precursor protoporphyrin in a griseofulvin-induced mouse model of erythropoietic protoporphyria. <i>Hepatology</i> , 2002 , 35, 299-306	11.2	19
12	Increased hepatobiliary and fecal cholesterol excretion upon activation of the liver X receptor is independent of ABCA1. <i>Journal of Biological Chemistry</i> , 2002 , 277, 33870-7	5.4	153
11	Stimulation of lipogenesis by pharmacological activation of the liver X receptor leads to production of large, triglyceride-rich very low density lipoprotein particles. <i>Journal of Biological Chemistry</i> , 2002 , 277, 34182-90	5.4	380
10	Fat malabsorption in essential fatty acid-deficient mice is not due to impaired bile formation. <i>American Journal of Physiology - Renal Physiology</i> , 2002 , 283, G900-8	5.1	23
9	Differential effects of streptozotocin-induced diabetes on expression of hepatic ABC-transporters in rats. <i>Gastroenterology</i> , 2002 , 122, 1842-52	13.3	64
8	The role of transhepatic bile salt flux in the control of hepatic secretion of triacylglycerol-rich lipoproteins in vivo in rodents. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2002 , 1573, 9-20	4	9
7	Hepatobiliary cholesterol transport is not impaired in Abca1-null mice lacking HDL. <i>Journal of Clinical Investigation</i> , 2001 , 108, 843-850	15.9	120
6	Hyperlipidemia and atherosclerosis associated with liver disease in ferrochelatase-deficient mice. <i>Journal of Lipid Research</i> , 2001 , 42, 41-50	6.3	32
5	Hepatic lipid accumulation, altered very low density lipoprotein formation and apolipoprotein E deposition in apolipoprotein E3-Leiden transgenic mice. <i>Journal of Hepatology</i> , 2000 , 33, 189-98	13.4	26
4	Apolipoprotein E participates in the regulation of very low density lipoprotein-triglyceride secretion by the liver. <i>Journal of Biological Chemistry</i> , 1999 , 274, 35711-8	5.4	103
3	3-Hydroxy-3-methylglutaryl-coenzyme A reductase inhibitors (statins) induce hepatic expression of the phospholipid translocase mdr2 in rats. <i>Gastroenterology</i> , 1999 , 117, 678-87	13.3	61

- 2 Biliary fibrosis associated with altered bile composition in a mouse model of erythropoietic protoporphyria. *Gastroenterology*, **1999**, 117, 696-705 133 83
- 1 Impaired secretion of very low density lipoprotein-triglycerides by apolipoprotein E- deficient mouse hepatocytes. *Journal of Clinical Investigation*, **1997**, 100, 2915-22 159 140