

# Yong-gui Yuan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/715927/publications.pdf>

Version: 2024-02-01

121  
papers

3,280  
citations

196777

29  
h-index

223390

49  
g-index

126  
all docs

126  
docs citations

126  
times ranked

4506  
citing authors

#	ARTICLE	IF	CITATIONS
1	Influence of sleep difficulty on post-traumatic stress symptoms among frontline medical staff during COVID-19 pandemic in China. <i>Psychology, Health and Medicine</i> , 2022, 27, 1924-1936.	1.3	3
2	Impaired robust interhemispheric function integration of depressive brain from REST meta-MDD database in China. <i>Bipolar Disorders</i> , 2022, 24, 400-411.	1.1	8
3	The Reliability and Validity of Post Stroke Depression Scale in Different Type of Post Stroke Depression Patients. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2022, 31, 106222.	0.7	3
4	The impact of HTR1A and HTR1B methylation combined with stress/genotype on early antidepressant efficacy. <i>Psychiatry and Clinical Neurosciences</i> , 2022, 76, 51-57.	1.0	6
5	Combination of spontaneous regional brain activity and HTR1A/1B DNA methylation to predict early responses to antidepressant treatments in MDD. <i>Journal of Affective Disorders</i> , 2022, 302, 249-257.	2.0	4
6	Bacteroides species differentially modulate depression-like behavior via gut-brain metabolic signaling. <i>Brain, Behavior, and Immunity</i> , 2022, 102, 11-22.	2.0	66
7	Characteristics of post-traumatic embitterment disorder of inpatients in a general hospital in China. <i>Clinical Psychology and Psychotherapy</i> , 2022, , .	1.4	1
8	Influence and interaction of resting state functional magnetic resonance and tryptophan hydroxylase-2 methylation on short-term antidepressant drug response. <i>BMC Psychiatry</i> , 2022, 22, 218.	1.1	5
9	Clinical Efficacy of the Chinese Herbal Medicine Shumian Capsule for Insomnia: A Randomized, Double-Blind, Placebo-Controlled Trial. <i>Neuropsychiatric Disease and Treatment</i> , 2022, Volume 18, 669-679.	1.0	2
10	Reliability and Validity of the Urdu Version of Psychosomatic Symptoms Scale in Pakistani Patients. <i>Frontiers in Psychology</i> , 2022, 13, 861859.	1.1	6
11	Spatio-Temporal Attention Graph Convolution Network for Functional Connectome Classification. , 2022, , .		2
12	The interaction of P11 methylation and early-life stress impacts the antidepressant response in patients with major depressive disorder. <i>Journal of Affective Disorders</i> , 2022, 312, 128-135.	2.0	4
13	Decreased cortical thickness of left premotor cortex as a treatment predictor in major depressive disorder. <i>Brain Imaging and Behavior</i> , 2021, 15, 1420-1426.	1.1	6
14	Reliability and validity of the Chinese version of the post-traumatic embitterment disorder self-rating scale (PTED-21) among inpatients in general hospital. <i>Clinical Psychology and Psychotherapy</i> , 2021, 28, 882-890.	1.4	3
15	Identification of first-episode unmedicated major depressive disorder using pretreatment features of dominant coactivation patterns. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 104, 110038.	2.5	8
16	Identification of specific neural circuit underlying the key cognitive deficit of remitted late-onset depression: A multi-modal MRI and machine learning study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 108, 110192.	2.5	7
17	Desynchronized Functional Activities Between Brain White and Gray Matter in Major Depression Disorder. <i>Journal of Magnetic Resonance Imaging</i> , 2021, 53, 1375-1386.	1.9	10
18	Asthma-Specific Temporal Variability Reveals the Effect of Group Cognitive Behavior Therapy in Asthmatic Patients. <i>Frontiers in Neurology</i> , 2021, 12, 615820.	1.1	0

#	ARTICLE	IF	CITATIONS
19	Disrupted hemispheric connectivity specialization in patients with major depressive disorder: Evidence from the REST-meta-MDD Project. <i>Journal of Affective Disorders</i> , 2021, 284, 217-228.	2.0	23
20	Clinimetric properties of the Chinese version of the Euthymia Scale. <i>Clinical Psychology and Psychotherapy</i> , 2021, , .	1.4	8
21	Spatio-temporal graph convolutional network for diagnosis and treatment response prediction of major depressive disorder from functional connectivity. <i>Human Brain Mapping</i> , 2021, 42, 3922-3933.	1.9	28
22	A Bibliometric Analysis of the One Hundred Most Cited Studies in Psychosomatic Research. <i>Psychotherapy and Psychosomatics</i> , 2021, 90, 425-430.	4.0	29
23	Coupling of spatial and directional functional network connectivity reveals a physiological basis for salience network hubs in asthma. <i>Brain Imaging and Behavior</i> , 2021, , 1.	1.1	3
24	Higher baseline serum adiponectin predicts better treatment remission in patients with generalized anxiety disorder treated with escitalopram. <i>Annals of Palliative Medicine</i> , 2021, 10, 7634-7643.	0.5	1
25	Effect of NEUROG3 polymorphism rs144643855 on regional spontaneous brain activity in major depressive disorder. <i>Behavioural Brain Research</i> , 2021, 409, 113310.	1.2	4
26	The Combination of Serum BDNF, Cortisol and IFN-Gamma Can Assist the Diagnosis of Major Depressive Disorder. <i>Neuropsychiatric Disease and Treatment</i> , 2021, Volume 17, 2819-2829.	1.0	13
27	Disrupted intrinsic functional brain topology in patients with major depressive disorder. <i>Molecular Psychiatry</i> , 2021, 26, 7363-7371.	4.1	82
28	Global topology alteration of the brain functional network affects the 8-week antidepressant response in major depressive disorder. <i>Journal of Affective Disorders</i> , 2021, 294, 491-496.	2.0	15
29	Brain structural alterations in MDD patients with gastrointestinal symptoms: Evidence from the REST-meta-MDD project. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2021, 111, 110386.	2.5	18
30	DNA methylations of brain-derived neurotrophic factor exon VI are associated with major depressive disorder and antidepressant-induced remission in females. <i>Journal of Affective Disorders</i> , 2021, 295, 101-107.	2.0	17
31	Sleep disturbance-related neuroimaging features as potential biomarkers for the diagnosis of major depressive disorder: A multicenter study based on machine learning. <i>Journal of Affective Disorders</i> , 2021, 295, 148-155.	2.0	9
32	Early Enhancement of Neuroplasticity Index, the Ratio of Serum Brain-Derived Neurotrophic Factor Level to HAMD-24 Score, in Predicting the Long-Term Antidepressant Efficacy. <i>Frontiers in Behavioral Neuroscience</i> , 2021, 15, 712445.	1.0	0
33	Clinical characteristics of comorbid post-traumatic embitterment disorder and major depressive disorder patients in China. <i>General Hospital Psychiatry</i> , 2021, 74, 147-147.	1.2	0
34	Distinct Features of Cerebral Blood Flow and Spontaneous Neural Activity as Integrated Predictors of Early Response to Antidepressants. <i>Frontiers in Psychiatry</i> , 2021, 12, 788398.	1.3	0
35	Disrupted structural brain connectome underlying the cognitive deficits in remitted late-onset depression. <i>Brain Imaging and Behavior</i> , 2020, 14, 1600-1611.	1.1	20
36	Altered Brain Entropy as a predictor of antidepressant response in major depressive disorder. <i>Journal of Affective Disorders</i> , 2020, 260, 716-721.	2.0	16

#	ARTICLE	IF	CITATIONS
37	Altered resting-state dynamic functional brain networks in major depressive disorder: Findings from the REST-meta-MDD consortium. <i>NeuroImage: Clinical</i> , 2020, 26, 102163.	1.4	76
38	Predicting Response to Group Cognitive Behavioral Therapy in Asthma by a Small Number of Abnormal Resting-State Functional Connections. <i>Frontiers in Neuroscience</i> , 2020, 14, 575771.	1.4	1
39	Aberrant amplitude low-frequency fluctuation (ALFF) and regional homogeneity (ReHo) in generalized anxiety disorder (GAD) and their roles in predicting treatment remission. <i>Annals of Translational Medicine</i> , 2020, 8, 1319-1319.	0.7	26
40	Biotypes of major depressive disorder: Neuroimaging evidence from resting-state default mode network patterns. <i>NeuroImage: Clinical</i> , 2020, 28, 102514.	1.4	51
41	The relationship of tryptophan hydroxylase-2 methylation to early-life stress and its impact on short-term antidepressant treatment response. <i>Journal of Affective Disorders</i> , 2020, 276, 850-858.	2.0	19
42	Influence of genetic polymorphisms in homocysteine and lipid metabolism systems on antidepressant drug response. <i>BMC Psychiatry</i> , 2020, 20, 408.	1.1	8
43	Dopamine Multilocus Genetic Profile, Spontaneous Activity of Left Superior Temporal Gyrus, and Early Therapeutic Effect in Major Depressive Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 591407.	1.3	9
44	The effect of mGlu2/3 receptors on synaptic activities to different types of GABAergic interneurons in the anterior cingulate cortex. <i>Neuropharmacology</i> , 2020, 175, 108180.	2.0	2
45	CACNA1C Gene rs11832738 Polymorphism Influences Depression Severity by Modulating Spontaneous Activity in the Right Middle Frontal Gyrus in Patients With Major Depressive Disorder. <i>Frontiers in Psychiatry</i> , 2020, 11, 73.	1.3	14
46	Development of the psychosomatic symptom scale (PSSS) and assessment of its reliability and validity in general hospital patients in China. <i>General Hospital Psychiatry</i> , 2020, 64, 1-8.	1.2	16
47	Predicting conversion to Alzheimer's disease among individual high-risk patients using the Characterizing AD Risk Events index model. <i>CNS Neuroscience and Therapeutics</i> , 2020, 26, 720-729.	1.9	4
48	Amygdala connectivity mediates the association between anxiety and depression in patients with major depressive disorder. <i>Brain Imaging and Behavior</i> , 2019, 13, 1146-1159.	1.1	41
49	Increased interhemispheric synchrony underlying the improved athletic performance of rowing athletes by transcranial direct current stimulation. <i>Brain Imaging and Behavior</i> , 2019, 13, 1324-1332.	1.1	8
50	Health anxiety in medical employees: A multicentre study. <i>Journal of International Medical Research</i> , 2019, 47, 4854-4861.	0.4	17
51	Serum BICC1 levels are significantly different in various mood disorders. <i>Neuropsychiatric Disease and Treatment</i> , 2019, Volume 15, 259-265.	1.0	4
52	Baseline serum C-reactive protein levels may predict antidepressant treatment responses in patients with major depressive disorder. <i>Journal of Affective Disorders</i> , 2019, 250, 432-438.	2.0	25
53	Shared genetic risk factors for depression and stroke. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2019, 93, 55-70.	2.5	30
54	Reduced default mode network functional connectivity in patients with recurrent major depressive disorder. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 9078-9083.	3.3	441

#	ARTICLE	IF	CITATIONS
55	Higher serum VGF protein levels discriminate bipolar depression from major depressive disorder. <i>Journal of Neuroscience Research</i> , 2019, 97, 597-606.	1.3	10
56	The roles of brain-derived neurotrophic factor (BDNF) and glial cell line-derived neurotrophic factor (GDNF) in predicting treatment remission in a Chinese Han population with generalized anxiety disorder. <i>Psychiatry Research</i> , 2019, 271, 319-324.	1.7	25
57	Risk factors associated with cognitions for late-onset depression based on anterior and posterior default mode sub-networks. <i>Journal of Affective Disorders</i> , 2018, 235, 544-550.	2.0	7
58	Increased temporal variability of striatum region facilitating the early antidepressant response in patients with major depressive disorder. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2018, 85, 39-45.	2.5	40
59	Association Analysis of the Brain-Derived Neurotrophic Factor Gene <i>Val66Met</i> Polymorphism and Gender with Efficacy of Antidepressants in the Chinese Han Population with Generalized Anxiety Disorder. <i>Genetic Testing and Molecular Biomarkers</i> , 2018, 22, 199-206.	0.3	4
60	Effects of tandospirone augmentation in major depressive disorder patients with high anxiety: A multicenter, randomized, parallel-controlled, open-label study. <i>Journal of Psychiatric Research</i> , 2018, 99, 104-110.	1.5	22
61	White Matter Integrity Disruptions Correlate With Cognitive Impairments in Asthma. <i>Journal of Magnetic Resonance Imaging</i> , 2018, 48, 748-756.	1.9	14
62	Abnormal ventral tegmental area-anterior cingulate cortex connectivity in Parkinson's disease with depression. <i>Behavioural Brain Research</i> , 2018, 347, 132-139.	1.2	24
63	Investigation of health anxiety and related factors in Chinese patients with physical disease. <i>Perspectives in Psychiatric Care</i> , 2018, 54, 185-191.	0.9	1
64	Distinctive pretreatment features of bilateral nucleus accumbens networks predict early response to antidepressants in major depressive disorder. <i>Brain Imaging and Behavior</i> , 2018, 12, 1042-1052.	1.1	19
65	Disrupted topology of hippocampal connectivity is associated with short-term antidepressant response in major depressive disorder. <i>Journal of Affective Disorders</i> , 2018, 225, 539-544.	2.0	25
66	Decreased cerebral blood flow in the primary motor cortex in major depressive disorder with psychomotor retardation. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2018, 81, 438-444.	2.5	37
67	Promoter haplotypes of interleukin-10 gene linked to cortex plasticity in subjects with risk of Alzheimer's disease. <i>NeuroImage: Clinical</i> , 2018, 17, 587-595.	1.4	6
68	Predictive brain networks for major depression in a semi-multimodal fusion hierarchical feature reduction framework. <i>Neuroscience Letters</i> , 2018, 665, 163-169.	1.0	12
69	Cognitive Deficit-Related Interhemispheric Asynchrony within the Medial Hub of the Default Mode Network Aids in Classifying the Hyperthyroid Patients. <i>Neural Plasticity</i> , 2018, 2018, 1-7.	1.0	2
70	Aberrant Default Mode Network Underlying the Cognitive Deficits in the Patients With Late-Onset Depression. <i>Frontiers in Aging Neuroscience</i> , 2018, 10, 310.	1.7	14
71	Altered Regional Cerebral Blood Flow of Right Cerebellum Posterior Lobe in Asthmatic Patients With or Without Depressive Symptoms. <i>Frontiers in Psychiatry</i> , 2018, 9, 225.	1.3	14
72	Clinical practice guidelines for post-stroke depression in China. <i>Revista Brasileira De Psiquiatria</i> , 2018, 40, 325-334.	0.9	36

#	ARTICLE	IF	CITATIONS
73	Abnormal spontaneous brain activity is associated with impaired emotion and cognition in hyperthyroidism: A rs-fMRI study. <i>Behavioural Brain Research</i> , 2018, 351, 188-194.	1.2	13
74	Reduced serum VGF levels were reversed by antidepressant treatment in depressed patients. <i>World Journal of Biological Psychiatry</i> , 2017, 18, 586-591.	1.3	15
75	The protein and mRNA expression levels of glial cell line-derived neurotrophic factor in post stroke depression and major depressive disorder. <i>Scientific Reports</i> , 2017, 7, 8674.	1.6	18
76	Combined serum levels of multiple proteins in tPA-BDNF pathway may aid the diagnosis of five mental disorders. <i>Scientific Reports</i> , 2017, 7, 6871.	1.6	27
77	Group Cognitive Behavior Therapy Reversed Abnormal Spontaneous Brain Activity in Adult Asthmatic Patients. <i>Psychotherapy and Psychosomatics</i> , 2017, 86, 178-180.	4.0	8
78	Aberrant Intra- and Internetwork Functional Connectivity in Depressed Parkinson's Disease. <i>Scientific Reports</i> , 2017, 7, 2568.	1.6	33
79	Disrupted reward circuits is associated with cognitive deficits and depression severity in major depressive disorder. <i>Journal of Psychiatric Research</i> , 2017, 84, 9-17.	1.5	64
80	A risk prediction model for post-stroke depression in Chinese stroke survivors based on clinical and socio-psychological features. <i>Oncotarget</i> , 2017, 8, 62891-62899.	0.8	15
81	Group Cognitive Behavior Therapy Reversed Insula Subregions Functional Connectivity in Asthmatic Patients. <i>Frontiers in Aging Neuroscience</i> , 2017, 9, 105.	1.7	10
82	Mediating Role of the Reward Network in the Relationship between the Dopamine Multilocus Genetic Profile and Depression. <i>Frontiers in Molecular Neuroscience</i> , 2017, 10, 292.	1.4	14
83	Abnormal Functional Connectivity of Ventral Anterior Insula in Asthmatic Patients with Depression. <i>Neural Plasticity</i> , 2017, 2017, 1-11.	1.0	8
84	Amygdala Atrophy and Its Functional Disconnection with the Cortico-Striatal-Pallidal-Thalamic Circuit in Major Depressive Disorder in Females. <i>PLoS ONE</i> , 2017, 12, e0168239.	1.1	40
85	New opinion on the subtypes of poststroke depression in Chinese stroke survivors. <i>Neuropsychiatric Disease and Treatment</i> , 2017, Volume 13, 707-713.	1.0	6
86	Abnormal brain functional connectivity leads to impaired mood and cognition in hyperthyroidism: a resting-state functional MRI study. <i>Oncotarget</i> , 2017, 8, 6283-6294.	0.8	30
87	Disrupted Interhemispheric Synchrony in Default Mode Network Underlying the Impairment of Cognitive Flexibility in Late-Onset Depression. <i>Frontiers in Aging Neuroscience</i> , 2016, 8, 230.	1.7	31
88	Management of Poststroke Neuropsychiatric Disorders. <i>Translational Neuroscience and Clinics</i> , 2016, 2, 244-251.	0.1	3
89	Aberrant topographical organization of the default mode network underlying the cognitive impairment of remitted late-onset depression. <i>Neuroscience Letters</i> , 2016, 629, 26-32.	1.0	21
90	Impaired interhemispheric synchrony in Parkinson's disease with depression. <i>Scientific Reports</i> , 2016, 6, 27477.	1.6	30

#	ARTICLE	IF	CITATIONS
91	Plastic modulation of episodic memory networks in the aging brain with cognitive decline. <i>Behavioural Brain Research</i> , 2016, 308, 38-45.	1.2	6
92	TPH-2 Polymorphisms Interact with Early Life Stress to Influence Response to Treatment with Antidepressant Drugs. <i>International Journal of Neuropsychopharmacology</i> , 2016, 19, pyw070.	1.0	23
93	Structural and Functional Connectivity of Default Mode Network underlying the Cognitive Impairment in Late-onset Depression. <i>Scientific Reports</i> , 2016, 6, 37617.	1.6	35
94	Correlation of 5-HTT, BDNF and NPSR1 gene polymorphisms with anxiety and depression in asthmatic patients. <i>International Journal of Molecular Medicine</i> , 2016, 38, 65-74.	1.8	22
95	Genetics pathway-based imaging approaches in Chinese Han population with Alzheimer's disease risk. <i>Brain Structure and Function</i> , 2016, 221, 433-446.	1.2	8
96	Immunity factor contributes to altered brain functional networks in individuals at risk for Alzheimer's disease: Neuroimaging-genetic evidence. <i>Brain, Behavior, and Immunity</i> , 2016, 56, 84-95.	2.0	5
97	Towards a multi protein and mRNA expression of biological predictive and distinguish model for post stroke depression. <i>Oncotarget</i> , 2016, 7, 54329-54338.	0.8	12
98	Multiple genetic imaging study of the association between cholesterol metabolism and brain functional alterations in individuals with risk factors for Alzheimer's disease. <i>Oncotarget</i> , 2016, 7, 15315-15328.	0.8	11
99	Association of specific frequency bands of functional MRI signal oscillations with motor symptoms and depression in Parkinson's disease. <i>Scientific Reports</i> , 2015, 5, 16376.	1.6	21
100	A comparative study of anhedonia components between major depression and schizophrenia in Chinese populations. <i>Annals of General Psychiatry</i> , 2015, 14, 24.	1.2	33
101	The reliability and validity of a Chinese-version Short Health Anxiety Inventory: an investigation of university students. <i>Neuropsychiatric Disease and Treatment</i> , 2015, 11, 1739.	1.0	35
102	Altered Resting-State Brain Activity and Connectivity in Depressed Parkinson's Disease. <i>PLoS ONE</i> , 2015, 10, e0131133.	1.1	53
103	Frequency-Dependent Amplitude Alterations of Resting-State Spontaneous Fluctuations in Late-Onset Depression. <i>BioMed Research International</i> , 2015, 2015, 1-9.	0.9	49
104	The BDNF Val66Met polymorphism, resting-state hippocampal functional connectivity and cognitive deficits in acute late-onset depression. <i>Journal of Affective Disorders</i> , 2015, 183, 22-30.	2.0	40
105	Tryptophan hydroxylase 2 gene is associated with cognition in late-onset depression in a Chinese Han population. <i>Neuroscience Letters</i> , 2015, 600, 98-103.	1.0	11
106	Changed Synaptic Plasticity in Neural Circuits of Depressive-Like and Escitalopram-Treated Rats. <i>International Journal of Neuropsychopharmacology</i> , 2015, 18, pyv046.	1.0	52
107	Abnormal cerebral functional connectivity in esophageal cancer patients with theory of mind deficits in resting state. <i>Journal of Cancer Research and Therapeutics</i> , 2015, 11, 438.	0.3	4
108	Investigation of health anxiety and its related factors in nursing students. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 1223.	1.0	24

#	ARTICLE	IF	CITATIONS
109	Association analysis of COMT/MTHFR polymorphisms and major depressive disorder in Chinese Han population. <i>Journal of Affective Disorders</i> , 2014, 161, 73-78.	2.0	31
110	Imbalanced hippocampal functional networks associated with remitted geriatric depression and apolipoprotein E $\epsilon$ 4 allele in nondemented elderly: A preliminary study. <i>Journal of Affective Disorders</i> , 2014, 164, 5-13.	2.0	48
111	Neuroimaging studies of depressive disorders in China since 2000. <i>Shanghai Archives of Psychiatry</i> , 2014, 26, 113-8.	0.7	0
112	Abnormal Functional Connectivity of Amygdala in Late-Onset Depression Was Associated with Cognitive Deficits. <i>PLoS ONE</i> , 2013, 8, e75058.	1.1	92
113	Influence and interaction of genetic polymorphisms in the serotonin system and life stress on antidepressant drug response. <i>Journal of Psychopharmacology</i> , 2012, 26, 349-359.	2.0	60
114	Topologically Convergent and Divergent Structural Connectivity Patterns between Patients with Remitted Geriatric Depression and Amnesic Mild Cognitive Impairment. <i>Journal of Neuroscience</i> , 2012, 32, 4307-4318.	1.7	282
115	Mapping the Altered Patterns of Cerebellar Resting-State Function in Longitudinal Amnesic Mild Cognitive Impairment Patients. <i>Journal of Alzheimer's Disease</i> , 2011, 23, 87-99.	1.2	51
116	Genetic variation in apolipoprotein E alters regional gray matter volumes in remitted late-onset depression. <i>Journal of Affective Disorders</i> , 2010, 121, 273-277.	2.0	43
117	Abnormal Integrity of Long Association Fiber Tracts Is Associated With Cognitive Deficits in Patients With Remitted Geriatric Depression. <i>Journal of Clinical Psychiatry</i> , 2010, 71, 1386-1390.	1.1	22
118	Larger regional white matter volume is associated with executive function deficit in remitted geriatric depression: An optimized voxel-based morphometry study. <i>Journal of Affective Disorders</i> , 2009, 115, 225-229.	2.0	22
119	Abnormal neural activity in the patients with remitted geriatric depression: A resting-state functional magnetic resonance imaging study. <i>Journal of Affective Disorders</i> , 2008, 111, 145-152.	2.0	122
120	Regional Gray Matter Changes Are Associated with Cognitive Deficits in Remitted Geriatric Depression: An Optimized Voxel-Based Morphometry Study. <i>Biological Psychiatry</i> , 2008, 64, 541-544.	0.7	80
121	White matter integrity of the whole brain is disrupted in first-episode remitted geriatric depression. <i>NeuroReport</i> , 2007, 18, 1845-1849.	0.6	63