

Koh-Ichi Nakashiro

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7169507/koh-ichi-nakashiro-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

92
papers

2,086
citations

27
h-index

43
g-index

95
ext. papers

2,266
ext. citations

3.8
avg, IF

4.14
L-index

#	Paper	IF	Citations
92	Oral squamous cell carcinoma may originate from bone marrow-derived stem cells. <i>Oncology Letters</i> , 2021 , 21, 170	2.6	1
91	MicroRNA-361-3p is a potent therapeutic target for oral squamous cell carcinoma. <i>Cancer Science</i> , 2020 , 111, 1645-1651	6.9	11
90	Valve Interstitial Cell-Specific Cyclooxygenase-1 Associated With Calcification of Aortic Valves. <i>Annals of Thoracic Surgery</i> , 2020 , 110, 40-49	2.7	2
89	Paclitaxel Potentiates the Anticancer Effect of Cetuximab by Enhancing Antibody-Dependent Cellular Cytotoxicity on Oral Squamous Cell Carcinoma Cells In Vitro. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	6
88	Locally advanced mammary analogue secretory carcinoma of the parotid gland. <i>International Journal of Oral and Maxillofacial Surgery</i> , 2019 , 48, 865-868	2.9	5
87	Management of cN0 case in oral cancer(Current status and issues). <i>Journal of Japanese Society of Oral Oncology</i> , 2019 , 31, 137-142	0	1
86	Distinct Regulation of CXCL10 Production by Cytokines in Human Salivary Gland Ductal and Acinar Cells. <i>Inflammation</i> , 2018 , 41, 1172-1181	5.1	5
85	Determination of the origin of oral squamous cell carcinoma by microarray analysis: Squamous epithelium or minor salivary gland?. <i>International Journal of Cancer</i> , 2018 , 143, 2551-2560	7.5	6
84	Successful management of an oro-cutaneous fistula and exposed mandibular plate with the vacuum-assisted closure system: A case report. <i>Experimental and Therapeutic Medicine</i> , 2018 , 16, 5315-5317	2.1	1
83	Inverse correlation between the number of CXCR3 macrophages and the severity of inflammatory lesions in Sjögren's syndrome salivary glands: A pilot study. <i>Journal of Oral Pathology and Medicine</i> , 2018 , 47, 710-718	3.3	12
82	The CUL3-SPOP-DAXX axis is a novel regulator of VEGFR2 expression in vascular endothelial cells. <i>Scientific Reports</i> , 2017 , 7, 42845	4.9	20
81	A rare recurrent case of hypervascular juvenile ossifying fibroma. <i>Journal of Oral and Maxillofacial Surgery, Medicine, and Pathology</i> , 2017 , 29, 550-554	0.4	
80	Prognostic impact of preoperative serum interleukin-6 levels in patients with early-stage oral squamous cell carcinoma, defined by sentinel node biopsy. <i>Oncology Letters</i> , 2017 , 14, 7965-7969	2.6	4
79	Current status and tasks of sentinel node biopsy in oral cancer (Review Article). <i>Journal of Japanese Society of Oral Oncology</i> , 2016 , 28, 71-75	0	1
78	Comprehensive assessment of the prognosis of pancreatic cancer: peripheral blood neutrophil-lymphocyte ratio and immunohistochemical analyses of the tumour site. <i>Scandinavian Journal of Gastroenterology</i> , 2016 , 51, 610-7	2.4	31
77	Objective validity of an implant-retained overdenture with a ball attachment system after marginal mandibulectomy. <i>British Journal of Oral and Maxillofacial Surgery</i> , 2016 , 54, e21-5	1.4	5
76	Therapeutic potential of targeting cell division cycle associated 5 for oral squamous cell carcinoma. <i>Oncotarget</i> , 2016 , 7, 2343-53	3.3	27

75	Annexin A8 is a novel molecular marker for detecting lymph node metastasis in oral squamous cell carcinoma. <i>Oncotarget</i> , 2016 , 7, 4882-9	3.3	14
74	Reversible interconversion and maintenance of mammary epithelial cell characteristics by the ligand-regulated EGFR system. <i>Scientific Reports</i> , 2016 , 6, 20209	4.9	10
73	Identification of Akt1 as a potent therapeutic target for oral squamous cell carcinoma. <i>International Journal of Oncology</i> , 2015 , 47, 1273-81	4.4	10
72	Ribonucleotide reductase M2 is a promising molecular target for the treatment of oral squamous cell carcinoma. <i>International Journal of Oncology</i> , 2015 , 46, 1971-7	4.4	13
71	Pathophysiology of lung injury induced by common bile duct ligation in mice. <i>PLoS ONE</i> , 2014 , 9, e945503	3.7	18
70	Prognostic significance of interleukin-8 and CD163-positive cell-infiltration in tumor tissues in patients with oral squamous cell carcinoma. <i>PLoS ONE</i> , 2014 , 9, e110378	3.7	44
69	Immunochemoradiotherapy for patients with oral squamous cell carcinoma: augmentation of OK-432-induced helper T cell 1 response by 5-FU and X-ray irradiation. <i>Neoplasia</i> , 2013 , 15, 805-14	6.4	14
68	Prognostic impact of expression of Bcl-2 and Bax genes in circulating immune cells derived from patients with head and neck carcinoma. <i>Neoplasia</i> , 2013 , 15, 305-14	6.4	14
67	Targeting Aurora kinase A suppresses the growth of human oral squamous cell carcinoma cells in vitro and in vivo. <i>Oral Oncology</i> , 2013 , 49, 551-9	4.4	13
66	Possible negative effect of tocilizumab on BRONJ in a patient with methotrexate-associated lymphoproliferative disorder. <i>Nihon Koku Geka Gakkai Zasshi</i> , 2013 , 59, 346-351	0.1	7
65	Growth inhibition and apoptosis by an active component of OK-432, a streptococcal agent, via Toll-like receptor 4 in human head and neck cancer cell lines. <i>Oral Oncology</i> , 2012 , 48, 678-85	4.4	10
64	Premetastatic vasculogenesis in oral squamous cell carcinoma xenograft-draining lymph nodes. <i>Oral Oncology</i> , 2012 , 48, 663-70	4.4	7
63	One-step nucleic acid amplification for detecting lymph node metastasis of head and neck squamous cell carcinoma. <i>Oral Oncology</i> , 2012 , 48, 958-963	4.4	26
62	Human FAT1 cadherin controls cell migration and invasion of oral squamous cell carcinoma through the localization of E-catenin. <i>Oncology Reports</i> , 2011 , 26, 587-92	3.5	50
61	Gene therapy for oral squamous cell carcinoma with IAI.3B promoter-driven oncolytic adenovirus-infected carrier cells. <i>Oncology Reports</i> , 2011 , 25, 795-802	3.5	2
60	Anti-tumor effect of small interfering RNA targeting the androgen receptor in human androgen-independent prostate cancer cells. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 391, 1075-9	3.4	18
59	Conditional deletion of Stat3 promotes neurogenesis and inhibits astroglialogenesis in neural stem cells. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 394, 843-7	3.4	63
58	Knockdown of Akt isoforms by RNA silencing suppresses the growth of human prostate cancer cells in vitro and in vivo. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 399, 79-83	3.4	30

57	Spindle cell carcinoma after irradiated oral squamous cell carcinoma treated with S-1. <i>Asian Journal of Oral and Maxillofacial Surgery</i> , 2010 , 22, 175-179		
56	Carrier cell-mediated cell lysis of squamous cell carcinoma cells by squamous cell carcinoma antigen 1 promoter-driven oncolytic adenovirus. <i>Journal of Gene Medicine</i> , 2010 , 12, 545-54	3.5	7
55	CD151 regulates HGF-stimulated morphogenesis of human breast cancer cells. <i>Biochemical and Biophysical Research Communications</i> , 2009 , 379, 1097-100	3.4	25
54	18F-Fluorodeoxyglucose Positron Emission Tomography/Computed Tomography for Diagnosis of Cervical Lymph Node Metastases of Oral Squamous Cell Carcinoma. <i>Asian Journal of Oral and Maxillofacial Surgery</i> , 2009 , 21, 88-95		0
53	Methotrexate-induced malignant lymphoma in the maxilla of a patient with rheumatoid arthritis. <i>Nihon Koku Geka Gakkai Zasshi</i> , 2009 , 55, 463-466	0.1	7
52	Role of Akt isoforms in HGF-induced invasive growth of human salivary gland cancer cells. <i>Biochemical and Biophysical Research Communications</i> , 2008 , 370, 123-8	3.4	25
51	Detection of lymph node micrometastases in patients with squamous carcinoma of the head and neck. <i>European Archives of Oto-Rhino-Laryngology</i> , 2008 , 265, 1147-53	3.5	43
50	Basic evidence of molecular targeted therapy for oral cancer and salivary gland cancer. <i>Head and Neck</i> , 2008 , 30, 800-9	4.2	55
49	The Meaning of FDG PET-CT and Sentinel Lymph Node Biopsy in Patients with cN0 Oral Squamous Cell Carcinoma. <i>Japanese Journal of Head and Neck Cancer</i> , 2008 , 34, 513-517	0.1	1
48	Stat3 as a molecular target in RNA interference-based treatment of oral squamous cell carcinoma. <i>Oncology Reports</i> , 2008 , 20, 873-8	3.5	14
47	Akt1 Is a Potent Molecular Target for Treatment of Oral Cancer. <i>Journal of Oral and Maxillofacial Surgery</i> , 2007 , 65, 34.e7	1.8	
46	Hypoxia enhances c-Met/HGF receptor expression and signaling by activating HIF-1alpha in human salivary gland cancer cells. <i>Oral Oncology</i> , 2006 , 42, 593-8	4.4	54
45	Inactivation of AR activates HGF/c-Met system in human prostatic carcinoma cells. <i>Biochemical and Biophysical Research Communications</i> , 2006 , 347, 1158-65	3.4	29
44	CXCR4 expression is associated with lymph-node metastasis of oral squamous cell carcinoma 2006 , 28, 61		5
43	PLZF regulates Pbx1 transcription and Pbx1-HoxC8 complex leads to androgen-independent prostate cancer proliferation. <i>Prostate</i> , 2006 , 66, 1092-9	4.2	41
42	CD151 forms a functional complex with c-Met in human salivary gland cancer cells. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 336, 408-16	3.4	54
41	A case of metastatic tongue cancer with replacement of the common carotid artery. <i>Nihon Koku Geka Gakkai Zasshi</i> , 2005 , 51, 368-371	0.1	1
40	PERSONALIZATION OF TUMOR MARKERS IN ORAL CANCER. <i>Japanese Journal of Head and Neck Cancer</i> , 2005 , 31, 493-497	0.1	

39	A case of remitting seronegative symmetrical synovitis with pitting edema (RS3PE) induced by UFT in a patient with metastatic foci of unknown origin. <i>Nihon Koku Geka Gakkai Zasshi</i> , 2005 , 51, 148-151	0.1	
38	CLINICAL RESULTS OF SENTINEL LYMPH NODE (SN) BIOPSY FOR ORAL CANCER-RELATIONSHIP BETWEEN SN LOCALIZATION AND METASTASIS IN TONGUE CANCER-. <i>Japanese Journal of Head and Neck Cancer</i> , 2005 , 31, 79-83	0.1	
37	Overexpression of cyclooxygenase-2 is associated with radioresistance in oral squamous cell carcinoma. <i>Oral Oncology</i> , 2004 , 40, 383-9	4.4	46
36	Expression of vascular endothelial growth factor A, B, C, and D in oral squamous cell carcinoma. <i>Oral Oncology</i> , 2004 , 40, 13-20	4.4	94
35	Gefitinib (Iressa® ZD1839), an epidermal growth factor receptor tyrosine kinase inhibitor, up-regulates p27KIP1 and induces G1 arrest in oral squamous cell carcinoma cell lines. <i>Oral Oncology</i> , 2004 , 40, 43-51	4.4	39
34	Growth-regulated oncogene-1 expression is associated with angiogenesis and lymph node metastasis in human oral cancer. <i>Oncology</i> , 2004 , 66, 316-22	3.6	38
33	Phenotypic switch from paracrine to autocrine role of hepatocyte growth factor in an androgen-independent human prostatic carcinoma cell line, CWR22R. <i>American Journal of Pathology</i> , 2004 , 165, 533-40	5.8	35
32	HER2/neu Expression in Oral Squamous Cell Carcinoma. <i>Asian Journal of Oral and Maxillofacial Surgery</i> , 2004 , 16, 172-176		2
31	Constitutive activation of Stat3 correlates with increased expression of the c-Met/HGF receptor in oral squamous cell carcinoma. <i>Oncology Reports</i> , 2004 , 12, 293	3.5	2
30	Basic and clinical studies on quantitative analysis of lymph node micrometastasis in oral cancer. <i>Oncology Reports</i> , 2004 , 11, 33-9	3.5	13
29	Evaluation of the chemosensitivity of head and neck cancer cells based on the diverse function of mutated-p53 2003 , 22, 383		
28	Vesnarinone: a differentiation-inducing anti-cancer drug. <i>Anti-Cancer Drugs</i> , 2003 , 14, 391-5	2.4	6
27	Enhancement of radiosensitivity in head and neck cancer cells by ZD1839 (IRESSA®), a selective epidermal growth factor receptor tyrosine kinase inhibitor. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 2003 , 26, e150-6	2.7	18
26	Skp2 and Jab1 expression are associated with inverse expression of p27(KIP1) and poor prognosis in oral squamous cell carcinomas. <i>Oncology</i> , 2003 , 65, 355-62	3.6	68
25	Immunohistochemical expression of hepatocyte growth factor and c-Met/HGF receptor in benign and malignant human prostate tissue. <i>Oncology Reports</i> , 2003 , 10, 1149	3.5	6
24	Flavopiridol, a cyclin dependent kinase (CDK) inhibitor, induces apoptosis by regulating Bcl-x in oral cancer cells. <i>Oral Oncology</i> , 2003 , 39, 49-55	4.4	20
23	Thiazolidinediones inhibit cell growth of human oral squamous cell carcinoma in vitro independent of peroxisome proliferator-activated receptor gamma. <i>Oral Oncology</i> , 2003 , 39, 855-61	4.4	15
22	Enhancement of tumor radioresponse by combined treatment with gefitinib (Iressa, ZD1839), an epidermal growth factor receptor tyrosine kinase inhibitor, is accompanied by inhibition of DNA damage repair and cell growth in oral cancer. <i>International Journal of Cancer</i> , 2003 , 107, 1030-7	7.5	78

21	Up-regulation of DNA-dependent protein kinase correlates with radiation resistance in oral squamous cell carcinoma. <i>Cancer Science</i> , 2003 , 94, 894-900	6.9	98
20	Posttranscriptional regulation of TSC-22 (TGF-beta-stimulated clone-22) gene by TGF-beta 1. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 305, 846-54	3.4	19
19	Possible role of stromal-cell-derived factor-1/CXCR4 signaling on lymph node metastasis of oral squamous cell carcinoma. <i>Experimental Cell Research</i> , 2003 , 290, 289-302	4.2	146
18	Gefitinib (Iressa), an epidermal growth factor receptor tyrosine kinase inhibitor, mediates the inhibition of lymph node metastasis in oral cancer cells. <i>Cancer Letters</i> , 2003 , 201, 149-55	9.9	40
17	P27KIP1, Skp2 AND JAB1 EXPRESSION AND THEIR CLINICAL IMPLICATION IN ORAL SQUAMOUS CELL CARCINOMAS. <i>Japanese Journal of Head and Neck Cancer</i> , 2003 , 29, 210-216		1
16	Process and present status of radioisotope-guided sentinel lymph node biopsy at our department. <i>Nihon Koku Geka Gakkai Zasshi</i> , 2003 , 49, 257-263	0.1	2
15	INTRAOPERATIVE RAPID DETECTION OF MICROMETASTASIS IN SENTINEL LYMPH NODES OF ORAL MALIGNANT TUMORS. <i>Japanese Journal of Head and Neck Cancer</i> , 2003 , 29, 64-69		3
14	Two cases of habitual temporomandibular dislocation surgically treated by the Buckley-Terry procedure using a T-type titanium miniplate. <i>Nihon Koku Geka Gakkai Zasshi</i> , 2003 , 49, 544-547	0.1	
13	Evaluation of the chemosensitivity of head and neck cancer cells based on the diverse function of mutated-p53. <i>International Journal of Oncology</i> , 2003 , 22, 383-9	1	3
12	Immunohistochemical expression of hepatocyte growth factor and c-Met/HGF receptor in benign and malignant human prostate tissue. <i>Oncology Reports</i> , 2003 , 10, 1149-53	3.5	25
11	Expression of peroxisome proliferator-activated receptor γ and the growth inhibitory effect of its synthetic ligands in human salivary gland cancer cell lines 2002 , 20, 599		
10	Infiltration of tumor-associated macrophages in human oral squamous cell carcinoma. <i>Oncology Reports</i> , 2002 , 9, 1219	3.5	7
9	EVALUATION OF THE CHEMOSENSITIVITY OF HEAD AND NECK CANCER BASED ON DIVERSE FUNCTION OF MUTATED-p53. <i>Japanese Journal of Head and Neck Cancer</i> , 2002 , 28, 264-268		
8	Dysfunction of the p53 tumor suppressor pathway in head and neck cancer. <i>International Journal of Oncology</i> , 2002 , 21, 119-26	1	7
7	Androgen receptor expression in androgen-independent prostate cancer cell lines. <i>Prostate</i> , 2001 , 47, 66-75	4.2	97
6	Role of HGF/c-met system in invasion and metastasis of oral squamous cell carcinoma cells in vitro and its clinical significance. <i>International Journal of Cancer</i> , 2001 , 93, 489-96	7.5	80
5	Role of peroxisome proliferator-activated receptor γ and its ligands in non-neoplastic and neoplastic human urothelial cells. <i>American Journal of Pathology</i> , 2001 , 159, 591-7	5.8	73
4	Two cases of Bell's palsy responding to treatment with the antiviral agent acyclovir.. <i>Nihon Koku Geka Gakkai Zasshi</i> , 2001 , 47, 695-698	0.1	

- | | | | |
|---|--|-----|----|
| 3 | Hepatocyte growth factor secreted by prostate-derived stromal cells stimulates growth of androgen-independent human prostatic carcinoma cells. <i>American Journal of Pathology</i> , 2000 , 157, 795-803 | 58 | 43 |
| 2 | Possible contribution of active MMP2 to lymph-node metastasis and secreted cathepsin L to bone invasion of newly established human oral-squamous-cancer cell lines. <i>International Journal of Cancer</i> , 1997 , 70, 120-7 | 7.5 | 93 |
| 1 | Basic and clinical studies on quantitative analysis of lymph node micrometastasis in oral cancer. <i>Oncology Reports</i> , | 3.5 | 4 |