Scott M Stevens

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7175424/publications.pdf

Version: 2024-02-01

64 papers

7,169 citations

236833 25 h-index 57 g-index

67 all docs

67
docs citations

67 times ranked

8143 citing authors

| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Antithrombotic Therapy for VTE Disease. Chest, 2016, 149, 315-352. | 0.4 | 4,060 |
| 2 | A Pharmacogenetic versus a Clinical Algorithm for Warfarin Dosing. New England Journal of Medicine, 2013, 369, 2283-2293. | 13.9 | 660 |
| 3 | Diagnosis of DVT. Chest, 2012, 141, e351S-e418S. | 0.4 | 570 |
| 4 | Guidance for the evaluation and treatment of hereditary and acquired thrombophilia. Journal of Thrombosis and Thrombolysis, 2016, 41, 154-164. | 1.0 | 230 |
| 5 | Time outside of therapeutic range in atrial fibrillation patients is associated with long-term risk of dementia. Heart Rhythm, 2014, 11, 2206-2213. | 0.3 | 130 |
| 6 | Long-Term Population-Based Cerebral Ischemic Event and Cognitive Outcomes of Direct Oral Anticoagulants Compared With Warfarin Among Long-term Anticoagulated Patients for Atrial Fibrillation. American Journal of Cardiology, 2016, 118, 210-214. | 0.7 | 123 |
| 7 | Withholding Anticoagulation after a Negative Result on Duplex Ultrasonography for Suspected Symptomatic Deep Venous Thrombosis. Annals of Internal Medicine, 2004, 140, 985. | 2.0 | 123 |
| 8 | Radiation and Chest CT Scan Examinations. Chest, 2012, 142, 750-760. | 0.4 | 110 |
| 9 | Reduction of Peripherally Inserted Central Catheter-Associated DVT. Chest, 2013, 143, 627-633. | 0.4 | 89 |
| 10 | Physician Alerts to Prevent Symptomatic Venous Thromboembolism in Hospitalized Patients. Circulation, 2009, 119, 2196-2201. | 1.6 | 88 |
| 11 | Follow-up of Incidental Pulmonary Nodules and the Radiology Report. Journal of the American College of Radiology, 2014, 11, 378-383. | 0.9 | 86 |
| 12 | Apixaban for the Secondary Prevention of Thrombosis Among Patients With Antiphospholipid Syndrome. Clinical and Applied Thrombosis/Hemostasis, 2016, 22, 239-247. | 0.7 | 85 |
| 13 | Adherence to PIOPED II Investigators' Recommendations for Computed Tomography Pulmonary Angiography. American Journal of Medicine, 2013, 126, 36-42. | 0.6 | 70 |
| 14 | Antiphospholipid antibodies and recurrent thrombosis after a first unprovoked venous thromboembolism. Blood, 2018, 131, 2151-2160. | 0.6 | 62 |
| 15 | Management of Low-Risk Pulmonary Embolism Patients Without Hospitalization. Chest, 2018, 154, 249-256. | 0.4 | 60 |
| 16 | Apixaban compared with warfarin to prevent thrombosis in thrombotic antiphospholipid syndrome: a randomized trial. Blood Advances, 2022, 6, 1661-1670. | 2.5 | 56 |
| 17 | Atrial Fibrillation Patients Treated With Longâ€Term Warfarin Anticoagulation Have Higher Rates of All Dementia Types Compared With Patients Receiving Longâ€Term Warfarin for Other Indications. Journal of the American Heart Association, 2016, 5, . | 1.6 | 43 |
| 18 | Atrial fibrillation incrementally increases dementia risk across all CHADS 2 and CHA 2 DS 2 VASc strata in patients receiving long-term warfarin. American Heart Journal, 2017, 188, 93-98. | 1.2 | 41 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Protocol Modification of Apixaban for the Secondary Prevention of Thrombosis Among Patients With Antiphospholipid Syndrome Study. Clinical and Applied Thrombosis/Hemostasis, 2018, 24, 192-192. | 0.7 | 41 |
| 20 | Percent Time With a Supratherapeutic INR in Atrial Fibrillation Patients Also Using an Antiplatelet Agent Is Associated With Longâ€Term Risk of Dementia. Journal of Cardiovascular Electrophysiology, 2015, 26, 1180-1186. | 0.8 | 40 |
| 21 | The Michigan Appropriateness Guide for Intravenous Catheters (MAGIC) initiative: A summary and review of peripherally inserted central catheter and venous catheter appropriate use. Journal of Hospital Medicine, 2016, 11, 306-310. | 0.7 | 36 |
| 22 | Longâ€term risk of recurrence in patients with a first unprovoked venous thromboembolism managed according to dâ€dimer results; A cohort study. Journal of Thrombosis and Haemostasis, 2019, 17, 1144-1152. | 1.9 | 34 |
| 23 | Assessment of the Safety and Efficiency of Using an Age-Adjusted D-dimer Threshold to Exclude Suspected Pulmonary Embolism. Chest, 2014, 146, 1444-1451. | 0.4 | 32 |
| 24 | Major Bleeding With Dabigatran and Rivaroxaban in Patients With Atrial Fibrillation. Clinical and Applied Thrombosis/Hemostasis, 2014, 20, 665-672. | 0.7 | 29 |
| 25 | Follow-up of Incidental Pulmonary Nodules and the Radiology Report. Journal of the American College of Radiology, 2016, 13, R18-R24. | 0.9 | 28 |
| 26 | Rationale and design of the impact of anticoagulation therapy on the Cognitive Decline and Dementia in Patients with Nonvalvular Atrial Fibrillation (CAF) Trial: A Vanguard study. Clinical Cardiology, 2019, 42, 506-512. | 0.7 | 18 |
| 27 | Deep Vein Thrombosis Prophylaxis in Hospitalized Medical Patients: Current Recommendations, General Rates of Implementation, and Initiatives for Improvement. Clinics in Chest Medicine, 2010, 31, 675-689. | 0.8 | 16 |
| 28 | Venous Thromboembolism in Critically Ill Medical Patients Receiving Chemoprophylaxis. Clinical and Applied Thrombosis/Hemostasis, 2016, 22, 265-273. | 0.7 | 16 |
| 29 | Computer surveillance of patients at high risk for and with venous thromboembolism. AMIA Annual Symposium proceedings, 2010, 2010, 217-21. | 0.2 | 16 |
| 30 | Electronic Alerts, Comparative Practitioner Metrics, and Education Improves Thromboprophylaxis and Reduces Thrombosis. American Journal of Medicine, 2016, 129, 1124.e17-1124.e26. | 0.6 | 15 |
| 31 | Comparative thrombosis risk of vascular access devices among critically ill medical patients. Thrombosis Research, 2018, 172, 54-60. | 0.8 | 15 |
| 32 | Natural Language Processing Performance for the Identification of Venous Thromboembolism in an Integrated Healthcare System. Clinical and Applied Thrombosis/Hemostasis, 2021, 27, 107602962110131. | 0.7 | 15 |
| 33 | Weight-based enoxaparin dosing and deep vein thrombosis in hospitalized trauma patients: A double-blind, randomized, pilot study. Surgery, 2018, 164, 144-149. | 1.0 | 14 |
| 34 | Age-adjusted versus clinical probability-adjusted D-dimer to exclude pulmonary embolism. Thrombosis Research, 2018, 167, 15-19. | 0.8 | 14 |
| 35 | Computerized Clinical Decision Support Improves Warfarin Management and Decreases Recurrent Venous Thromboembolism. Clinical and Applied Thrombosis/Hemostasis, 2015, 21, 197-203. | 0.7 | 13 |
| 36 | Electronic alerts, comparative practitioner metrics, and education improve thromboprophylaxis and reduce venous thrombosis in community hospitals. Research and Practice in Thrombosis and Haemostasis, 2018, 2, 481-489. | 1.0 | 12 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Effect of Low-Intensity vs Standard-Intensity Warfarin Prophylaxis on Venous Thromboembolism or Death Among Patients Undergoing Hip or Knee Arthroplasty. JAMA - Journal of the American Medical Association, 2019, 322, 834. | 3.8 | 9 |
| 38 | The use of a fixed high sensitivity to evaluate five D-dimer assays' ability to rule out deep venous thrombosis: a novel approach. British Journal of Haematology, 2005, 131, 341-347. | 1.2 | 8 |
| 39 | Withholding Anticoagulation Following a Single Negative Whole-Leg Ultrasound in Patients at High Pretest Probability for Deep Vein Thrombosis. Clinical and Applied Thrombosis/Hemostasis, 2013, 19, 79-85. | 0.7 | 8 |
| 40 | Thrombophilic Evaluation in Patients with Acute Pulmonary Embolism. Seminars in Respiratory and Critical Care Medicine, 2017, 38, 107-120. | 0.8 | 8 |
| 41 | Diagnosing deep vein thrombosis in cancer patients with suspected symptoms: An individual participant data metaâ€analysis. Journal of Thrombosis and Haemostasis, 2020, 18, 2245-2252. | 1.9 | 6 |
| 42 | Preemptive Anticoagulation in Patients With a High Pretest Probability of Pulmonary Embolism. Chest, 2018, 153, 1153-1159. | 0.4 | 5 |
| 43 | Depression as a Driving Force for Low Time in Therapeutic Range and Dementia in Patients With and Without Atrial Fibrillation. American Journal of Cardiology, 2021, 153, 58-64. | 0.7 | 5 |
| 44 | Intermittent pneumatic compression in patients with stroke. Lancet, The, 2013, 382, 484-486. | 6.3 | 4 |
| 45 | Role of thrombophilia testing: con. Journal of Thrombosis and Thrombolysis, 2015, 39, 379-391. | 1.0 | 4 |
| 46 | The Population-Based Long-Term Impact of Anticoagulant and Antiplatelet Therapies in Low-Risk Patients With Atrial Fibrillation. American Journal of Cardiology, 2017, 120, 75-82. | 0.7 | 4 |
| 47 | Postâ€discharge thrombosis and bleeding in medical patients: A novel risk score derived from ubiquitous biomarkers. Research and Practice in Thrombosis and Haemostasis, 2021, 5, e12560. | 1.0 | 4 |
| 48 | Review: the Wells rule is more useful than individual clinical features for predicting risk of deep venous thrombosis. Evidence-Based Medicine, 2006, 11, 56-56. | 0.6 | 3 |
| 49 | New evidence on old drugs; warfarin versus aspirin after bioprosthetic aortic valve placement. Thrombosis Research, 2017, 150, 102-103. | 0.8 | 2 |
| 50 | Predicting postdischarge hospitalâ€associated venous thromboembolism among medical patients using a validated mortality risk score derived from common biomarkers. Research and Practice in Thrombosis and Haemostasis, 2020, 4, 872-878. | 1.0 | 2 |
| 51 | The Wells rule was not useful in ruling out deep venous thrombosis in a primary care setting. Evidence-Based Medicine, 2006, 11, 57-57. | 0.6 | 1 |
| 52 | Review: Gestalt or clinical decision rules have limited sensitivity and specificity for detecting acute PE. Annals of Internal Medicine, 2012, 156, JC1. | 2.0 | 1 |
| 53 | Timing of parenteral anticoagulation after thrombolysis for the treatment of pulmonary embolism. Thrombosis Research, 2020, 195, 58-61. | 0.8 | 1 |
| 54 | Something old, something new…. Blood, 2020, 135, 1307-1308. | 0.6 | 1 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Safety of Excluding Suspected Deep Vein Thrombosis with a Single Whole-Leg Compression Ultrasound: Systematic Review and Meta-Analysis Blood, 2009, 114, 243-243. | 0.6 | 1 |
| 56 | Single Complete Compression Ultrasonography for Suspected Deep Venous Thrombosis: Ideal in Routine Clinical Practice?. Annals of Internal Medicine, 2004, 141, 889. | 2.0 | 1 |
| 57 | Review: The Wells rule is more useful than individual clinical features for predicting risk for deep venous thrombosis. ACP Journal Club, 2006, 144, 46. | 0.1 | 1 |
| 58 | Response to Letter Regarding Article, "Physician Alerts to Prevent Symptomatic Venous Thromboembolism in Hospitalized Patients― Circulation, 2009, 120, . | 1.6 | 0 |
| 59 | Ultrasound of the Whole Arm to Manage Suspected Upper-Extremity Deep Venous Thrombosis. JAMA Internal Medicine, 2015, 175, 1227. | 2.6 | 0 |
| 60 | The Wells rule was not useful in ruling out deep venous thrombosis in a primary care setting. ACP Journal Club, 2006, 144, 47. | 0.1 | 0 |
| 61 | Concomitant VTE increased risks for mortality and hemorrhage in older patients with cancer, with risk varying by cancer type. ACP Journal Club, 2007, 147, 51. | 0.1 | 0 |
| 62 | Review: the Wells rule is more useful than individual clinical features for predicting risk for deep venous thrombosis. ACP Journal Club, 2006, 144, 46-7. | 0.1 | 0 |
| 63 | The Wells rule was not useful in ruling out deep venous thrombosis in a primary care setting. ACP Journal Club, 2006, 144, 46-7. | 0.1 | 0 |
| 64 | Concomitant VTE increased risks for mortality and hemorrhage in older patients with cancer, with risk varying by cancer type. ACP Journal Club, 2007, 147, 51. | 0.1 | 0 |