

Kazuyuki Tobe

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7188341/kazuyuki-tobe-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

121
papers

5,563
citations

35
h-index

73
g-index

131
ext. papers

6,527
ext. citations

6
avg, IF

5.08
L-index

#	Paper	IF	Citations
121	Insulin resistance and growth retardation in mice lacking insulin receptor substrate-1. <i>Nature</i> , 1994 , 372, 182-6	50.4	914
120	Overexpression of monocyte chemoattractant protein-1 in adipose tissues causes macrophage recruitment and insulin resistance. <i>Journal of Biological Chemistry</i> , 2006 , 281, 26602-14	5.4	638
119	Regulatory mechanisms for adipose tissue M1 and M2 macrophages in diet-induced obese mice. <i>Diabetes</i> , 2009 , 58, 2574-82	0.9	530
118	Impaired insulin signaling in endothelial cells reduces insulin-induced glucose uptake by skeletal muscle. <i>Cell Metabolism</i> , 2011 , 13, 294-307	24.6	298
117	Tyrosine phosphorylation of the EGF receptor by the kinase Jak2 is induced by growth hormone. <i>Nature</i> , 1997 , 390, 91-6	50.4	252
116	Insulin receptor substrate 2 plays a crucial role in beta cells and the hypothalamus. <i>Journal of Clinical Investigation</i> , 2004 , 114, 917-27	15.9	187
115	Efficacy and safety of monotherapy with the novel sodium/glucose cotransporter-2 inhibitor tofogliflozin in Japanese patients with type 2 diabetes mellitus: a combined Phase 2 and 3 randomized, placebo-controlled, double-blind, parallel-group comparative study. <i>Cardiovascular Diabetology</i> , 2014 , 13, 65	8.7	135
114	SIRT1-Mediated eNAMPT Secretion from Adipose Tissue Regulates Hypothalamic NAD ⁺ and Function in Mice. <i>Cell Metabolism</i> , 2015 , 21, 706-17	24.6	129
113	Treatment with SRT1720, a SIRT1 activator, ameliorates fatty liver with reduced expression of lipogenic enzymes in MSG mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2009 , 297, E1179-86	6	125
112	CD206 M2-like macrophages regulate systemic glucose metabolism by inhibiting proliferation of adipocyte progenitors. <i>Nature Communications</i> , 2017 , 8, 286	17.4	116
111	Increased expression of the sterol regulatory element-binding protein-1 gene in insulin receptor substrate-2(-/-) mouse liver. <i>Journal of Biological Chemistry</i> , 2001 , 276, 38337-40	5.4	104
110	Genome-wide association studies in the Japanese population identify seven novel loci for type 2 diabetes. <i>Nature Communications</i> , 2016 , 7, 10531	17.4	99
109	Telmisartan improves insulin resistance and modulates adipose tissue macrophage polarization in high-fat-fed mice. <i>Endocrinology</i> , 2011 , 152, 1789-99	4.8	79
108	Implications of altered NAD metabolism in metabolic disorders. <i>Journal of Biomedical Science</i> , 2019 , 26, 34	13.3	77
107	Japanese Clinical Practice Guideline for Diabetes 2019. <i>Diabetology International</i> , 2020 , 11, 165-223	2.3	75
106	Vascular endothelial growth factor induces activation and subcellular translocation of focal adhesion kinase (p125FAK) in cultured rat cardiac myocytes. <i>Circulation Research</i> , 1999 , 84, 1194-202	15.7	73
105	Genetic risk score constructed using 14 susceptibility alleles for type 2 diabetes is associated with the early onset of diabetes and may predict the future requirement of insulin injections among Japanese individuals. <i>Diabetes Care</i> , 2012 , 35, 1763-70	14.6	71

104	Serum levels of vascular endothelial growth factor in patients with acute myocardial infarction undergoing reperfusion therapy. <i>Clinical Science</i> , 1997 , 92, 453-4	6.5	67
103	Identification of a 190-kDa protein as a novel substrate for the insulin receptor kinase functionally similar to insulin receptor substrate-1. <i>Journal of Biological Chemistry</i> , 1995 , 270, 5698-701	5.4	66
102	The RNA Methyltransferase Complex of WTAP, METTL3, and METTL14 Regulates Mitotic Clonal Expansion in Adipogenesis. <i>Molecular and Cellular Biology</i> , 2018 , 38,	4.8	65
101	Long-term safety and efficacy of tofogliflozin, a selective inhibitor of sodium-glucose cotransporter 2, as monotherapy or in combination with other oral antidiabetic agents in Japanese patients with type 2 diabetes mellitus: multicenter, open-label, randomized controlled trials. <i>Expert Opinion on Pharmacotherapy</i> , 2014 , 15, 749-66	4	65
100	HIF-1 α in Myeloid Cells Promotes Adipose Tissue Remodeling Toward Insulin Resistance. <i>Diabetes</i> , 2016 , 65, 3649-3659	0.9	59
99	Signal transduction mechanism of insulin and insulin-like growth factor-1. <i>Endocrine Journal</i> , 1996 , 43 Suppl, S33-41	2.9	59
98	In vivo depletion of CD206+ M2 macrophages exaggerates lung injury in endotoxemic mice. <i>American Journal of Pathology</i> , 2015 , 185, 162-71	5.8	58
97	Mediator complex recruits epigenetic regulators via its two cyclin-dependent kinase subunits to repress transcription of immune response genes. <i>Journal of Biological Chemistry</i> , 2013 , 288, 20955-20965	5.4	56
96	A subset of cerebrovascular pericytes originates from mature macrophages in the very early phase of vascular development in CNS. <i>Scientific Reports</i> , 2017 , 7, 3855	4.9	52
95	Both insulin signaling defects in the liver and obesity contribute to insulin resistance and cause diabetes in <i>Irs2</i> ($-/-$) mice. <i>Journal of Biological Chemistry</i> , 2004 , 279, 25039-49	5.4	50
94	Growth hormone-induced tyrosine phosphorylation of EGF receptor as an essential element leading to MAP kinase activation and gene expression. <i>Endocrine Journal</i> , 1998 , 45 Suppl, S27-31	2.9	47
93	Sirtuin1 Maintains Actin Cytoskeleton by Deacetylation of Cortactin in Injured Podocytes. <i>Journal of the American Society of Nephrology: JASN</i> , 2015 , 26, 1939-59	12.7	46
92	Japanese Clinical Practice Guideline for Diabetes 2019. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1020-1076	10.76	46
91	Nmnat3 Is Dispensable in Mitochondrial NAD Level Maintenance In Vivo. <i>PLoS ONE</i> , 2016 , 11, e0147037	3.7	43
90	Expression of DGAT2 in white adipose tissue is regulated by central leptin action. <i>Journal of Biological Chemistry</i> , 2005 , 280, 3331-7	5.4	42
89	Vascular endothelial growth factor (VEGF) activates Raf-1, mitogen-activated protein (MAP) kinases, and S6 kinase (p90rsk) in cultured rat cardiac myocytes. <i>Journal of Cellular Physiology</i> , 1998 , 175, 239-46	7	39
88	Respiratory involvement in IgG4-related Mikulicz's disease. <i>Modern Rheumatology</i> , 2012 , 22, 31-39	3.3	37
87	CD103 T cells constrain lung fibrosis induced by CD103 tissue-resident pathogenic CD4 T cells. <i>Nature Immunology</i> , 2019 , 20, 1469-1480	19.1	36

86	The mechanism of insulin-induced signal transduction mediated by the insulin receptor substrate family. <i>Endocrine Journal</i> , 1999 , 46, S25-34	2.9	35
85	CD11c+ M1-like macrophages (M ₁) but not CD206+ M2-like M ₂ are involved in folliculogenesis in mice ovary. <i>Scientific Reports</i> , 2018 , 8, 8171	4.9	32
84	Astaxanthin stimulates mitochondrial biogenesis in insulin resistant muscle via activation of AMPK pathway. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2020 , 11, 241-258	10.3	31
83	Bofutsushosan improves gut barrier function with a bloom of <i>Akkermansia muciniphila</i> and improves glucose metabolism in mice with diet-induced obesity. <i>Scientific Reports</i> , 2020 , 10, 5544	4.9	28
82	Hepatic Sdf2l1 controls feeding-induced ER stress and regulates metabolism. <i>Nature Communications</i> , 2019 , 10, 947	17.4	28
81	Safety and efficacy of ipragliflozin in Japanese patients with type 2 diabetes in real-world clinical practice: interim results of the STELLA-LONG TERM post-marketing surveillance study. <i>Expert Opinion on Pharmacotherapy</i> , 2018 , 19, 189-201	4	27
80	Monoclonal antibody against citrullinated peptides obtained from rheumatoid arthritis patients reacts with numerous citrullinated microbial and food proteins. <i>Arthritis and Rheumatology</i> , 2015 , 67, 2020-31	9.5	26
79	FTO Gene Polymorphism Is Associated with Type 2 Diabetes through Its Effect on Increasing the Maximum BMI in Japanese Men. <i>PLoS ONE</i> , 2016 , 11, e0165523	3.7	24
78	Linkage of CD8 T cell exhaustion with high-fat diet-induced tumorigenesis. <i>Scientific Reports</i> , 2019 , 9, 12284	4.9	23
77	Baseline characteristics and interim (3-month) efficacy and safety data from STELLA-LONG TERM, a long-term post-marketing surveillance study of ipragliflozin in Japanese patients with type 2 diabetes in real-world clinical practice. <i>Expert Opinion on Pharmacotherapy</i> , 2016 , 17, 1985-94	4	21
76	Overexpression of Nmnat3 efficiently increases NAD and NGD levels and ameliorates age-associated insulin resistance. <i>Aging Cell</i> , 2018 , 17, e12798	9.9	19
75	Secretory units of islets in transplantation index is a useful predictor of insulin requirement in Japanese type 2 diabetic patients. <i>Journal of Diabetes Investigation</i> , 2014 , 5, 570-80	3.9	19
74	Mesenchymal stromal cells in bone marrow express adiponectin and are efficiently targeted by an adiponectin promoter-driven Cre transgene. <i>International Immunology</i> , 2019 , 31, 729-742	4.9	18
73	Partial depletion of CD206-positive M2-like macrophages induces proliferation of beige progenitors and enhances browning after cold stimulation. <i>Scientific Reports</i> , 2018 , 8, 14567	4.9	18
72	Effect of ipragliflozin on liver function in Japanese type 2 diabetes mellitus patients: a subgroup analysis of the STELLA-LONG TERM study (3-month interim results). <i>Endocrine Journal</i> , 2019 , 66, 31-41	2.9	17
71	Differential effects of DNA tumor virus nuclear oncogene products on adipocyte differentiation. <i>FEBS Letters</i> , 1987 , 215, 345-9	3.8	16
70	Proteomic analysis of serum biomarkers for prediabetes using the Long-Evans Agouti rat, a spontaneous animal model of type 2 diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2017 , 8, 661-673	3.9	15
69	SIRT1 and FOXO1 mRNA expression in PBMC correlates to physical activity in COPD patients. <i>International Journal of COPD</i> , 2017 , 12, 3237-3244	3	14

68	Elevated levels of plasma lactate dehydrogenase is an unfavorable prognostic factor in patients with epidermal growth factor receptor mutation-positive non-small cell lung cancer, receiving treatment with gefitinib or erlotinib. <i>Molecular and Clinical Oncology</i> , 2016 , 4, 774-778	1.6	13
67	Insulin- and Lipopolysaccharide-Mediated Signaling in Adipose Tissue Macrophages Regulates Postprandial Glycemia through Akt-mTOR Activation. <i>Molecular Cell</i> , 2020 , 79, 43-53.e4	17.6	12
66	Safety and Effectiveness of Ipragliflozin for Type 2 Diabetes in Japan: 12-Month Interim Results of the STELLA-LONG TERM Post-Marketing Surveillance Study. <i>Advances in Therapy</i> , 2019 , 36, 923-949	4.1	12
65	Astaxanthin Ameliorated Parvalbumin-Positive Neuron Deficits and Alzheimer's Disease-Related Pathological Progression in the Hippocampus of Mice. <i>Frontiers in Pharmacology</i> , 2020 , 11, 307	5.6	11
64	M2 macrophages in metabolism. <i>Diabetology International</i> , 2016 , 7, 342-351	2.3	11
63	M2-like macrophages serve as a niche for adipocyte progenitors in adipose tissue. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 1394-1400	3.9	11
62	Inactivation of DNA-dependent protein kinase promotes heat-induced apoptosis independently of heat-shock protein induction in human cancer cell lines. <i>PLoS ONE</i> , 2013 , 8, e58325	3.7	11
61	Semaglutide once a week in adults with overweight or obesity, with or without type 2 diabetes in an east Asian population (STEP 6): a randomised, double-blind, double-dummy, placebo-controlled, phase 3a trial.. <i>Lancet Diabetes and Endocrinology</i> , 2022 ,	18.1	11
60	Plasma neuron-specific enolase level as a prognostic marker in patients with non-small cell lung cancer receiving gefitinib. <i>Molecular and Clinical Oncology</i> , 2015 , 3, 802-806	1.6	10
59	Safety and efficacy of ipragliflozin in elderly versus non-elderly Japanese patients with type 2 diabetes mellitus: a subgroup analysis of the STELLA-LONG TERM study. <i>Expert Opinion on Pharmacotherapy</i> , 2018 , 19, 327-336	4	9
58	Sodium-glucose cotransporter 2 inhibitor, tofogliflozin, shows better improvements of blood glucose and insulin secretion in patients with high insulin levels at baseline. <i>Journal of Diabetes Investigation</i> , 2018 , 9, 862-869	3.9	9
57	Identifying pre-disease signals before metabolic syndrome in mice by dynamical network biomarkers. <i>Scientific Reports</i> , 2019 , 9, 8767	4.9	9
56	Replication Study in a Japanese Population to Evaluate the Association between 10 SNP Loci, Identified in European Genome-Wide Association Studies, and Type 2 Diabetes. <i>PLoS ONE</i> , 2015 , 10, e0126363	3.7	9
55	Reduced serum level of leukocyte cell-derived chemotaxin 2 is associated with the presence of diabetic retinopathy. <i>Clinica Chimica Acta</i> , 2016 , 463, 145-149	6.2	8
54	Sirt1 activator induces proangiogenic genes in preadipocytes to rescue insulin resistance in diet-induced obese mice. <i>Scientific Reports</i> , 2018 , 8, 11370	4.9	8
53	Replication Study in a Japanese Population of Six Susceptibility Loci for Type 2 Diabetes Originally Identified by a Transethnic Meta-Analysis of Genome-Wide Association Studies. <i>PLoS ONE</i> , 2016 , 11, e0154093	3.7	8
52	NAD+ Metabolism Regulates Preadipocyte Differentiation by Enhancing [Ketoglutarate-Mediated Histone H3K9 Demethylation at the PPAR[Promoter. <i>Frontiers in Cell and Developmental Biology</i> , 2020 , 8, 586179	5.7	8
51	Urinary afamin levels are associated with the progression of diabetic nephropathy. <i>Diabetes Research and Clinical Practice</i> , 2019 , 147, 37-46	7.4	8

50	CD206+ M2-Like Macrophages Are Essential for Successful Implantation. <i>Frontiers in Immunology</i> , 2020 , 11, 557184	8.4	7
49	Real-World Evidence for Long-Term Safety and Effectiveness of Ipragliflozin in Japanese Patients with Type 2 Diabetes Mellitus: final Results of a 3-Year Post-Marketing Surveillance Study (STELLA-LONG TERM). <i>Expert Opinion on Pharmacotherapy</i> , 2021 , 22, 373-387	4	7
48	Deletion of PHGDH in adipocytes improves glucose intolerance in diet-induced obese mice. <i>Biochemical and Biophysical Research Communications</i> , 2018 , 504, 309-314	3.4	7
47	Deletion of SIRT1 in myeloid cells impairs glucose metabolism with enhancing inflammatory response to adipose tissue hypoxia. <i>Diabetology International</i> , 2016 , 7, 59-68	2.3	6
46	Ratio of low molecular weight serum adiponectin to the total adiponectin value is associated with type 2 diabetes through its relation to increasing insulin resistance. <i>PLoS ONE</i> , 2018 , 13, e0192609	3.7	6
45	Outcome and Prognostic Factors in Patients with Small Cell Lung Cancer who Receive Third-line Chemotherapy. <i>Tumori</i> , 2014 , 100, 507-511	1.7	6
44	Roles of Insulin Receptor Substrate-1 and Shc on Insulin-Like Growth Factor I Receptor Signaling in Early Passages of Cultured Human Fibroblasts		6
43	Stimulation of the phosphorylation of cytoskeletal 350-kDa and 300-kDa proteins by insulin-like growth factor-I, platelet-derived growth factor and phorbol ester in rat 3Y1 cells. <i>Cell Structure and Function</i> , 1988 , 13, 417-23	2.2	6
42	Lactate dehydrogenase and body mass index are prognostic factors in patients with recurrent small cell lung cancer receiving amrubicin. <i>Tumori</i> , 2016 , 102, 606-609	1.7	6
41	Pioglitazone Ameliorates Smooth Muscle Cell Proliferation in Cuff-Induced Neointimal Formation by Both Adiponectin-Dependent and -Independent Pathways. <i>Scientific Reports</i> , 2016 , 6, 34707	4.9	5
40	Safety and effectiveness of ipragliflozin in elderly versus non-elderly Japanese type 2 diabetes mellitus patients: 12 month interim results of the STELLA-LONG TERM study. <i>Current Medical Research and Opinion</i> , 2019 , 35, 1901-1910	2.5	5
39	Suppression of Dynamical Network Biomarker Signals at the Predisease State () before Metabolic Syndrome in Mice by a Traditional Japanese Medicine (Kampo Formula) Bofutsushosan. <i>Evidence-based Complementary and Alternative Medicine</i> , 2020 , 2020, 9129134	2.3	5
38	Diastolic Cardiac Function Improvement by Liraglutide Is Mainly Body Weight Reduction Dependent but Independently Contributes to B-Type Natriuretic Peptide Reduction in Patients with Type 2 Diabetes with Preserved Ejection Fraction. <i>Journal of Diabetes Research</i> , 2021 , 2021, 8838026	3.9	5
37	CD206+ macrophage is an accelerator of endometriotic-like lesion via promoting angiogenesis in the endometriosis mouse model. <i>Scientific Reports</i> , 2021 , 11, 853	4.9	5
36	Genome-wide association studies identify two novel loci conferring susceptibility to diabetic retinopathy in Japanese patients with type 2 diabetes. <i>Human Molecular Genetics</i> , 2021 , 30, 716-726	5.6	5
35	Impact of body mass index on the efficacy and safety of ipragliflozin in Japanese patients with type2 diabetes mellitus: A subgroup analysis of 3-month interim results from the Specified Drug Use Results Survey of Ipragliflozin Treatment in Type2 Diabetic Patients: Long-term Use study. <i>Journal of Diabetes Research</i> , 2021 , 2021, 8838026	3.9	4
34	Physiologic Target, Molecular Evolution, and Pathogenic Functions of a Monoclonal Anti-Citrullinated Protein Antibody Obtained From a Patient With Rheumatoid Arthritis. <i>Arthritis and Rheumatology</i> , 2020 , 72, 2040-2049	9.5	4
33	Clinical manifestations of a sporadic maturity-onset diabetes of the young (MODY) 5 with a whole deletion of HNF1B based on 17q12 microdeletion. <i>Endocrine Journal</i> , 2019 , 66, 1113-1116	2.9	4

32	Generation and characterization of a Meflin-CreERT2 transgenic line for lineage tracing in white adipose tissue. <i>PLoS ONE</i> , 2021 , 16, e0248267	3.7	4
31	Anti-obesity effect of a hop-derived prenylflavonoid isoxanthohumol in a high-fat diet-induced obese mouse model. <i>Bioscience of Microbiota, Food and Health</i> , 2020 , 39, 175-182	3.2	3
30	Safety and effectiveness of ipragliflozin in Japanese patients with type 2 diabetes mellitus and impaired renal function: subgroup analysis of a 3-year post-marketing surveillance study (STELLA-LONG TERM). <i>Diabetology International</i> , 2021 , 12, 181-196	2.3	3
29	Calorie restriction-mediated restoration of hypothalamic signal transducer and activator of transcription 3 (STAT3) phosphorylation is not effective for lowering the body weight set point in IRS-2 knockout obese mice. <i>Diabetology International</i> , 2015 , 6, 321-335	2.3	2
28	Autoantibodies reactive to PEP08 are clinically related with morbidity and severity of interstitial lung disease in connective tissue diseases. <i>European Journal of Immunology</i> , 2018 , 48, 1717-1727	6.1	2
27	Family history of diabetes in both parents is strongly associated with impaired β cell function in Japanese type 2 diabetes patients. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 564-572	3.9	2
26	Gut microbiota, determined by dietary nutrients, drive modification of the plasma lipid profile and insulin resistance. <i>iScience</i> , 2021 , 24, 102445	6.1	2
25	Anti-programmed death ligand 1 therapy-induced type 1 diabetes presenting with multiple islet-related autoantibodies. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 253-254	3.9	2
24	Oral Administration of Nicotinamide Mononucleotide Is Safe and Efficiently Increases Blood Nicotinamide Adenine Dinucleotide Levels in Healthy Subjects.. <i>Frontiers in Nutrition</i> , 2022 , 9, 868640	6.2	2
23	Macrophage-specific hypoxia-inducible factor-1 α deletion suppresses the development of liver tumors in high-fat diet-fed obese and diabetic mice. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 1411-1418	3.9	1
22	Replication study of the association of rs7578597 in THADA, rs10886471 in GRK5, and rs7403531 in RASGRP1 with susceptibility to type 2 diabetes among a Japanese population. <i>Diabetology International</i> , 2015 , 6, 306-312	2.3	1
21	Isoxanthohumol, a hop-derived flavonoid, alters the metabolomics profile of mouse feces. <i>Bioscience of Microbiota, Food and Health</i> , 2020 , 39, 100-108	3.2	1
20	The PREDICTIVETM Study: a multinational, prospective observational study to evaluate the safety and efficacy of insulin detemir treatment in patients with type 1 and 2 diabetes—data from the Japan cohort. <i>Diabetology International</i> , 2012 , 3, 11-20	2.3	1
19	Wild-type ATTR amyloidosis of the ureter in a 56-year-old woman with rheumatoid arthritis and Sjögren's syndrome. <i>International Journal of Clinical and Experimental Pathology</i> , 2015 , 8, 8624-7	1.4	1
18	Appetite Loss as an Adverse Effect During Treatment with EGFR-TKIs in Elderly Patients with Non-small Cell Lung Cancer. <i>Anticancer Research</i> , 2016 , 36, 4951-4	2.3	1
17	Involuntary Measurement System for Respiratory Waveform for Prevention of Accidental Drowning during Bathing. <i>Advanced Biomedical Engineering</i> , 2013 , 2, 17-24	0.7	1
16	Type 1 diabetes woman with repeated miscarriages successfully gave birth after introducing an insulin pump with a predictive low glucose suspend feature. <i>Diabetology International</i> , 2021 , 12, 324-329	2.3	1
15	Real-world evidence for long-term safety and effectiveness of ipragliflozin in treatment-naïve versus non-naïve Japanese patients with type 2 diabetes mellitus: subgroup analysis of a 3-year post-marketing surveillance study (STELLA-LONG TERM). <i>Diabetology International</i> , 2021 , 12, 430-444	2.3	1

14	High prevalence of fragmented QRS on electrocardiography in Japanese patients with diabetes irrespective of metabolic syndrome. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 1680-1688	3.9	1
13	Effect of ipragliflozin on liver function in Japanese type 2 diabetes mellitus patients: subgroup analysis of a 3-year post-marketing surveillance study (STELLA-LONG TERM). <i>Endocrine Journal</i> , 2021 , 68, 905-918	2.9	1
12	TCR function analysis using a novel system reveals the multiple unconventional tumor-reactive T cells in human breast cancer-infiltrating lymphocytes. <i>European Journal of Immunology</i> , 2021 , 51, 2306-2316	6.1	1
11	Vascular endothelial growth factor (VEGF) activates Raf-1, mitogen-activated protein (MAP) kinases, and S6 kinase (p90rsk) in cultured rat cardiac myocytes 1998 , 175, 239		1
10	Isoliquiritigenin attenuates adipose tissue inflammation and metabolic syndrome by modifying gut bacteria composition in mice.. <i>Molecular Nutrition and Food Research</i> , 2022 , e2101119	5.9	1
9	A Comparison of Line Blots, Enzyme-linked Immunosorbent, and RNA-immunoprecipitation Assays of Antisyntetase Antibodies in Serum Samples from 44 Patients.. <i>Internal Medicine</i> , 2022 , 61, 313-322	1.1	0
8	Fate of adipocyte progenitors during adipogenesis in mice fed a high-fat diet. <i>Molecular Metabolism</i> , 2021 , 54, 101328	8.8	0
7	A Case of Systemic Lupus Erythematosus and Antineutrophil Cytoplasmic Antibodies-Associated Vasculitis Overlap Syndrome. <i>Case Reports in Rheumatology</i> , 2021 , 2021, 6690658	0.8	0
6	Pulmonary reversed halo sign as a presenting manifestation of rheumatoid arthritis. <i>Joint Bone Spine</i> , 2015 , 82, 207	2.9	
5	Safety and Effectiveness of Ipragliflozin in Elderly Versus Non-elderly Japanese Patients with Type2 Diabetes: Subgroup Analysis of STELLA-LONGTERM. <i>Diabetes Therapy</i> , 2021 , 12, 1359-1378	3.6	
4	Widespread Mechanic's Hands in Antisyntetase Syndrome With Anti-OJ Antibody. <i>Journal of Rheumatology</i> , 2021 , 48, 1341	4.1	
3	A case of wild-type transthyretin cardiac amyloidosis with rheumatoid arthritis. <i>Modern Rheumatology Case Reports</i> , 2021 , 5, 206-213	0.4	
2	A Case of Pulmonary Infarction Resembling Pneumonia during Immunosuppressive Treatment for Rheumatoid Arthritis. <i>Case Reports in Rheumatology</i> , 2021 , 2021, 5983580	0.8	
1	A Case of Granulomatosis with Polyangiitis with Various Breast Lesions as the Initial Symptoms: A Case-Based Review. <i>Case Reports in Rheumatology</i> , 2021 , 2021, 4416072	0.8	