

Felix Mauch

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/71919/publications.pdf>

Version: 2024-02-01

57
papers

10,773
citations

81743

39
h-index

149479

56
g-index

60
all docs

60
docs citations

60
times ranked

9596
citing authors

#	ARTICLE	IF	CITATIONS
1	Dual control of MAPK activities by AP2C1 and MKP1 MAPK phosphatases regulates defence responses in Arabidopsis. <i>Journal of Experimental Botany</i> , 2022, 73, 2369-2384.	2.4	12
2	Silica nanoparticles enhance disease resistance in Arabidopsis plants. <i>Nature Nanotechnology</i> , 2021, 16, 344-353.	15.6	172
3	Expression of a Fungal Lectin in Arabidopsis Enhances Plant Growth and Resistance Toward Microbial Pathogens and a Plant-Parasitic Nematode. <i>Frontiers in Plant Science</i> , 2021, 12, 657451.	1.7	13
4	Marasmius oreades agglutinin enhances resistance of Arabidopsis against plant-parasitic nematodes and a herbivorous insect. <i>BMC Plant Biology</i> , 2021, 21, 402.	1.6	1
5	Combined Abiotic Stresses Repress Defense and Cell Wall Metabolic Genes and Render Plants More Susceptible to Pathogen Infection. <i>Plants</i> , 2021, 10, 1946.	1.6	10
6	A <i>Phytophthora</i> effector protein promotes symplastic cell-to-cell trafficking by physical interaction with plasmodesmata-localised callose synthases. <i>New Phytologist</i> , 2020, 227, 1467-1478.	3.5	30
7	The potential of antagonistic moroccan <i>Streptomyces</i> isolates for the biological control of damping-off disease of pea (<i>Pisum sativum</i> L.) caused by <i>Aphanomyces euteiches</i> . <i>Journal of Phytopathology</i> , 2019, 167, 82-90.	0.5	10
8	A conserved RxLR effector interacts with host RABA-type GTPases to inhibit vesicle-mediated secretion of antimicrobial proteins. <i>Plant Journal</i> , 2018, 95, 187-203.	2.8	42
9	Protein phosphatase AP2C1 negatively regulates basal resistance and defense responses to <i>Pseudomonas syringae</i> . <i>Journal of Experimental Botany</i> , 2017, 68, erw485.	2.4	41
10	The sterol-binding activity of PATHOGENESIS-RELATED PROTEIN 1 reveals the mode of action of an antimicrobial protein. <i>Plant Journal</i> , 2017, 89, 502-509.	2.8	156
11	Regulatory and Functional Aspects of Indolic Metabolism in Plant Systemic Acquired Resistance. <i>Molecular Plant</i> , 2016, 9, 662-681.	3.9	62
12	Pathogen and Circadian Controlled 1 (PCC1) regulates polar lipid content, ABA-related responses, and pathogen defence in Arabidopsis thaliana. <i>Journal of Experimental Botany</i> , 2013, 64, 3385-3395.	2.4	42
13	Export of Salicylic Acid from the Chloroplast Requires the Multidrug and Toxin Extrusion-Like Transporter EDS5. <i>Plant Physiology</i> , 2013, 162, 1815-1821.	2.3	195
14	Immunocytochemical determination of the subcellular distribution of ascorbate in plants. <i>Planta</i> , 2011, 233, 1-12.	1.6	125
15	Glutathione Deficiency of the Arabidopsis Mutant <i>pad2-1</i> Affects Oxidative Stress-Related Events, Defense Gene Expression, and the Hypersensitive Response. <i>Plant Physiology</i> , 2011, 157, 2000-2012.	2.3	90
16	Disease resistance of Arabidopsis to <i>Phytophthora brassicae</i> is established by the sequential action of indole glucosinolates and camalexin. <i>Plant Journal</i> , 2010, 62, 840-851.	2.8	180
17	Indolic secondary metabolites protect Arabidopsis from the oomycete pathogen <i>Phytophthora brassicae</i> . <i>Plant Signaling and Behavior</i> , 2010, 5, 1099-1101.	1.2	25
18	The chloroplast protein RPH1 plays a role in the immune response of Arabidopsis to <i>Phytophthora brassicae</i> . <i>Plant Journal</i> , 2009, 58, 287-298.	2.8	39

#	ARTICLE	IF	CITATIONS
19	The glutathione-deficient mutant <i>pad2-1</i> accumulates lower amounts of glucosinolates and is more susceptible to the insect herbivore <i>Spodoptera littoralis</i> . <i>Plant Journal</i> , 2008, 55, 774-786.	2.8	182
20	Evolution of the cutinase gene family: Evidence for lateral gene transfer of a candidate <i>Phytophthora</i> virulence factor. <i>Gene</i> , 2008, 408, 1-8.	1.0	67
21	Subcellular immunocytochemical analysis detects the highest concentrations of glutathione in mitochondria and not in plastids. <i>Journal of Experimental Botany</i> , 2008, 59, 4017-4027.	2.4	123
22	The PP2C-Type Phosphatase AP2C1, Which Negatively Regulates MPK4 and MPK6, Modulates Innate Immunity, Jasmonic Acid, and Ethylene Levels in <i>Arabidopsis</i> . <i>Plant Cell</i> , 2007, 19, 2213-2224.	3.1	302
23	Priming: Getting Ready for Battle. <i>Molecular Plant-Microbe Interactions</i> , 2006, 19, 1062-1071.	1.4	1,241
24	Identification of PAD2 as a γ -glutamylcysteine synthetase highlights the importance of glutathione in disease resistance of <i>Arabidopsis</i> . <i>Plant Journal</i> , 2006, 49, 159-172.	2.8	329
25	Large-Scale Gene Discovery in the Oomycete <i>Phytophthora infestans</i> Reveals Likely Components of Phytopathogenicity Shared with True Fungi. <i>Molecular Plant-Microbe Interactions</i> , 2005, 18, 229-243.	1.4	160
26	Sulphur Deficiency Causes a Reduction in Antimicrobial Potential and Leads to Increased Disease Susceptibility of Oilseed Rape. <i>Journal of Phytopathology</i> , 2005, 153, 27-36.	0.5	61
27	The role of abscisic acid in plant-pathogen interactions. <i>Current Opinion in Plant Biology</i> , 2005, 8, 409-414.	3.5	706
28	Crosstalk and differential response to abiotic and biotic stressors reflected at the transcriptional level of effector genes from secondary metabolism. <i>Plant Molecular Biology</i> , 2004, 54, 817-835.	2.0	111
29	Quantification of induced resistance against <i>Phytophthora</i> species expressing GFP as a vital marker: β -aminobutyric acid but not BTH protects potato and <i>Arabidopsis</i> from infection. <i>Molecular Plant Pathology</i> , 2003, 4, 237-248.	2.0	97
30	The Rapid Induction of Glutathione S-Transferases AtGSTF2 and AtGSTF6 by Avirulent <i>Pseudomonas syringae</i> is the Result of Combined Salicylic Acid and Ethylene Signaling. <i>Plant and Cell Physiology</i> , 2003, 44, 750-757.	1.5	66
31	Expression Profile Matrix of <i>Arabidopsis</i> Transcription Factor Genes Suggests Their Putative Functions in Response to Environmental Stresses[W]. <i>Plant Cell</i> , 2002, 14, 559-574.	3.1	849
32	Probing the diversity of the <i>Arabidopsis</i> glutathione S-transferase gene family. <i>Plant Molecular Biology</i> , 2002, 49, 515-532.	2.0	465
33	Characterization of an <i>Arabidopsis-Phytophthora</i> Pathosystem: resistance requires a functional PAD2 gene and is independent of salicylic acid, ethylene and jasmonic acid signalling. <i>Plant Journal</i> , 2001, 28, 293-305.	2.8	161
34	Manipulation of salicylate content in <i>Arabidopsis thaliana</i> by the expression of an engineered bacterial salicylate synthase. <i>Plant Journal</i> , 2001, 25, 67-77.	2.8	110
35	Construction and application of a microprojectile system for the transfection of organotypic brain slices. <i>Journal of Neuroscience Methods</i> , 2000, 101, 171-179.	1.3	6
36	Constitutive expression of the defense-related <i>Rir1b</i> gene in transgenic rice plants confers enhanced resistance to the rice blast fungus <i>Magnaporthe grisea</i> . <i>Plant Molecular Biology</i> , 2000, 43, 59-66.	2.0	37

#	ARTICLE	IF	CITATIONS
37	Characterization of the rice pathogen-related protein Rir1a and regulation of the corresponding gene. <i>Plant Molecular Biology</i> , 1998, 38, 577-586.	2.0	23
38	Mechanosensitive Expression of a Lipoxygenase Gene in Wheat. <i>Plant Physiology</i> , 1997, 114, 1561-1566.	2.3	53
39	Characterization of a rice gene induced by <i>Pseudomonas syringae</i> pv. <i>syringae</i> : requirement for the bacterial <i>lemA</i> gene function. <i>Physiological and Molecular Plant Pathology</i> , 1995, 46, 71-81.	1.3	40
40	Quantitative field resistance of wheat to powdery mildew and defense reactions at the seedling stage: identification of a potential marker. <i>Physiological and Molecular Plant Pathology</i> , 1995, 47, 185-199.	1.3	18
41	Differential Induction of Distinct Glutathione-S-Transferases of Wheat by Xenobiotics and by Pathogen Attack. <i>Plant Physiology</i> , 1993, 102, 1193-1201.	2.3	234
42	Ethylene-induced chitinase and β -1,3-glucanase accumulate specifically in the lower epidermis and along vascular strands of bean leaves. <i>Planta</i> , 1992, 186, 367-75.	1.6	39
43	<i>Research Notes</i> Sequence and Expression of a Wheat Gene that Encodes a Novel Protein Associated with Pathogen Defense. <i>Molecular Plant-Microbe Interactions</i> , 1992, 5, 516.	1.4	45
44	A wheat glutathione-S-transferase gene with transposon-like sequences in the promoter region. <i>Plant Molecular Biology</i> , 1991, 16, 1089-1091.	2.0	15
45	Sequence and tissue-specific expression of a putative peroxidase gene from wheat (<i>Triticum aestivum</i>) Tj ETQq1 1 0.784314 ggBT /Ov	2.0	50
46	Cloning and sequencing of cDNAs encoding a pathogen-induced putative peroxidase of wheat (<i>Triticum aestivum</i> L.). <i>Plant Molecular Biology</i> , 1991, 16, 329-331.	2.0	81
47	Sequence of a wheat cDNA encoding a pathogen-induced thaumatin-like protein. <i>Plant Molecular Biology</i> , 1991, 17, 283-285.	2.0	73
48	A Pathogen-Induced Wheat Gene Encodes a Protein Homologous to Glutathione-S-Transferases. <i>Molecular Plant-Microbe Interactions</i> , 1991, 4, 14.	1.4	124
49	Functional Implications of the Subcellular Localization of Ethylene-Induced Chitinase and β -1,3-Glucanase in Bean Leaves. <i>Plant Cell</i> , 1989, 1, 447.	3.1	192
50	Antifungal Hydrolases in Pea Tissue. <i>Plant Physiology</i> , 1988, 88, 936-942.	2.3	1,120
51	Antifungal Hydrolases in Pea Tissue. <i>Plant Physiology</i> , 1988, 87, 325-333.	2.3	304
52	Colorimetric assay for chitinase. <i>Methods in Enzymology</i> , 1988, , 430-435.	0.4	203
53	Chitinase from <i>Phaseolus vulgaris</i> leaves. <i>Methods in Enzymology</i> , 1988, 161, 479-484.	0.4	10
54	Plant chitinases are potent inhibitors of fungal growth. <i>Nature</i> , 1986, 324, 365-367.	13.7	871

#	ARTICLE	IF	CITATIONS
55	Ethylene: Symptom, Not Signal for the Induction of Chitinase and β -1,3-Glucanase in Pea Pods by Pathogens and Elicitors. <i>Plant Physiology</i> , 1984, 76, 607-611.	2.3	305
56	Chitinase in bean leaves: induction by ethylene, purification, properties, and possible function. <i>Planta</i> , 1983, 157, 22-31.	1.6	649
57	Potential of Moroccan isolates of plant growth promoting streptomycetes for biocontrol of the root rot disease of pea plants caused by the oomycete pathogen <i>Aphanomyces euteiches</i> .. <i>Biocontrol Science and Technology</i> , 0, , 1-18.	0.5	2