

K Thomas Jensen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7193082/publications.pdf>

Version: 2024-02-01

56
papers

2,208
citations

186265
28
h-index

214800
47
g-index

56
all docs

56
docs citations

56
times ranked

1572
citing authors

#	ARTICLE	IF	CITATIONS
1	Impacts and effects of a historical high and ENSO linked freshwater inflow in the tropical estuary Nha Phu, southeast Vietnam. <i>Regional Studies in Marine Science</i> , 2018, 17, 28-37.	0.7	3
2	The most vagile host as the main determinant of population connectivity in marine macroparasites. <i>Marine Ecology - Progress Series</i> , 2015, 520, 85-99.	1.9	23
3	Impact of trematodes on the population structure and shell shape of the estuarine mud snail <i>Hydrobia ulvae</i> from a Southern European estuary. <i>Marine Ecology</i> , 2014, 35, 1-10.	1.1	0
4	Seasonal and spatial distribution of mesozooplankton in a tropical estuary, Nha Phu, South Central Viet Nam. <i>Biologia (Poland)</i> , 2014, 69, 80-91.	1.5	10
5	Growth Performance of Indian Major Carps and Their Hybrids in Polyculture in Bangladesh. <i>Journal of Applied Aquaculture</i> , 2014, 26, 157-168.	1.4	2
6	Cytogenetic characterisation of the razor shells <i>Ensis directus</i> (Conrad, 1843) and <i>E. minor</i> (Chenu.) <i>Tj ETQq 0 0 rgBT /Overlock 10 TF 5</i>	1.3	8
7	Population genetic analysis of <i>Ensis directus</i> unveils high genetic variation in the introduced range and reveals a new species from the NW Atlantic. <i>Marine Biology</i> , 2012, 159, 2209-2227.	1.5	10
8	The cockle <i>Cerastoderma edule</i> at Northeast Atlantic shores: genetic signatures of glacial refugia. <i>Marine Biology</i> , 2012, 159, 221-230.	1.5	33
9	Trematode fauna of <i>Hydrobia ulvae</i> (Gastropoda: Prosobranchia) in a eutrophic temperate estuary. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2011, 91, 913-921.	0.8	6
10	The linked units of 5S rDNA and U1 snDNA of razor shells (Mollusca: Bivalvia: Pharidae). <i>Heredity</i> , 2011, 107, 127-142.	2.6	17
11	Climate influences parasite-mediated competitive release. <i>Parasitology</i> , 2011, 138, 1436-1441.	1.5	13
12	The introduced clam <i>Ensis americanus</i> in the Wadden Sea: field experiment on impact of bird predation and tidal level on survival and growth. <i>Helgoland Marine Research</i> , 2010, 64, 93-100.	1.3	17
13	Digenean trematode species in the cockle <i>Cerastoderma edule</i> : identification key and distribution along the north-eastern Atlantic shoreline. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2009, 89, 543-556.	0.8	65
14	Invaders interfere with native parasite-host interactions. <i>Biological Invasions</i> , 2009, 11, 1421-1429.	2.4	93
15	Parasites in the northern Wadden Sea: a conservative ecosystem component over 4 decades. <i>Helgoland Marine Research</i> , 2008, 62, 37-47.	1.3	16
16	The role of biotic factors in the transmission of free-living endohelminth stages. <i>Parasitology</i> , 2008, 135, 407-426.	1.5	195
17	Ambient fauna impairs parasite transmission in a marine parasite-host system. <i>Parasitology</i> , 2008, 135, 1111-1116.	1.5	67
18	Metazoan parasites in an intermediate host population near its southern border: the common cockle (<i>Cerastoderma edule</i>) and its trematodes in a Moroccan coastal lagoon (Merja Zerga). <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2008, 88, 357-364.	0.8	26

#	ARTICLE	IF	CITATIONS
19	Production of marine trematode cercariae: a potentially overlooked path of energy flow in benthic systems. <i>Marine Ecology - Progress Series</i> , 2008, 372, 147-155.	1.9	70
20	Effects of restoration management on the estuarine isopod <i>Cyathura carinata</i> : mediation by trematodes and habitat change. <i>Marine Biology</i> , 2007, 151, 109-118.	1.5	7
21	Effect of intermediate host size (<i>Cerastoderma edule</i>) on infectivity of cercariae of <i>Himasthla quissetensis</i> (Echinostomatidae: Trematoda). <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2005, 85, 809-812.	0.8	21
22	Impact of microphallid trematodes on the survivorship, growth, and reproduction of an isopod (<i>Cyathura carinata</i>). <i>Journal of Experimental Marine Biology and Ecology</i> , 2005, 318, 191-199.	1.5	21
23	Biologically induced differences in erodibility and aggregation of subtidal and intertidal sediments: a possible cause for seasonal changes in sediment deposition. <i>Journal of Marine Systems</i> , 2005, 55, 123-138.	2.1	70
24	Repetitive sequences in the ITS1 region of ribosomal DNA in congeneric microphallid species (Trematoda: Digenea). <i>Parasitology Research</i> , 2005, 97, 420-423.	1.6	20
25	Infection characteristics of a trematode in an estuarine isopod: influence of substratum. <i>Hydrobiologia</i> , 2005, 539, 149-155.	2.0	3
26	Surface activity of <i>Corophium volutator</i> : A role for parasites?. <i>Journal of Sea Research</i> , 2005, 54, 176-184.	1.6	3
27	A New Video and Digital Camera System for Studies of the Dynamics of Microtopographic Features on Tidal Flats. <i>Marine Georesources and Geotechnology</i> , 2004, 22, 115-122.	2.1	3
28	Effects of food concentration on clearance rate and energy budget of the Arctic bivalve <i>Hiatella arctica</i> (L) at subzero temperature. <i>Journal of Experimental Marine Biology and Ecology</i> , 2004, 311, 171-183.	1.5	28
29	Use of ITS rDNA for discriminating of larval stages of two microphallid (Digenea) species using <i>Hydrobia ulvae</i> (Pennant, 1777) and <i>Corophium volutator</i> (Pallas, 1766) as intermediate hosts. <i>Parasitology Research</i> , 2004, 93, 304-10.	1.6	15
30	Trematodes in a <i>Cyathura carinata</i> population from a temperate intertidal estuary: infection patterns and impact on host. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2004, 84, 1151-1158.	0.8	13
31	In situ growth of juvenile cockles, <i>Cerastoderma edule</i> , experimentally infected with larval trematodes (<i>Himasthla interrupta</i>). <i>Journal of Sea Research</i> , 2003, 50, 37-43.	1.6	33
32	Enhanced erodibility of fine-grained marine sediments by <i>Hydrobia ulvae</i> . <i>Journal of Sea Research</i> , 2002, 48, 51-58.	1.6	64
33	Annual growth bands in the bivalve <i>Hiatella arctica</i> validated by a mark-recapture study in NE Greenland. <i>Polar Biology</i> , 2002, 25, 794-796.	1.2	18
34	Growth and production of <i>Hiatella arctica</i> (Bivalvia) in a high-Arctic fjord (Young Sound, Northeast) Tj ETQq0 0 0 rgPT /Overlock 10 Tf 50	1.9	45
35	Macrozoobenthic community structure in a high-arctic East Greenland fjord. <i>Polar Biology</i> , 2000, 23, 792-801.	1.2	60
36	Effect of intermediate host size (<i>Cerastoderma edule</i>) on infectivity of cercariae of three <i>Himasthla</i> species (Echinostomatidae, Trematoda). <i>Journal of Experimental Marine Biology and Ecology</i> , 1999, 238, 259-269.	1.5	46

#	ARTICLE	IF	CITATIONS
37	Reduced survivorship of <i>Himasthla</i> (Trematoda, Digenea)-infected cockles (<i>Cerastoderma edule</i>) exposed to oxygen depletion. <i>Journal of Sea Research</i> , 1999, 42, 325-331.	1.6	48
38	Infectivity of <i>Himasthla</i> spp. (Trematoda) in cockle (<i>Cerastoderma edule</i>) spat. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 1999, 79, 265-271.	0.8	34
39	Influence of trematode infections on in situ growth rates of <i>Littorina littorea</i> . <i>Journal of the Marine Biological Association of the United Kingdom</i> , 1999, 79, 425-430.	0.8	30
40	The influence of the trematode <i>Microphallus claviformis</i> on two congeneric intermediate host species (<i>Corophium</i>): infection characteristics and host survival. <i>Journal of Experimental Marine Biology and Ecology</i> , 1998, 227, 35-48.	1.5	38
41	Change of Topography and Sediment Characteristics on an Intertidal Mud-Flat Following Mass-Mortality of the Amphipod <i>Corophium Volutator</i> . <i>Journal of the Marine Biological Association of the United Kingdom</i> , 1998, 78, 1167-1180.	0.8	66
42	Infection characteristics of <i>Himasthla elongata</i> cercariae in cockles as a function of water current. <i>Diseases of Aquatic Organisms</i> , 1998, 34, 63-70.	1.0	53
43	Title is missing!. <i>Hydrobiologia</i> , 1997, 355, 61-70.	2.0	25
44	Parasite transmission between soft-bottom invertebrates: temperature mediated infection rates and mortality in <i>Corophium volutator</i> . <i>Marine Ecology - Progress Series</i> , 1997, 151, 123-134.	1.9	98
45	Red List of macrofaunal benthic invertebrates of the Wadden Sea. <i>Helgoländer Meeresuntersuchungen</i> , 1996, 50, 69-76.	0.2	7
46	The effect of larval trematodes on the survival rates of two species of mud snails (hydrobiidae) experimentally exposed to desiccation, freezing and anoxia. <i>Helgoländer Meeresuntersuchungen</i> , 1996, 50, 327-335.	0.2	27
47	The enigma of gigantism: effect of larval trematodes on growth, fecundity, egestion and locomotion in <i>Hydrobia ulvae</i> (pennant) (gastropoda: prosobranchia). <i>Journal of Experimental Marine Biology and Ecology</i> , 1994, 181, 53-66.	1.5	115
48	Field and laboratory experiments on interactions among an infaunal polychaete, <i>Nereis diversicolor</i> , and two amphipods, <i>Corophium volutator</i> & <i>C. arenarium</i> : effects on survival, recruitment and migration. <i>Journal of Experimental Marine Biology and Ecology</i> , 1993, 168, 259-278.	1.5	23
49	Density-dependent growth in cockles (<i>Cerastoderma edule</i>): evidence from interannual comparisons. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 1993, 73, 333-342.	0.8	44
50	Dynamics and growth of the cockle, <i>Cerastoderma edule</i> , on an intertidal mud-flat in the Danish Wadden sea: Effects of submersion time and density. <i>Journal of Sea Research</i> , 1992, 28, 335-345.	1.0	74
51	Mass mortality in two common soft-bottom invertebrates, <i>Hydrobia ulvae</i> and <i>Corophium volutator</i> -the possible role of trematodes. <i>Helgoländer Meeresuntersuchungen</i> , 1992, 46, 329-339.	0.2	82
52	Macrozoobenthos on an intertidal mudflat in the Danish Wadden Sea: Comparisons of surveys made in the 1930s, 1940s and 1980s. <i>Helgoländer Meeresuntersuchungen</i> , 1992, 46, 363-376.	0.2	32
53	Choice of microhabitat in tactile foraging dunlins <i>Calidris alpina</i> : the importance of sediment penetrability. <i>Marine Ecology - Progress Series</i> , 1992, 85, 1-8.	1.9	57
54	A field experiment on competition between <i>Corophium volutator</i> (Pallas) and <i>Corophium arenarium</i> Crawford (Crustacea: Amphipoda): effects on survival, reproduction and recruitment. <i>Journal of Experimental Marine Biology and Ecology</i> , 1990, 137, 1-24.	1.5	50

#	ARTICLE	IF	CITATIONS
55	The importance of some epibenthic predators on the density of juvenile benthic macrofauna in the Danish Wadden Sea. <i>Journal of Experimental Marine Biology and Ecology</i> , 1985, 89, 157-174.	1.5	119
56	The presence of the bivalve <i>Cerastoderma edule</i> affects migration, survival and reproduction of the amphipod <i>Corophium volutator</i> . <i>Marine Ecology - Progress Series</i> , 1985, 25, 269-277.	1.9	42