Akira Miyauchi

List of Publications by Citations

Source: https://exaly.com/author-pdf/7200780/akira-miyauchi-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

60 13,850 364 104 h-index g-index citations papers 16,031 6.45 3.5 374 L-index avg, IF ext. citations ext. papers

#	Paper	IF	Citations
364	Electrophysiologic recurrent laryngeal nerve monitoring during thyroid and parathyroid surgery: international standards guideline statement. <i>Laryngoscope</i> , 2011 , 121 Suppl 1, S1-16	3.6	636
363	An observation trial without surgical treatment in patients with papillary microcarcinoma of the thyroid. <i>Thyroid</i> , 2003 , 13, 381-7	6.2	582
362	An observational trial for papillary thyroid microcarcinoma in Japanese patients. <i>World Journal of Surgery</i> , 2010 , 34, 28-35	3.3	523
361	Patient age is significantly related to the progression of papillary microcarcinoma of the thyroid under observation. <i>Thyroid</i> , 2014 , 24, 27-34	6.2	472
360	External branch of the superior laryngeal nerve monitoring during thyroid and parathyroid surgery: International Neural Monitoring Study Group standards guideline statement. <i>Laryngoscope</i> , 2013 , 123 Suppl 4, S1-14	3.6	218
359	Clinical aspects of primary thyroid lymphoma: diagnosis and treatment based on our experience of 119 cases. <i>Thyroid</i> , 1993 , 3, 93-9	6.2	209
358	Prognostic significance of extrathyroid extension of papillary thyroid carcinoma: massive but not minimal extension affects the relapse-free survival. <i>World Journal of Surgery</i> , 2006 , 30, 780-6	3.3	205
357	Incidences of Unfavorable Events in the Management of Low-Risk Papillary Microcarcinoma of the Thyroid by Active Surveillance Versus Immediate Surgery. <i>Thyroid</i> , 2016 , 26, 150-5	6.2	200
356	Risk factors for recurrence to the lymph node in papillary thyroid carcinoma patients without preoperatively detectable lateral node metastasis: validity of prophylactic modified radical neck dissection. <i>World Journal of Surgery</i> , 2007 , 31, 2085-91	3.3	197
355	BRAF mutation in papillary thyroid carcinoma in a Japanese population: its lack of correlation with high-risk clinicopathological features and disease-free survival of patients. <i>Endocrine Journal</i> , 2009 , 56, 89-97	2.9	194
354	Comparison of methimazole and propylthiouracil in patients with hyperthyroidism caused by GravesQdisease. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007 , 92, 2157-62	5.6	194
353	Papillary microcarcinoma of the thyroid: how should it be treated?. <i>World Journal of Surgery</i> , 2004 , 28, 1115-21	3.3	191
352	A Clinical Framework to Facilitate Risk Stratification When Considering an Active Surveillance Alternative to Immediate Biopsy and Surgery in Papillary Microcarcinoma. <i>Thyroid</i> , 2016 , 26, 144-9	6.2	185
351	Prognostic impact of serum thyroglobulin doubling-time under thyrotropin suppression in patients with papillary thyroid carcinoma who underwent total thyroidectomy. <i>Thyroid</i> , 2011 , 21, 707-16	6.2	183
350	Preoperative ultrasonographic examination for lymph node metastasis: usefulness when designing lymph node dissection for papillary microcarcinoma of the thyroid. <i>World Journal of Surgery</i> , 2004 , 28, 498-501	3.3	171
349	Prognostic factors and treatment outcomes for anaplastic thyroid carcinoma: ATC Research Consortium of Japan cohort study of 677 patients. <i>World Journal of Surgery</i> , 2012 , 36, 1247-54	3.3	170
348	Prognostic factors for recurrence of papillary thyroid carcinoma in the lymph nodes, lung, and bone: analysis of 5,768 patients with average 10-year follow-up. <i>World Journal of Surgery</i> , 2012 , 36, 127	74 ² 8	162

(2018-1985)

347	Intrathyroidal epithelial thymoma: an entity distinct from squamous cell carcinoma of the thyroid. <i>World Journal of Surgery</i> , 1985 , 9, 128-35	3.3	154
346	Clinical significance of lymph node metastasis of thyroid papillary carcinoma located in one lobe. World Journal of Surgery, 2006 , 30, 1821-8	3.3	151
345	Clinical Trials of Active Surveillance of Papillary Microcarcinoma of the Thyroid. <i>World Journal of Surgery</i> , 2016 , 40, 516-22	3.3	148
344	Ultrasonographically and anatomopathologically detectable node metastases in the lateral compartment as indicators of worse relapse-free survival in patients with papillary thyroid carcinoma. World Journal of Surgery, 2005, 29, 917-20	3.3	148
343	Usefulness of thyroglobulin measurement in fine-needle aspiration biopsy specimens for diagnosing cervical lymph node metastasis in patients with papillary thyroid cancer. <i>World Journal of Surgery</i> , 2005 , 29, 483-5	3.3	143
342	Accurate and simple method of diagnosing thyroid nodules the modified technique of ultrasound-guided fine needle aspiration biopsy. <i>Thyroid</i> , 1995 , 5, 141-5	6.2	129
341	Distinct clinical, serological, and sonographic characteristics of hashimoto@thyroiditis based with and without IgG4-positive plasma cells. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, 1309	- 1 5	124
340	Clinical characteristics of 852 patients with subacute thyroiditis before treatment. <i>Internal Medicine</i> , 2008 , 47, 725-9	1.1	117
339	Prognosis of patients with papillary thyroid carcinoma having clinically apparent metastasis to the lateral compartment. <i>Endocrine Journal</i> , 2009 , 56, 759-66	2.9	116
338	Prognostic factors and therapeutic strategies for differentiated carcinomas of the thyroid. <i>Endocrine Journal</i> , 2009 , 56, 177-92	2.9	114
337	Cribriform-morular variant of papillary thyroid carcinoma: a pathological and molecular genetic study with evidence of frequent somatic mutations in exon 3 of the beta-catenin gene. <i>Journal of Pathology</i> , 2003 , 199, 58-67	9.4	112
336	A therapeutic strategy for incidentally detected papillary microcarcinoma of the thyroid. <i>Nature Clinical Practice Endocrinology and Metabolism</i> , 2007 , 3, 240-8		108
335	Induction chemotherapy with weekly paclitaxel administration for anaplastic thyroid carcinoma. <i>Thyroid</i> , 2010 , 20, 7-14	6.2	105
334	Risk factors contributing to a poor prognosis of papillary thyroid carcinoma: validity of UICC/AJCC TNM classification and stage grouping. <i>World Journal of Surgery</i> , 2007 , 31, 838-48	3.3	105
333	Insights into the Management of Papillary Microcarcinoma of the Thyroid. <i>Thyroid</i> , 2018 , 28, 23-31	6.2	98
332	Thyroid cancer detected by ultrasound-guided fine-needle aspiration biopsy. <i>World Journal of Surgery</i> , 1996 , 20, 848-53; discussion 853	3.3	96
331	Analysis of 754 cases of antithyroid drug-induced agranulocytosis over 30 years in Japan. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 4776-83	5.6	95
330	Estimation of the lifetime probability of disease progression of papillary microcarcinoma of the thyroid during active surveillance. <i>Surgery</i> , 2018 , 163, 48-52	3.6	95

329	Clinicopathologic significance of intrathyroidal epithelial thymoma/carcinoma showing thymus-like differentiation: a collaborative study with Member Institutes of The Japanese Society of Thyroid Surgery. <i>American Journal of Clinical Pathology</i> , 2007 , 127, 230-6	1.9	92
328	Relation of doubling time of plasma calcitonin levels to prognosis and recurrence of medullary thyroid carcinoma. <i>Annals of Surgery</i> , 1984 , 199, 461-6	7.8	88
327	Excellent prognosis of patients with solitary T1N0M0 papillary thyroid carcinoma who underwent thyroidectomy and elective lymph node dissection without radioiodine therapy. <i>World Journal of Surgery</i> , 2010 , 34, 1285-90	3.3	86
326	Improvement in phonation after reconstruction of the recurrent laryngeal nerve in patients with thyroid cancer invading the nerve. <i>Surgery</i> , 2009 , 146, 1056-62	3.6	84
325	Cribriform-morular variant of papillary thyroid carcinoma: clue to early detection of familial adenomatous polyposis-associated colon cancer. <i>World Journal of Surgery</i> , 2004 , 28, 886-9	3.3	83
324	Methimazole-induced agranulocytosis in patients with GravesQdisease is more frequent with an initial dose of 30 mg daily than with 15 mg daily. <i>Thyroid</i> , 2009 , 19, 559-63	6.2	82
323	Quantitative measurement of thyroid blood flow for differentiation of painless thyroiditis from GravesQdisease. <i>Clinical Endocrinology</i> , 2007 , 67, 41-5	3.4	81
322	Prevalence and prognostic significance of poor differentiation and tall cell variant in papillary carcinoma in Japan. <i>World Journal of Surgery</i> , 2008 , 32, 1535-43; discussion 1544-5	3.3	81
321	Biological behavior and prognosis of familial papillary thyroid carcinoma. Surgery, 2009, 145, 100-5	3.6	80
320	TSH-suppressive doses of levothyroxine are required to achieve preoperative native serum triiodothyronine levels in patients who have undergone total thyroidectomy. <i>European Journal of Endocrinology</i> , 2012 , 167, 373-8	6.5	80
319	Diagnosis of medullary thyroid carcinoma by calcitonin measurement in fine-needle aspiration biopsy specimens. <i>Thyroid</i> , 2007 , 17, 635-8	6.2	80
318	Overall Survival of Papillary Thyroid Carcinoma Patients: A Single-Institution Long-Term Follow-Up of 5897 Patients. <i>World Journal of Surgery</i> , 2018 , 42, 615-622	3.3	79
317	Ultrasonographic evaluation of thyroid nodules in 900 patients: comparison among ultrasonographic, cytological, and histological findings. <i>Thyroid</i> , 2007 , 17, 1269-76	6.2	79
316	Lateral and mediastinal lymph node dissection in differentiated thyroid carcinoma: indications, benefits, and risks. <i>World Journal of Surgery</i> , 2007 , 31, 905-15	3.3	79
315	Comparison of the costs of active surveillance and immediate surgery in the management of low-risk papillary microcarcinoma of the thyroid. <i>Endocrine Journal</i> , 2017 , 64, 59-64	2.9	77
314	International neuromonitoring study group guidelines 2018: Part II: Optimal recurrent laryngeal nerve management for invasive thyroid cancer-incorporation of surgical, laryngeal, and neural electrophysiologic data. <i>Laryngoscope</i> , 2018 , 128 Suppl 3, S18-S27	3.6	74
313	Prognosis and prognostic factors of follicular carcinoma in Japan: importance of postoperative pathological examination. <i>World Journal of Surgery</i> , 2007 , 31, 1417-24	3.3	73
312	Distinct histopathological features of Hashimoto@thyroiditis with respect to IgG4-related disease. <i>Modern Pathology</i> , 2012 , 25, 1086-97	9.8	71

311	IgG4-related disease of the thyroid glands. Endocrine Journal, 2012, 59, 273-81	2.9	70	
310	Prognostic factors of minimally invasive follicular thyroid carcinoma: extensive vascular invasion significantly affects patient prognosis. <i>Endocrine Journal</i> , 2013 , 60, 637-42	2.9	69	
309	Prospective study of vocal fold function after loss of the neuromonitoring signal in thyroid surgery: The International Neural Monitoring Study Group@POLT study. <i>Laryngoscope</i> , 2016 , 126, 1260-6	3.6	69	
308	The FOXE1 and NKX2-1 loci are associated with susceptibility to papillary thyroid carcinoma in the Japanese population. <i>Journal of Medical Genetics</i> , 2011 , 48, 645-8	5.8	68	
307	Prospective trial of unilateral surgery for nonhereditary medullary thyroid carcinoma in patients without germline RET mutations. <i>World Journal of Surgery</i> , 2002 , 26, 1023-8	3.3	68	
306	Thyroid function in early pregnancy in Japanese healthy women: relation to urinary iodine excretion, emesis, and fetal and child development. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 1683-8	5.6	66	
305	The role of ansa-to-recurrent-laryngeal nerve anastomosis in operations for thyroid cancer. <i>The European Journal of Surgery</i> , 1998 , 164, 927-33		63	
304	Papillary thyroid microcarcinoma might progress during pregnancy. <i>Thyroid</i> , 2014 , 24, 840-4	6.2	58	
303	Establishment of an intraoperative staging system (iStage) by improving UICC TNM classification system for papillary thyroid carcinoma. <i>World Journal of Surgery</i> , 2010 , 34, 2570-80	3.3	58	
302	Recovery of parathyroid function after total thyroidectomy: long-term follow-up study. <i>ANZ Journal of Surgery</i> , 2005 , 75, 532-6	1	57	
301	Nonoperative management of low-risk differentiated thyroid carcinoma. <i>Current Opinion in Oncology</i> , 2015 , 27, 15-20	4.2	56	
300	Serial changes in liver function tests in patients with thyrotoxicosis induced by Graves Qdisease and painless thyroiditis. <i>Thyroid</i> , 2008 , 18, 283-7	6.2	56	
299	Ill-defined edge on ultrasonographic examination can be a marker of aggressive characteristic of papillary thyroid microcarcinoma. <i>World Journal of Surgery</i> , 2005 , 29, 1007-11; discussion 1011-2	3.3	55	
298	Usefulness of ultrasonography for diagnosis of malignant lymphoma of the thyroid. <i>Thyroid</i> , 2006 , 16, 983-7	6.2	53	
297	Patient Experience of Thyroid Cancer Active Surveillance in Japan. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2019 , 145, 363-370	3.9	52	
296	Tumor size is the strongest predictor of microscopic lymph node metastasis and lymph node recurrence of N0 papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2013 , 60, 113-7	2.9	51	
295	Our experience of treatment of cribriform morular variant of papillary thyroid carcinoma; difference in clinicopathological features of FAP-associated and sporadic patients. <i>Endocrine Journal</i> , 2011 , 58, 685-9	2.9	50	
294	Prognostic significance of ki-67 labeling index in papillary thyroid carcinoma. <i>World Journal of Surgery</i> , 2010 , 34, 3015-21	3.3	50	

293	Effect of levo-thyroxine replacement on non-high-density lipoprotein cholesterol in hypothyroid patients. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007 , 92, 608-11	5.6	50
292	Trends in the Implementation of Active Surveillance for Low-Risk Papillary Thyroid Microcarcinomas at Kuma Hospital: Gradual Increase and Heterogeneity in the Acceptance of This New Management Option. <i>Thyroid</i> , 2018 , 28, 488-495	6.2	48
291	Extranodal tumor extension to adjacent organs predicts a worse cause-specific survival in patients with papillary thyroid carcinoma. <i>World Journal of Surgery</i> , 2007 , 31, 1194-201	3.3	48
290	Indications and Strategy for Active Surveillance of Adult Low-Risk Papillary Thyroid Microcarcinoma: Consensus Statements from the Japan Association of Endocrine Surgery Task Force on Management for Papillary Thyroid Microcarcinoma. <i>Thyroid</i> , 2021 , 31, 183-192	6.2	48
289	Differentiation of postpartum Graves Q hyrotoxicosis from postpartum destructive thyrotoxicosis using antithyrotropin receptor antibodies and thyroid blood flow. <i>Thyroid</i> , 2014 , 24, 1027-31	6.2	47
288	Biochemical Markers Reflecting Thyroid Function in Athyreotic Patients on Levothyroxine Monotherapy. <i>Thyroid</i> , 2017 , 27, 484-490	6.2	46
287	Effects of Pregnancy on Papillary Microcarcinomas of the Thyroid Re-Evaluated in the Entire Patient Series at Kuma Hospital. <i>Thyroid</i> , 2016 , 26, 156-60	6.2	46
286	Revisiting Low-Risk Thyroid Papillary Microcarcinomas Resected Without Observation: Was Immediate Surgery Necessary?. <i>World Journal of Surgery</i> , 2016 , 40, 523-8	3.3	44
285	Cribriform-morular variant of papillary thyroid carcinomacytological and immunocytochemical findings of 18 cases. <i>Diagnostic Cytopathology</i> , 2010 , 38, 890-6	1.4	44
284	BRAF(V600E) mutation analysis in papillary thyroid carcinoma: is it useful for all patients?. <i>World Journal of Surgery</i> , 2014 , 38, 679-87	3.3	43
283	Prognosis of low-risk papillary thyroid carcinoma patients: its relationship with the size of primary tumors. <i>Endocrine Journal</i> , 2012 , 59, 119-25	2.9	43
282	Prevalence and biological behaviour of variants of papillary thyroid carcinoma: experience at a single institute. <i>Pathology</i> , 2008 , 40, 617-22	1.6	43
281	Ratio of serum free triiodothyronine to free thyroxine in GravesQhyperthyroidism and thyrotoxicosis caused by painless thyroiditis. <i>Endocrine Journal</i> , 2005 , 52, 537-42	2.9	43
280	Mutational analysis of the APC gene in cribriform-morula variant of papillary thyroid carcinoma. World Journal of Surgery, 2006 , 30, 775-9	3.3	42
279	Needle tract implantation of papillary thyroid carcinoma after fine-needle aspiration biopsy. <i>World Journal of Surgery</i> , 2005 , 29, 1544-9	3.3	42
278	Alternative surgical strategies and favorable outcomes in patients with medullary thyroid carcinoma in Japan: experience of a single institution. <i>World Journal of Surgery</i> , 2009 , 33, 58-66	3.3	41
277	Morules in cribriform-morular variant of papillary thyroid carcinoma: Immunohistochemical characteristics and distinction from squamous metaplasia. <i>Apmis</i> , 2004 , 112, 275-82	3.4	41
276	Impact of positional changes in neural monitoring endotracheal tube on amplitude and latency of electromyographic response in monitored thyroid surgery: Results from the Porcine Experiment. Head and Neck, 2016, 38 Suppl 1, E1004-8	4.2	41

(2008-2016)

275	Pathological characteristics of low-risk papillary thyroid microcarcinoma with progression during active surveillance. <i>Endocrine Journal</i> , 2016 , 63, 805-810	2.9	40
274	Relationship between prognosis of papillary thyroid carcinoma patient and age: a retrospective single-institution study. <i>Endocrine Journal</i> , 2012 , 59, 399-405	2.9	40
273	Clinicopathologic characteristics and prognosis of diffuse sclerosing variant of papillary thyroid carcinoma in Japan: an 18-year experience at a single institution. <i>World Journal of Surgery</i> , 2009 , 33, 958	3-62	40
272	Natural history of papillary thyroid microcarcinoma: Kinetic analyses on tumor volume during active surveillance and before presentation. <i>Surgery</i> , 2019 , 165, 25-30	3.6	40
271	Prognostic significance of changes in serum thyroglobulin antibody levels of pre- and post-total thyroidectomy in thyroglobulin antibody-positive papillary thyroid carcinoma patients. <i>Endocrine Journal</i> , 2013 , 60, 871-6	2.9	39
270	Prognosis of patients with papillary thyroid carcinoma showing postoperative recurrence to the central neck. <i>World Journal of Surgery</i> , 2011 , 35, 767-72	3.3	39
269	Subclassification of non-solid-type papillary thyroid carcinoma identification of high-risk group in common type. <i>Cancer Science</i> , 2008 , 99, 1908-15	6.9	39
268	Prognosis of patients with benign thyroid diseases accompanied by incidental papillary carcinoma undetectable on preoperative imaging tests. <i>World Journal of Surgery</i> , 2007 , 31, 1672-6	3.3	39
267	The relationship of psychological factors to the prognosis of hyperthyroidism in antithyroid drug-treated patients with GravesQdisease. <i>Clinical Endocrinology</i> , 2003 , 58, 550-5	3.4	39
266	Introducing the reporting system for thyroid fine-needle aspiration cytology according to the new guidelines of the Japan Thyroid Association. <i>Endocrine Journal</i> , 2014 , 61, 539-52	2.9	38
265	Sporadic congenital hyperthyroidism due to a germline mutation in the thyrotropin receptor gene (Leu 512 Gln) in a Japanese patient. <i>Endocrine Journal</i> , 2006 , 53, 735-40	2.9	38
264	Loss of cellular polarity/cohesiveness in the invasive front of papillary thyroid carcinoma, a novel predictor for lymph node metastasis; possible morphological indicator of epithelial mesenchymal transition. <i>Journal of Clinical Pathology</i> , 2011 , 64, 325-9	3.9	36
263	Validation of the Acoustic Voice Quality Index in the Japanese Language. Journal of Voice, 2017, 31, 260	Dæg-20	50 ₅ 9
262	Diagnostic significance of PAX8 in thyroid squamous cell carcinoma. <i>Endocrine Journal</i> , 2015 , 62, 991-5	2.9	35
261	Voice quality after surgical treatment for thyroid cancer. <i>Thyroid</i> , 2013 , 23, 847-53	6.2	35
260	Initial treatment with 15 mg of prednisolone daily is sufficient for most patients with subacute thyroiditis in Japan. <i>Thyroid</i> , 2013 , 23, 269-72	6.2	35
259	Prognosis and prognostic factors of patients with papillary carcinoma showing distant metastasis at surgery (M1 patients) in Japan. <i>Endocrine Journal</i> , 2010 , 57, 523-31	2.9	35
258	Evaluation of a new rapid and fully automated electrochemiluminescence immunoassay for thyrotropin receptor autoantibodies. <i>Thyroid</i> , 2008 , 18, 1157-64	6.2	35

257	3,5,3@Triiodothyronine thyrotoxicosis due to increased conversion of administered levothyroxine in patients with massive metastatic follicular thyroid carcinoma. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 2239-42	5.6	35
256	Relationship of biochemically persistent disease and thyroglobulin-doubling time to age at surgery in patients with papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2013 , 60, 415-421	2.9	34
255	Prognosis and prognostic factors of papillary thyroid carcinoma in patients under 20 years. <i>Endocrine Journal</i> , 2012 , 59, 539-45	2.9	34
254	Unilateral surgery supported by germline RET oncogene mutation analysis in patients with sporadic medullary thyroid carcinoma. <i>World Journal of Surgery</i> , 2000 , 24, 1367-72	3.3	34
253	Ultrasound image analysis using deep learning algorithm for the diagnosis of thyroid nodules. <i>Medicine (United States)</i> , 2019 , 98, e15133	1.8	33
252	The common genetic variant rs944289 on chromosome 14q13.3 associates with risk of both malignant and benign thyroid tumors in the Japanese population. <i>Thyroid</i> , 2015 , 25, 333-40	6.2	33
251	Loss of cellular polarity/cohesiveness in the invasive front of papillary thyroid carcinoma and periostin expression. <i>Cancer Letters</i> , 2009 , 281, 188-95	9.9	33
250	The Safety and Efficacy of Weekly Paclitaxel Administration for Anaplastic Thyroid Cancer Patients: A Nationwide Prospective Study. <i>Thyroid</i> , 2016 , 26, 1293-9	6.2	33
249	TERT promoter mutations and Ki-67 labeling index as a prognostic marker of papillary thyroid carcinomas: combination of two independent factors. <i>Scientific Reports</i> , 2017 , 7, 41752	4.9	32
248	Noninvasive follicular thyroid neoplasm with papillary-like nuclear features: a single-institutional experience in Japan. <i>Endocrine Journal</i> , 2017 , 64, 1149-1155	2.9	32
247	Prognosis and growth activity depend on patient age in clinical and subclinical papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2014 , 61, 205-13	2.9	32
246	Prognostic impact of extrathyroid extension and clinical lymph node metastasis in papillary thyroid carcinoma depend on carcinoma size. <i>World Journal of Surgery</i> , 2010 , 34, 3007-14	3.3	32
245	Outcome of vocal cord function after partial layer resection of the recurrent laryngeal nerve in patients with invasive papillary thyroid cancer. <i>Surgery</i> , 2014 , 155, 184-9	3.6	31
244	Recovery of parathyroid function after total thyroidectomy. <i>Surgery Today</i> , 2000 , 30, 333-8	3	31
243	Prospective randomized study on injury of the external branch of the superior laryngeal nerve during thyroidectomy comparing intraoperative nerve monitoring and a conventional technique. Head and Neck, 2015 , 37, 1456-60	4.2	30
242	Insights and clinical questions about the active surveillance of low-risk papillary thyroid microcarcinomas [Review]. <i>Endocrine Journal</i> , 2016 , 63, 323-8	2.9	30
241	Type 1 and type 2 iodothyronine deiodinases in the thyroid gland of patients with 3,5,3@triiodothyronine-predominant Graves@disease. <i>European Journal of Endocrinology</i> , 2011 , 164, 95-	1605	30
240	Transient vocal cord paralysis after fine-needle aspiration biopsy of thyroid tumor. <i>Thyroid</i> , 2006 , 16, 697-9	6.2	30

(2013-2017)

239	TERT Promoter Mutations Were Not Found in Papillary Thyroid Microcarcinomas That Showed Disease Progression on Active Surveillance. <i>Thyroid</i> , 2017 , 27, 1206-1207	6.2	29	
238	Prognostic values of clinical lymph node metastasis and macroscopic extrathyroid extension in papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2014 , 61, 745-50	2.9	29	
237	Clinical outcomes of patients with papillary thyroid carcinoma after the detection of distant recurrence. World Journal of Surgery, 2010, 34, 2333-7	3.3	29	
236	A novel thyrotropin receptor germline mutation (Asp617Tyr) causing hereditary hyperthyroidism. <i>Endocrine Journal</i> , 2007 , 54, 927-34	2.9	29	
235	Four cases of GravesQdisease which developed after painful HashimotoQ thyroiditis. <i>Internal Medicine</i> , 2006 , 45, 385-9	1.1	29	
234	Changes in serum TSH receptor antibody (TRAb) values in patients with GravesQdisease after total or subtotal thyroidectomy. <i>Endocrine Journal</i> , 2003 , 50, 595-601	2.9	29	
233	Serum concentrations of remnant-like particles in hypothyroid patients before and after thyroxine replacement. <i>Clinical Endocrinology</i> , 2003 , 58, 621-6	3.4	29	
232	Investigation of the prognosis of patients with papillary thyroid carcinoma by tumor size. <i>Endocrine Journal</i> , 2012 , 59, 457-64	2.9	28	
231	Prevalence of TSH receptor and Gsalpha mutations in 45 autonomously functioning thyroid nodules in Japan. <i>Endocrine Journal</i> , 2009 , 56, 791-8	2.9	28	
230	Papillary carcinoma obscured by complication with subacute thyroiditis: sequential ultrasonographic and histopathological findings in five cases. <i>Thyroid</i> , 2008 , 18, 1221-5	6.2	28	
229	Lack of association between BRAF V600E mutation and mitogen-activated protein kinase activation in papillary thyroid carcinoma. <i>Pathology International</i> , 2007 , 57, 12-20	1.8	28	
228	Is an Increase in Thyroid Nodule Volume a Risk Factor for Malignancy?. <i>Thyroid</i> , 2015 , 25, 804-11	6.2	27	
227	Age- and Gender-Specific Risk of Thyroid Cancer in Patients With Familial Adenomatous Polyposis. Journal of Clinical Endocrinology and Metabolism, 2016 , 101, 4611-4617	5.6	27	
226	Needle tract implantation of follicular neoplasm after fine-needle aspiration biopsy: report of a case. <i>Thyroid</i> , 2006 , 16, 1059-62	6.2	27	
225	Stretching exercises to reduce symptoms of postoperative neck discomfort after thyroid surgery: prospective randomized study. <i>World Journal of Surgery</i> , 2005 , 29, 775-9	3.3	27	
224	Intrathyroid epithelial thymoma (ITET) and carcinoma showing thymus-like differentiation (CASTLE): CD5-positive neoplasms mimicking squamous cell carcinoma of the thyroid. <i>Histology and Histopathology</i> , 2013 , 28, 543-56	1.4	27	
223	Frequent BRAF and Absence of TERT Promoter Mutations Characterize Sporadic Pediatric Papillary Thyroid Carcinomas in Japan. <i>Endocrine Pathology</i> , 2017 , 28, 103-111	4.2	26	
222	Ki-67 labeling index is a predictor of postoperative persistent disease and cancer growth and a prognostic indicator in papillary thyroid carcinoma. <i>European Thyroid Journal</i> , 2013 , 2, 57-64	4.2	26	

221	Tumor thrombus of thyroid malignancies in veins: importance of detection by ultrasonography. <i>Thyroid</i> , 2011 , 21, 527-31	6.2	26
220	The Effectiveness of Prophylactic Modified Neck Dissection for Reducing the Development of Lymph Node Recurrence of Papillary Thyroid Carcinoma. <i>World Journal of Surgery</i> , 2017 , 41, 2283-2289	3.3	25
219	Active Surveillance Versus Immediate Surgery: Questionnaire Survey on the Current Treatment Strategy for Adult Patients with Low-Risk Papillary Thyroid Microcarcinoma in Japan. <i>Thyroid</i> , 2019 , 29, 1563-1571	6.2	25
218	Effect of the presence of remnant thyroid tissue on the serum thyroid hormone balance in thyroidectomized patients. <i>European Journal of Endocrinology</i> , 2015 , 173, 333-40	6.5	25
217	2015 American Thyroid Association Guidelines for Thyroid Nodules and Differentiated Thyroid Cancer and Their Implementation in Various Care Settings. <i>Thyroid</i> , 2016 , 26, 319-21	6.2	25
216	Evaluation of surgical results and prediction of prognosis in patients with medullary thyroid carcinoma by analysis of serum calcitonin levels. <i>World Journal of Surgery</i> , 1988 , 12, 610-5	3.3	25
215	Relationship of biochemically persistent disease and thyroglobulin-doubling time to age at surgery in patients with papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2013 , 60, 415-21	2.9	25
214	Preoperative diagnostic algorithm of primary thyroid lymphoma using ultrasound, aspiration cytology, and flow cytometry. <i>Endocrine Journal</i> , 2017 , 64, 859-865	2.9	24
213	Permanent hypoparathyroidism after completion total thyroidectomy as a second surgery: How do we avoid it?. <i>Endocrine Journal</i> , 2014 , 61, 403-8	2.9	24
212	Disturbed metabolism of remnant lipoproteins in patients with subclinical hypothyroidism. <i>American Journal of Medicine</i> , 2004 , 117, 696-9	2.4	24
211	Conservative Surveillance Management of Low-Risk Papillary Thyroid Microcarcinoma. Endocrinology and Metabolism Clinics of North America, 2019 , 48, 215-226	5.5	23
210	Thyroxine treatment may be useful for subclinical hypothyroidism in patients with female infertility. <i>Endocrine Journal</i> , 2015 , 62, 87-92	2.9	23
209	Cytological characteristics of papillary thyroid carcinoma on LBC specimens, compared with conventional specimens. <i>Diagnostic Cytopathology</i> , 2015 , 43, 108-13	1.4	23
208	Rapid bioassay for detection of thyroid-stimulating antibodies using cyclic adenosine monophosphate-gated calcium channel and aequorin. <i>European Thyroid Journal</i> , 2015 , 4, 14-9	4.2	23
207	Distant metastasis at diagnosis and large tumor size are significant prognostic factors of widely invasive follicular thyroid carcinoma. <i>Endocrine Journal</i> , 2013 , 60, 829-33	2.9	23
206	Distribution of IgG4- and/or IgG-positive plasma cells in Hashimoto@thyroiditis: an immunohistochemical study. <i>Pathobiology</i> , 2010 , 77, 267-72	3.6	23
205	Diagnosis of follicular carcinoma of the thyroid: role of sonography in preoperative diagnosis of follicular nodules. <i>Journal of Medical Ultrasonics (2001)</i> , 2005 , 32, 153-8	1.4	23
204	Cribriform-morular variant of papillary thyroid carcinoma. Report of a case showing morules with peculiar nuclear clearing. <i>Acta Cytologica</i> , 2004 , 48, 431-6	3	22

203	Metastatic carcinoma to the thyroid gland from renal cell carcinoma: role of ultrasonography in preoperative diagnosis. <i>Thyroid Research</i> , 2015 , 8, 4	2.4	21
202	Prognostic significance of patient age in minimally and widely invasive follicular thyroid carcinoma: investigation of three age groups. <i>Endocrine Journal</i> , 2014 , 61, 265-71	2.9	21
201	Biological behavior and prognosis of encapsulated papillary carcinoma of the thyroid: experience of a Japanese hospital for thyroid care. <i>World Journal of Surgery</i> , 2008 , 32, 1789-94	3.3	21
200	Exploratory analysis of prognostic factors for lenvatinib in radioiodine-refractory differentiated thyroid cancer. <i>Head and Neck</i> , 2019 , 41, 3023-3032	4.2	20
199	Prognosis and prognostic factors of differentiated thyroid carcinoma after the appearance of metastasis refractory to radioactive iodine therapy. <i>Endocrine Journal</i> , 2014 , 61, 821-4	2.9	20
198	Changes in serum thyroglobulin antibody levels as a dynamic prognostic factor for early-phase recurrence of thyroglobulin antibody-positive papillary thyroid carcinoma after total thyroidectomy. <i>Endocrine Journal</i> , 2014 , 61, 961-5	2.9	20
197	Active Surveillance as First-Line Management of Papillary Microcarcinoma. <i>Annual Review of Medicine</i> , 2019 , 70, 369-379	17.4	20
196	The Acoustic Voice Quality Index Version 03.01 for the Japanese-speaking Population. <i>Journal of Voice</i> , 2019 , 33, 125.e1-125.e12	1.9	20
195	Fine-needle aspiration cytology for medullary thyroid carcinoma: a single institutional experience in Japan. <i>Endocrine Journal</i> , 2017 , 64, 1099-1104	2.9	19
194	mRNA Expression as a Novel Prognostic Marker in Papillary Thyroid Carcinomas. <i>Thyroid</i> , 2019 , 29, 110	5 -1.1 14	19
193	Serum calcitonin levels with calcium loading tests before and after total thyroidectomy in patients with thyroid diseases other than medullary thyroid carcinoma. <i>Endocrine Journal</i> , 2011 , 58, 217-21	2.9	19
192	Significance of prophylactic modified radical neck dissection for patients with low-risk papillary thyroid carcinoma measuring 1.1-3.0 cm: first report of a trial at Kuma Hospital. <i>Surgery Today</i> , 2011 , 41, 1486-91	3	19
191	Thyroid ultrasonography. World Journal of Surgery, 2010 , 34, 1171-80	3.3	19
190	Results of a phase I clinical study using dendritic cell vaccinations for thyroid cancer. <i>Thyroid</i> , 2007 , 17, 53-8	6.2	19
189	Association of Patient Age With Progression of Low-risk Papillary Thyroid Carcinoma Under Active Surveillance: A Systematic Review and Meta-analysis. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2020 , 146, 552-560	3.9	19
188	Kinetic Analysis of Growth Activity in Enlarging Papillary Thyroid Microcarcinomas. <i>Thyroid</i> , 2019 , 29, 1765-1773	6.2	18
187	Static and dynamic prognostic factors of papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2014 , 61, 1145	- 521 .9	18
186	Intrathyroidal epithelial thymoma/carcinoma showing thymus-like differentiation; comparison with thymic lymphoepithelioma-like carcinoma and a possibility of development from a multipotential stem cell. <i>Apmis</i> , 2013 , 121, 523-30	3.4	18

185	Prognostic factors of papillary thyroid carcinoma vary according to sex and patient age. <i>World Journal of Surgery</i> , 2011 , 35, 2684-90	3.3	18
184	Local prognosis of patients with papillary thyroid carcinoma who were intra-operatively diagnosed as having minimal invasion of the trachea: a 17-year experience in a single institute. <i>Asian Journal of Surgery</i> , 2009 , 32, 102-8	1.6	18
183	Innervation of the cricothyroid muscle by the recurrent laryngeal nerve. <i>Head and Neck</i> , 2016 , 38 Suppl 1, E441-5	4.2	17
182	Prognostic significance of young age in papillary thyroid carcinoma: analysis of 5,733 patients with 150 monthsQmedian follow-up. <i>Endocrine Journal</i> , 2014 , 61, 491-7	2.9	17
181	Attenuation of postmenopausal bone loss in patients with transient hypoparathyroidism after total thyroidectomy. <i>World Journal of Surgery</i> , 2013 , 37, 2860-5	3.3	17
180	Static Prognostic Factors and Appropriate Surgical Designs for Patients with Medullary Thyroid Carcinoma: The Second Report from a Single-Institution Study in Japan. <i>World Journal of Surgery</i> , 2018 , 42, 3954-3966	3.3	16
179	Moderate Frequency of Anti-Thyroglobulin Antibodies in the Early Phase of Subacute Thyroiditis. <i>European Thyroid Journal</i> , 2019 , 8, 268-272	4.2	16
178	Anaplastic thyroid carcinomas incidentally found on postoperative pathological examination. <i>World Journal of Surgery</i> , 2014 , 38, 2311-6	3.3	16
177	Functional characterization of the novel BRAF complex mutation, BRAF(V600delinsYM), identified in papillary thyroid carcinoma. <i>International Journal of Cancer</i> , 2013 , 132, 738-43	7.5	16
176	Prognostic Significance of the Proportion of Tall Cell Components in Papillary Thyroid Carcinoma. <i>World Journal of Surgery</i> , 2017 , 41, 742-747	3.3	16
175	Immunoglobulin G4 thyroiditis in a GravesQdisease patient with a large goiter developing hypothyroidism. <i>Thyroid</i> , 2013 , 23, 1496-7	6.2	16
174	Role of ultrasonography in patients with cytologically follicular thyroid tumor. <i>Auris Nasus Larynx</i> , 2011 , 38, 508-11	2.2	16
173	Cytologic findings of primary thyroid MALT lymphoma with extreme plasma cell differentiation: FNA cytology of two cases. <i>Diagnostic Cytopathology</i> , 2009 , 37, 815-9	1.4	16
172	Lateral mobilization of the recurrent laryngeal nerve to facilitate tracheal surgery in patients with thyroid cancer invading the trachea near Berry@ligament. <i>World Journal of Surgery</i> , 2007 , 31, 2081-4	3.3	16
171	Paediatric follicular thyroid carcinoma - indolent cancer with low prevalence of RAS mutations and absence of PAX8-PPARG fusion in a Japanese population. <i>Histopathology</i> , 2017 , 71, 760-768	7.3	15
170	Papillary thyroid carcinoma with desmoid-type fibromatosis: A clinical, pathological, and immunohistochemical study of 14 cases. <i>Endocrine Journal</i> , 2017 , 64, 1017-1023	2.9	15
169	Reappraisal of "cyst fluid only" on thyroid fine-needle aspiration cytology. <i>Endocrine Journal</i> , 2017 , 64, 759-765	2.9	15
168	Immunohistochemical detection of NRAS(Q61R) protein in follicular-patterned thyroid tumors. <i>Human Pathology</i> , 2016 , 53, 51-7	3.7	15

167	Gene rearrangement of immunoglobulin as a marker of thyroid lymphoma. <i>World Journal of Surgery</i> , 1998 , 22, 558-61	3.3	15	
166	Ansa-recurrent nerve anastomosis for vocal cord paralysis due to mediastinal lesions. <i>Annals of Thoracic Surgery</i> , 1994 , 57, 1020-1	2.7	15	
165	Tumor protrusion with intensive blood signals on ultrasonography is a strongly suggestive finding of follicular thyroid carcinoma. <i>Medical Ultrasonography</i> , 2016 , 18, 25-9	1.4	15	
164	Innervation of the cricothyroid muscle by extralaryngeal branches of the recurrent laryngeal nerve. <i>Laryngoscope</i> , 2016 , 126, 1157-62	3.6	14	
163	Cytological findings for the diagnosis of primary thyroid mucosa-associated lymphoid tissue lymphoma by fine needle aspiration. <i>Acta Cytologica</i> , 2015 , 59, 26-36	3	13	
162	Diagnostic value of GATA-3 in cytological identification of parathyroid tissues. <i>Endocrine Journal</i> , 2016 , 63, 621-6	2.9	13	
161	Reference values of serum calcitonin with calcium stimulation tests by electrochemiluminescence immunoassay before/after total thyroidectomy in Japanese patients with thyroid diseases other than medullary thyroid carcinoma. <i>Endocrine Journal</i> , 2016 , 63, 627-32	2.9	13	
160	Surgical Methods and Experiences of Surgeons did not Significantly Affect the Recovery in Phonation Following Reconstruction of the Recurrent Laryngeal Nerve. <i>World Journal of Surgery</i> , 2016 , 40, 2948-2955	3.3	13	
159	Calcifications in Thyroid Tumors on Ultrasonography: Calcification Types and Relationship with Histopathological Type. <i>Ultrasound International Open</i> , 2018 , 4, E45-E51	2.1	13	
158	Evaluation of cytologically benign solitary thyroid nodules by ultrasonography: a retrospective analysis of 1877 cases. <i>Auris Nasus Larynx</i> , 2013 , 40, 308-11	2.2	13	
157	Genotype Analyses in the Japanese and Belarusian Populations Reveal Independent Effects of rs965513 and rs1867277 but Do Not Support the Role of FOXE1 Polyalanine Tract Length in Conferring Risk for Papillary Thyroid Carcinoma. <i>Thyroid</i> , 2017 , 27, 224-235	6.2	13	
156	Final pathology findings after immediate or delayed surgery in patients with cytologically benign or follicular thyroid nodules. <i>World Journal of Surgery</i> , 2011 , 35, 558-62	3.3	13	
155	Right-sided aortic arch and aberrant left subclavian artery with or without a left nonrecurrent inferior laryngeal nerve. <i>Head and Neck</i> , 2016 , 38, E2508-11	4.2	13	
154	Thyroid Fine-Needle Aspiration and Smearing Techniques. VideoEndocrinology, 2018, 5,	1.6	13	
153	Calcitonin doubling time in medullary thyroid carcinoma after the detection of distant metastases keenly predicts patients Quarcinoma death. <i>Endocrine Journal</i> , 2016 , 63, 663-7	2.9	12	
152	Laryngeal approach to the recurrent laryngeal nerve involved by thyroid cancer at the ligament of Berry. <i>Surgery</i> , 2012 , 152, 57-60	3.6	12	
151	Cytologic findings and differential diagnoses of primary thyroid MALT lymphoma with striking plasma cell differentiation and amyloid deposition. <i>Diagnostic Cytopathology</i> , 2014 , 42, 73-7	1.4	12	
150	Characteristic sonographic features of cribriform papillary thyroid carcinoma for differentiation from other thyroid nodules. <i>Journal of Medical Ultrasonics (2001)</i> , 2015 , 42, 83-7	1.4	11	

149	Graves@isease Patients with Persistent Hyperthyroidism and Diffuse Lymphoplasmacytic Infiltration in the Thyroid Show No Histopathological Compatibility with IgG4-Related Disease. <i>PLoS ONE</i> , 2015 , 10, e0134143	3.7	11
148	Undetectable serum thyroglobulin levels in patients with medullary thyroid carcinoma after total thyroidectomy without radioiodine ablation. <i>Thyroid</i> , 2012 , 22, 680-2	6.2	11
147	A case with diffuse sclerosing variant of papillary carcinoma of the thyroid: characteristic features on ultrasonography. <i>Journal of Medical Ultrasonics (2001)</i> , 2006 , 33, 159-61	1.4	11
146	Thyroid nodular lesion: analysis of cancer risk based on Kuma Hospital experience. <i>Pathology International</i> , 2003 , 53, 579-83	1.8	11
145	Is Less More? A Microsimulation Model Comparing Cost-effectiveness of the Revised American Thyroid Association@ 2015 to 2009 Guidelines for the Management of Patients With Thyroid Nodules and Differentiated Thyroid Cancer. <i>Annals of Surgery</i> , 2020 , 271, 765-773	7.8	11
144	Diagnosis and surgical indications of oxyphilic follicular tumors in Japan: Surgical specimens and cytology. <i>Endocrine Journal</i> , 2016 , 63, 977-982	2.9	10
143	Prevalence of thyrotropin receptor germline mutations and clinical courses in 89 hyperthyroid patients with diffuse goiter and negative anti-thyrotropin receptor antibodies. <i>Thyroid</i> , 2014 , 24, 789-95	6.2	10
142	The thyroid function of GravesQlisease patients is aggravated by depressive personality during antithyroid drug treatment. <i>BioPsychoSocial Medicine</i> , 2011 , 5, 9	2.8	10
141	A Novel Germline Mutation of KEAP1 (R483H) Associated with a Non-Toxic Multinodular Goiter. <i>Frontiers in Endocrinology</i> , 2016 , 7, 131	5.7	10
140	Periodic Granulocyte Count Measuring Is Useful for Detecting Asymptomatic Agranulocytosis in Antithyroid Drug-Treated Patients with Graves@isease. <i>European Thyroid Journal</i> , 2016 , 5, 253-260	4.2	10
139	Impact of the modification of the diagnostic criteria in the 2017 Bethesda System for Reporting Thyroid Cytopathology: a report of a single institution in Japan. <i>Endocrine Journal</i> , 2018 , 65, 1193-1198	2.9	10
138	Concept and design of a nationwide prospective feasibility/efficacy/safety study of weekly paclitaxel for patients with pathologically confirmed anaplastic thyroid cancer (ATCCJ-PTX-P2). <i>BMC Cancer</i> , 2015 , 15, 475	4.8	9
137	Comparative histopathological analysis of sporadic pediatric papillary thyroid carcinoma from Japan and Ukraine. <i>Endocrine Journal</i> , 2017 , 64, 977-993	2.9	9
136	Airtight tracheocutaneostomy after window resection of the trachea for invasive papillary thyroid carcinoma: experience of 109 cases. <i>World Journal of Surgery</i> , 2014 , 38, 660-6	3.3	9
135	Prognosis and prognostic factors of patients with papillary thyroid carcinoma requiring resection of recurrent laryngeal nerve due to carcinoma extension. <i>Endocrine Journal</i> , 2012 , 59, 247-52	2.9	9
134	Subclinical nonautoimmune hyperthyroidism in a family segregates with a thyrotropin receptor mutation with weakly increased constitutive activity. <i>Thyroid</i> , 2010 , 20, 1307-14	6.2	9
133	Germline RET mutation carriers in Japanese patients with apparently sporadic medullary thyroid carcinoma: A single institution experience. <i>Auris Nasus Larynx</i> , 2016 , 43, 551-5	2.2	9
132	Appropriateness of the revised Japanese guidelinesQisk classification for the prognosis of papillary thyroid carcinoma: a retrospective analysis of 5,845 papillary thyroid carcinoma patients. <i>Endocrine Journal</i> , 2019 , 66, 127-134	2.9	9

131	GravesQdisease and mental disorders. Journal of Clinical and Translational Endocrinology, 2020, 19, 100	2 0:7 .4	9
130	Risk of malignancy and clinical outcomes of cyst fluid only nodules in the thyroid based on ultrasound and aspiration cytology. <i>Diagnostic Cytopathology</i> , 2020 , 48, 30-34	1.4	9
129	Thrombotic Microangiopathy with Severe Proteinuria Induced by Lenvatinib for Radioactive Iodine-Refractory Papillary Thyroid Carcinoma. <i>Case Reports in Oncology</i> , 2018 , 11, 735-741	1	9
128	Immunohistochemical and Molecular Analyses Focusing on Mesenchymal Cells in Papillary Thyroid Carcinoma with Desmoid-Type Fibromatosis. <i>Pathobiology</i> , 2018 , 85, 300-303	3.6	9
127	Comparison of thyroglobulin and thyroid peroxidase antibodies measured by five different kits in autoimmune thyroid diseases. <i>Endocrine Journal</i> , 2017 , 64, 955-961	2.9	8
126	Re-evaluation of MIB-1 immunostaining for diagnosing hyalinizing trabecular tumour of the thyroid: semi-automated techniques with manual antigen retrieval are more accurate than fully automated techniques. <i>Endocrine Journal</i> , 2018 , 65, 239-244	2.9	8
125	Thyroid sclerosing mucoepidermoid carcinoma with eosinophilia distinct from the salivary type. <i>Endocrine Journal</i> , 2018 , 65, 427-436	2.9	8
124	Chromophobe renal cell carcinoma-like thyroid carcinoma: A novel clinicopathologic entity possibly associated with tuberous sclerosis complex. <i>Endocrine Journal</i> , 2017 , 64, 843-850	2.9	8
123	Optimal needle size for thyroid fine needle aspiration cytology. <i>Endocrine Journal</i> , 2019 , 66, 143-147	2.9	8
122	Cribriform-Morular Variant of Papillary Thyroid Carcinoma Shows High Ki-67 Labeling Indices, despite Its Excellent Prognosis. <i>Pathobiology</i> , 2019 , 86, 248-253	3.6	7
121	Routine Chest Computed Tomography at Presentation Does Not Identify Distant Metastasis in cT1aN0 Papillary Thyroid Carcinoma. <i>Thyroid</i> , 2020 , 30, 1620-1624	6.2	7
120	Comparative frequency of four different types of pregnancy-associated thyrotoxicosis in a single thyroid centre. <i>Thyroid Research</i> , 2017 , 10, 4	2.4	7
119	Excellent Prognosis of Central Lymph Node Recurrence-Free Survival for cN0M0 Papillary Thyroid Carcinoma Patients Who Underwent Routine Prophylactic Central Node Dissection. <i>World Journal of Surgery</i> , 2018 , 42, 2462-2468	3.3	7
118	Identification of Cytological Features Distinguishing Mucosa-Associated Lymphoid Tissue Lymphoma from Reactive Lymphoid Proliferation Using Thyroid Liquid-Based Cytology. <i>Acta</i> <i>Cytologica</i> , 2018 , 62, 93-98	3	7
117	Kinetic analyses of changes in serum TSH receptor antibody values after total thyroidectomy in patients with GravesQdisease. <i>Endocrine Journal</i> , 2016 , 63, 179-85	2.9	7
116	Prognostic impact of Ki-67 labeling index in minimally invasive follicular thyroid carcinoma. <i>Endocrine Journal</i> , 2016 , 63, 913-917	2.9	7
115	Papillary thyroid carcinoma with honeycomb-like multiple small cysts: characteristic features on ultrasonography. <i>European Thyroid Journal</i> , 2013 , 2, 270-4	4.2	7
114	Prognostic significance of carcinoma extension from primary lesions and metastatic nodes in papillary thyroid carcinoma: appropriateness of three subdivisions of extension. <i>Endocrine Journal</i> , 2011 , 58, 961-7	2.9	7

113	Prevalence and diagnostic significance of noninvasive follicular thyroid neoplasm with papillary-like nuclear features among tumors previously diagnosed as follicular adenoma: a single-institutional study in Japan. <i>Endocrine Journal</i> , 2020 , 67, 1071-1075	2.9	7
112	Hashimoto@Thyroiditis with Increased IgG4-Positive Plasma Cells: Using Thyroid-Specific Diagnostic Criteria May Identify Early Phase IgG4 Thyroiditis. <i>Thyroid</i> , 2020 , 30, 251-261	6.2	7
111	Needle Tract Implantation Following Fine-Needle Aspiration of Thyroid Cancer. <i>World Journal of Surgery</i> , 2020 , 44, 378-384	3.3	7
110	Thyroid Cancer Active Surveillance Program Retention and Adherence in Japan. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2021 , 147, 77-84	3.9	7
109	"Nodule in Nodule" on Thyroid Ultrasonography: Possibility of Follicular Carcinoma Transformed from Benign Thyroid Tumor. <i>European Thyroid Journal</i> , 2017 , 6, 101-107	4.2	6
108	Role of immediate recurrent laryngeal nerve reconstruction in surgery for thyroid cancers with fixed vocal cords. <i>Head and Neck</i> , 2017 , 39, 427-431	4.2	6
107	Important cytological findings for distinction between follicular variant and conventional papillary thyroid carcinoma, including noninvasive follicular thyroid tumors with papillary-like nuclear features. <i>Endocrine Journal</i> , 2019 , 66, 475-483	2.9	6
106	Serum calcitonin reference values for calcium stimulation tests by electrochemiluminescence immunoassay in Japanese men with non-medullary thyroid carcinoma. <i>Surgery Today</i> , 2018 , 48, 223-228	3 ³	6
105	Active surveillance of low-risk papillary thyroid microcarcinomas in Japan and other countries: a review. <i>Expert Review of Endocrinology and Metabolism</i> , 2020 , 15, 5-12	4.1	6
104	Active surveillance of low-risk papillary thyroid microcarcinomas. <i>Gland Surgery</i> , 2020 , 9, 1663-1673	2.2	6
103	Federated Learning for Thyroid Ultrasound Image Analysis to Protect Personal Information: Validation Study in a Real Health Care Environment. <i>JMIR Medical Informatics</i> , 2021 , 9, e25869	3.6	6
102	Diagnostic clues indicating tall cell variants of papillary thyroid carcinoma in fine needle aspiration. <i>Diagnostic Cytopathology</i> , 2019 , 47, 452-457	1.4	6
101	Prognostic Value of Extranodal Tumor Extension in Papillary Thyroid Carcinoma: Proposal for Upstaging of Cases with Extranodal Tumor Extension. <i>World Journal of Surgery</i> , 2020 , 44, 638-643	3.3	6
100	Marked Decrease Over Time in Conversion Surgery After Active Surveillance of Low-Risk Papillary Thyroid Microcarcinoma. <i>Thyroid</i> , 2021 , 31, 217-223	6.2	6
99	Fine-needle aspiration of parathyroid adenomas: Indications as a diagnostic approach. <i>Diagnostic Cytopathology</i> , 2021 , 49, 70-76	1.4	6
98	Calcitonin measurement in fine-needle aspirate washout fluid by electrochemiluminescence immunoassay for thyroid tumors. <i>Thyroid Research</i> , 2018 , 11, 15	2.4	6
97	Prognostic value of the 8 tumor-node-metastasis classification for follicular carcinoma and poorly differentiated carcinoma of the thyroid in Japan. <i>Endocrine Journal</i> , 2018 , 65, 621-627	2.9	6
96	Successful treatment switch from lenvatinib to sorafenib in a patient with radioactive iodine-refractory differentiated thyroid cancer intolerant to lenvatinib due to severe proteinuria. <i>Auris Nasus Larynx</i> , 2018 , 45, 1249-1252	2.2	6

95	A Novel Diagnostic Method for Thyroid Follicular Tumors Based on Immunofluorescence Analysis of p53-Binding Protein 1 Expression: Detection of Genomic Instability. <i>Thyroid</i> , 2019 , 29, 657-665	6.2	5	
94	Type 1 and type 2 iodothyronine deiodinases in the thyroid gland of patients with huge goitrous Hashimoto@thyroiditis. <i>Endocrine</i> , 2019 , 64, 584-590	4	5	
93	Cytoplasmic Lipid Accumulation Characteristic of the Cribriform Variant of Papillary Thyroid Carcinoma. <i>Pathobiology</i> , 2017 , 84, 251-257	3.6	5	
92	Prognostic value of the 8 edition of the tumor-node-metastasis classification for patients with papillary thyroid carcinoma: a single-institution study at a high-volume center in Japan. <i>Endocrine Journal</i> , 2018 , 65, 707-716	2.9	5	
91	Histopathological diagnoses of "accessory" thyroid nodules diagnosed as benign by fine-needle aspiration cytology and ultrasonography. <i>Thyroid</i> , 2012 , 22, 299-303	6.2	5	
90	Clinical significance and prognostic impact of subcutaneous or intrastrap muscular recurrence of papillary thyroid carcinoma. <i>Journal of Thyroid Research</i> , 2012 , 2012, 819797	2.6	5	
89	Transient hyperthyroidism after withdrawal of antithyroid drugs in patients with GravesQdisease. <i>Endocrine Journal</i> , 2004 , 51, 213-7	2.9	5	
88	Spontaneous Deceleration and Acceleration of Growth Rate in Medullary Thyroid Carcinomas Suggested by Changes in Calcitonin Doubling Times Over Long-Term Surveillance. <i>World Journal of Surgery</i> , 2019 , 43, 504-512	3.3	5	
87	Aggressive cribriform-morular variant of papillary thyroid carcinoma: Report of an unusual case with pulmonary metastasis displaying poorly differentiated features. <i>Pathology International</i> , 2018 , 68, 700-705	1.8	5	
86	Flow cytometric, gene rearrangement, and karyotypic analyses of 110 cases of primary thyroid lymphoma: a single-institutional experience in Japan. <i>Endocrine Journal</i> , 2019 , 66, 1083-1091	2.9	4	
85	Papillary Thyroid Carcinoma in Ukraine After Chernobyl and in Japan After Fukushima: Different Histopathological Scenarios. <i>Thyroid</i> , 2021 , 31, 1322-1334	6.2	4	
84	Case report: exceptionally rapid growth character of hobnail variant of papillary thyroid carcinoma: a report of four cases. <i>Endocrine Journal</i> , 2020 , 67, 1047-1053	2.9	4	
83	Older Age Significantly Affects Mortality of Patients with Papillary Thyroid Carcinoma Only When They Have High-Risk Features. <i>World Journal of Surgery</i> , 2020 , 44, 1885-1891	3.3	4	
82	Papillary thyroid carcinomas are highly obscured by inflammatory hypoechoic regions caused by subacute thyroiditis: a longitudinal evaluation of 710 patients using ultrasonography. <i>Endocrine Journal</i> , 2020 , 67, 569-574	2.9	4	
81	The thyroid cancer policy model: A mathematical simulation model of papillary thyroid carcinoma in The U.S. population. <i>PLoS ONE</i> , 2017 , 12, e0177068	3.7	4	
80	Sialic Acid-Binding Immunoglobulin-Like Lectin1 as a Novel Predictive Biomarker for Relapse in Graves@isease: A Multicenter Study. <i>Thyroid</i> , 2018 , 28, 50-59	6.2	4	
79	Malignant melanoma arising in melanin-producing medullary thyroid carcinoma. <i>International Journal of Surgery Case Reports</i> , 2016 , 20, 118-22	0.8	4	
78	Spontaneous Slowing and Regressing of Tumor Growth in Childhood/Adolescent Papillary Thyroid Carcinomas Suggested by the Postoperative Thyroglobulin-Doubling Time. <i>Journal of Thyroid Research</i> , 2018 , 2018, 6470251	2.6	4	

77	Control of Lung Metastases and Colon Polyposis with Lenvatinib Therapy in a Patient with Cribriform-Morular Variant of Papillary Thyroid Carcinoma and an Gene Mutation: A Case Study. <i>Thyroid</i> , 2019 , 29, 1511-1517	6.2	4
76	Thyroid function related symptoms during levothyroxine monotherapy in athyreotic patients. <i>Endocrine Journal</i> , 2019 , 66, 953-960	2.9	4
75	Occurrence of thyroxine tablet (Thyradin S(\square)) - induced liver dysfunction in a patient with subclinical hypothyroidism. <i>Endocrine Journal</i> , 2015 , 62, 719-24	2.9	4
74	Acoustic Breathiness Index for the Japanese-Speaking Population: Validation Study and Exploration of Affecting Factors. <i>Journal of Speech, Language, and Hearing Research</i> , 2019 , 62, 2617-263	3 ^{2.8}	4
73	Quality of Life in Patients with Low-Risk Papillary Thyroid Microcarcinoma: Active Surveillance Versus Immediate Surgery. <i>Endocrine Practice</i> , 2020 , 26, 1451-1457	3.2	4
72	High Prevalence of DICER1 Mutations and Low Frequency of Gene Fusions in Pediatric Follicular-Patterned Tumors of the Thyroid. <i>Endocrine Pathology</i> , 2021 , 32, 336-346	4.2	4
71	Differentiating between benign follicular nodules and follicular neoplasms in thyroid liquid-based cytology preparations. <i>Diagnostic Cytopathology</i> , 2016 , 44, 659-64	1.4	4
70	Subclassification of Tumor Extension and Nodal Metastasis in Papillary Thyroid Cancer to Improve Prognostic Accuracy of the Eighth Edition of the Tumor-Node-Metastasis Classification. <i>World Journal of Surgery</i> , 2020 , 44, 336-345	3.3	4
69	Identification of Recurrent TERT Promoter Mutations in Intrathyroid Thymic Carcinomas. <i>Endocrine Pathology</i> , 2020 , 31, 274-282	4.2	3
68	Partial prediction of postpartum GravesQhyrotoxicosis by sensitive bioassay for thyroid-stimulating antibody measured in early pregnancy. <i>Endocrine Journal</i> , 2016 , 63, 929-932	2.9	3
67	Derivation of thyroid lymphoepithelial cysts from follicular cells. <i>Endocrine Journal</i> , 2018 , 65, 579-586	2.9	3
66	Quantitative measurement of anti-TSH receptor antibodies is useful for the evaluation of treatment effect. <i>Thyroid</i> , 2007 , 17, 85-6	6.2	3
65	Utility of monoclonal PAX8 antibody for distinguishing intrathyroid thymic carcinoma from follicular cell-derived thyroid carcinoma. <i>Endocrine Journal</i> , 2018 , 65, 1171-1175	2.9	3
64	Sonographic Evaluation of Nodules Newly Detected in the Neck After Thyroidectomy: Suture Granuloma Versus Recurrent Carcinoma. <i>Ultrasound International Open</i> , 2018 , 4, E124-E130	2.1	3
63	Can Ultrasound Alone Predict Papillary Thyroid Carcinoma with Desmoid-Type Fibromatosis? A Retrospective Analysis of 13 Cases, Focusing on the Stromal Area. <i>Ultrasound International Open</i> , 2018 , 4, E39-E44	2.1	3
62	Warthin-like papillary thyroid carcinoma with immunoglobulin G4-positive plasma cells possibly related to Hashimoto@thyroiditis. <i>Endocrine Journal</i> , 2018 , 65, 175-180	2.9	3
61	The association between thyroid hormone balance and thyroid volume in patients with Hashimoto thyroiditis. <i>Endocrine Journal</i> , 2019 , 66, 763-768	2.9	2
60	Characteristics and natural course of hypoechoic thyroid lesions diagnosed as possible thyroid lymphomas by fine needle aspiration cytology. <i>Thyroid Research</i> , 2018 , 11, 8	2.4	2

59	Serum Thyroid Hormone Balance in Levothyroxine Monotherapy-Treated Patients with Atrophic Thyroid After Radioiodine Treatment for Graves Disease. <i>Thyroid</i> , 2019 , 29, 1364-1370	6.2	2
58	Up-regulation of urinary-type plasminogen activator correlates with high-risk papillary thyroid carcinoma with BRAF(V600E) mutation and its possible molecular mechanism. <i>Pathology Research and Practice</i> , 2014 , 210, 733-8	3.4	2
57	Tandem Germline RET Mutations in a Family Pathogenetic for Multiple Endocrine Neoplasia 2B, Confirmed by a Natural Experiment. <i>European Thyroid Journal</i> , 2014 , 3, 272-7	4.2	2
56	PSYCHOLOGICAL CHARACTERISTICS OF THE NEO-FFI AND THE TREE DRAWING TEST IN PATIENTS WITH THYROID DISEASE. <i>Psychologia</i> , 2013 , 56, 138-153	0.2	2
55	CD44v8-10 and CD44s Are Age-dependently Expressed in Primary Cultured Papillary Thyroid Carcinoma Cells and Are Associated with Cell Proliferation. <i>Kobe Journal of Medical Sciences</i> , 2019 , 65, E1-E9	0.6	2
54	The Japanese reporting system for thyroid aspiration cytology 2019 (JRSTAC2019). <i>Gland Surgery</i> , 2020 , 9, 1653-1662	2.2	2
53	Cytological findings of primary thyroid MALT lymphoma with extreme plasma cell differentiation in fine needle aspiration specimens. <i>The Journal of the Japanese Society of Clinical Cytology</i> , 2013 , 52, 507-	517	2
52	Intraoperative Neural Injury Management: Transection and Segmental Defects 2016 , 253-258		2
51	Findings and significance of mountain range-like clusters derived from follicular colonization for the diagnosis of MALT lymphoma. <i>The Journal of the Japanese Society of Clinical Cytology</i> , 2013 , 52, 428	-436	2
50	Tumor Volume Kinetic Analyses Might Explain Excellent Prognoses in Young Patients with Papillary Thyroid Carcinoma. <i>Journal of Thyroid Research</i> , 2020 , 2020, 4652767	2.6	2
49	53BP1 expression as a biomarker to differentiate thyroid follicular tumors. <i>Endocrine Connections</i> , 2021 , 10, 309-315	3.5	2
48	Sorafenib and Lenvatinib Treatment for Metastasis/Recurrence of Radioactive Iodine-refractory Differentiated Thyroid Carcinoma. <i>In Vivo</i> , 2021 , 35, 1057-1064	2.3	2
47	Prognostic Significance of Neutrophil-to-Lymphocyte Ratio in Differentiated Thyroid Carcinoma Having Distant Metastasis: A Comparison With Thyroglobulin-doubling Rate and Tumor Volume-doubling Rate. <i>In Vivo</i> , 2021 , 35, 1125-1132	2.3	2
46	Minute Hyalinizing Trabecular Tumor Misinterpreted as a Blood Vessel on Ultrasonography: A Case Report. <i>Ultrasound International Open</i> , 2017 , 3, E42-E44	2.1	1
45	Thyroid Tubercle of Zuckerkandl May Not Arise from the Ultimobranchial Body: Results from Histological Analysis. <i>Pathobiology</i> , 2020 , 87, 193-197	3.6	1
44	Evaluation of ^ ^ldquo;Benign^ ^rdquo; in Thyroid Bethesda System. <i>The Journal of the Japanese Society of Clinical Cytology</i> , 2014 , 53, 257-263	О	1
43	Long-Term Follow-Up of a Patient with Sporadic Nonautoimmune Hyperthyroidism Due to a Thyrotropin-Receptor Mutation (D619G). <i>AACE Clinical Case Reports</i> , 2018 , 4, 84-89	0.7	1
42	Active Surveillance as the Initial Course of Action in Low-Risk Papillary Microcarcinoma 2017 , 135-141		1

41	CA19-9 May Be a Prognostic Factor for Medullary Thyroid Carcinoma. <i>Clinical Thyroidology</i> , 2020 , 32, 24-26	0.1	1
40	Subset analysis of the Japanese risk classification guidelines for papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2020 , 67, 275-282	2.9	1
39	Histological alterations following fine-needle aspiration for parathyroid adenoma: Incidence and diagnostic problems. <i>Pathology International</i> , 2021 , 71, 400-405	1.8	1
38	Prognostic significance of vascular invasion and cell-proliferation activity in widely invasive follicular carcinoma of the thyroid. <i>Endocrine Journal</i> , 2021 , 68, 881-888	2.9	1
37	Phantom Nodules Detected by Ultrasound Examination of the Neck: The Possibility of Ectopic Cervical Thymic Tissue in Adults. <i>Ultrasound International Open</i> , 2018 , 4, E119-E123	2.1	1
36	Clinical significance and cytological detection of tracheal puncture following thyroid fine-needle aspiration: A retrospective study. <i>Diagnostic Cytopathology</i> , 2021 , 49, 1116-1121	1.4	1
35	Poorly Differentiated Thyroid Carcinoma Coexisting with Graves Qisease Involving T3 Thyrotoxicosis due to Increased D1 and D2 Activities. <i>Thyroid</i> , 2021 , 31, 1592-1596	6.2	1
34	Prevalence and Treatment Outcomes of Marine-Lenhart Syndrome in Japan <i>European Thyroid Journal</i> , 2021 , 10, 461-467	4.2	O
33	Clinicopathological features of primary thyroid Burkitt@lymphoma: a systematic review and meta-analysis. <i>Diagnostic Pathology</i> , 2020 , 15, 13	3	О
32	Thyroid Lymphoepithelial Cysts Mimicking Calcified or Solid Nodules on Ultrasonography. <i>Ultrasound International Open</i> , 2019 , 5, E60-E64	2.1	O
31	Ultrasound findings of noninvasive follicular thyroid neoplasm with papillary-like nuclear features compared with those of follicular variant of papillary carcinoma and encapsulated papillary carcinoma: a single-institution study in Japan. <i>Endocrine Journal</i> , 2020 , 67, 1147-1155	2.9	О
30	Proposed algorithm for cytological diagnosis of thyroid follicular lesions and the reporting system. <i>The Journal of the Japanese Society of Clinical Cytology</i> , 2014 , 53, 264-270	Ο	O
29	Kinetic analysis of the growth rate of sporadic and hereditary medullary thyroid carcinoma: comparing the postoperative calcitonin-doubling rate with the hypothetical preoperative tumor volume-doubling rate. <i>Thyroid Research</i> , 2020 , 13, 13	2.4	О
28	Preoperative detection of the TERT promoter mutations in papillary thyroid carcinomas. <i>Clinical Endocrinology</i> , 2021 , 95, 790-799	3.4	О
27	Exploratory Analysis to Predict Optimal Tumor Burden for Starting Lenvatinib in Patients With Radioiodine-Refractory Differentiated Thyroid Cancer. <i>Frontiers in Oncology</i> , 2021 , 11, 638123	5.3	О
26	Increased Cardiovascular Risks in Patients with TSH-Suppressed Differentiated Thyroid Cancer: A 26-Year Nationwide Swedish Cohort Study. <i>Clinical Thyroidology</i> , 2020 , 32, 178-180	0.1	O
25	Evaluation of E-Cadherin and Ecatenin Immunoreactivity for Determining Undifferentiated Cells in Anaplastic Thyroid Carcinoma. <i>Pathobiology</i> , 2021 , 88, 351-358	3.6	О
24	Proportion of serum thyroid hormone concentrations within the reference ranges in athyreotic patients on levothyroxine monotherapy: a retrospective study <i>Thyroid Research</i> , 2022 , 15, 9	2.4	O

23	Phonation Following Reconstruction of the Recurrent Laryngeal Nerve: Reply. <i>World Journal of Surgery</i> , 2017 , 41, 331-332	3.3
22	Management of Papillary Microcarcinoma of the Thyroid with A Short Column (Management of Tumors Diagnosed as Follicular Neoplasm on Cytology) 2019 , 83-90	
21	Re: "A Prospective Mixed-Methods Study of Decision-Making on Surgery or Active Surveillance for Low-Risk Papillary Thyroid Cancer" by Sawka. <i>Thyroid</i> , 2020 , 30, 1541-1542	6.2
20	Advances in the Diagnosis and Management of Papillary Thyroid Microcarcinoma 2020 , 1-10	
19	A Patient with Papillary Thyroid Carcinoma and Biochemical Evidence of Disease at Follow-Up Visits and Increasing Serum Tg Values at the Follow-Up Assessments 2016 , 135-143	
18	Comment on: "Derivation of thyroid lymphoepithelial cysts from follicular cells". <i>Endocrine Journal</i> , 2018 , 65, 877-878	2.9
17	Letter to the Editor: Reply. World Journal of Surgery, 2017, 41, 2645-2646	3.3
16	LC-1000 Flow Cytometry System Improves Risk Stratification of Thyroid Nodules with Suspected Follicular Neoplasm <i>JMA Journal</i> , 2022 , 5, 124-126	1
15	Management of Low-Risk Papillary Thyroid Carcinoma and Papillary Microcarcinoma: The Japanese Experience 2018 , 121-129	
14	A Patient with Papillary Thyroid Carcinoma and Biochemical Evidence of Disease at Follow-Up Visits and Increasing Serum Tg Values at Follow-Up Assessments 2021 , 119-129	
13	Diagnostic significance of apical membranous and cytoplasmic dot-like CD26 expression in encapsulated follicular variant of papillary thyroid carcinoma: a useful marker for capsular invasion. <i>Endocrine Journal</i> , 2020 , 67, 1207-1214	2.9
12	Can Prophylactic Central Neck Dissection for Papillary Thyroid Microcarcinoma Be Avoided to Decrease Thyroid Cancer Overtreatment?. <i>Clinical Thyroidology</i> , 2020 , 32, 477-479	0.1
11	Serum IgMs to Thyroglobulin Are Protective Against Autoimmune Thyroid Disease After Subacute Thyroiditis. <i>Clinical Thyroidology</i> , 2020 , 32, 321-323	0.1
10	Selective prophylactic lateral node dissection improves the ipsilateral lateral node recurrence-free survival: A retrospective single-center cohort study. <i>Annals of Medicine and Surgery</i> , 2020 , 57, 190-195	2
9	Neutrophil E o I lymphocyte Ratio as a Mortality Predictor in Radioactive Iodine R efractory Differentiated Thyroid Carcinoma following Lenvatinib Therapy. <i>Clinical Thyroidology</i> , 2021 , 33, 29-31	0.1
8	Papillary Carcinoma Observation 2021 , 199-203.e1	
7	Response Letter to Editorials: active surveillance of low-risk papillary microcarcinoma of the thyroid. <i>Annals of Thyroid</i> , 2018 , 3, 14-14	0.5
6	Re-Evaluation of 162 Malignant Thyroid Nodules that were Interpreted as Benign Based on Ultrasound Findings. <i>Ultrasound International Open</i> , 2018 , 4, E110-E116	2.1

5	Reply to: "Influence of specialized clinic on initial treatment choice in localized thyroid cancer". European Journal of Surgical Oncology, 2018 , 44, 2001-2002	3.6
4	Recurrent Laryngeal Nerve Paralysis [Management of Recurrent Laryngeal Nerve Injuries 2022 , 73-80	
3	Metastatic papillary thyroid carcinoma presenting with elevated serum levels of carbohydrate antigen 19-9 (CA19-9): a case report <i>Surgical Case Reports</i> , 2022 , 8, 45	0.8
2	Papillary Thyroid Carcinoma with Honeycomb-Like Growth: Clinicopathological Characteristics and Diagnostic Significance as a Novel Variant. <i>Pathobiology</i> , 2021 , 1-9	3.6
1	Melanin-producing medullary thyroid carcinoma with transformation to melanoma: A case report <i>Molecular and Clinical Oncology</i> , 2022 , 16, 34	1.6