

Akira Miyauchi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7200780/akira-miyauchi-publications-by-year.pdf>

Version: 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

364
papers

13,850
citations

60
h-index

104
g-index

374
ext. papers

16,031
ext. citations

3.5
avg, IF

6.45
L-index

#	Paper	IF	Citations
364	LC-1000 Flow Cytometry System Improves Risk Stratification of Thyroid Nodules with Suspected Follicular Neoplasm.. <i>JMA Journal</i> , 2022 , 5, 124-126	1	
363	Recurrent Laryngeal Nerve Paralysis [Management of Recurrent Laryngeal Nerve Injuries 2022 , 73-80		
362	Metastatic papillary thyroid carcinoma presenting with elevated serum levels of carbohydrate antigen 19-9 (CA19-9): a case report.. <i>Surgical Case Reports</i> , 2022 , 8, 45	0.8	
361	Melanin-producing medullary thyroid carcinoma with transformation to melanoma: A case report.. <i>Molecular and Clinical Oncology</i> , 2022 , 16, 34	1.6	
360	Proportion of serum thyroid hormone concentrations within the reference ranges in athyreotic patients on levothyroxine monotherapy: a retrospective study.. <i>Thyroid Research</i> , 2022 , 15, 9	2.4	0
359	Papillary Thyroid Carcinoma in Ukraine After Chernobyl and in Japan After Fukushima: Different Histopathological Scenarios. <i>Thyroid</i> , 2021 , 31, 1322-1334	6.2	4
358	Prevalence and Treatment Outcomes of Marine-Lenhart Syndrome in Japan.. <i>European Thyroid Journal</i> , 2021 , 10, 461-467	4.2	0
357	A Patient with Papillary Thyroid Carcinoma and Biochemical Evidence of Disease at Follow-Up Visits and Increasing Serum Tg Values at Follow-Up Assessments 2021 , 119-129		
356	53BP1 expression as a biomarker to differentiate thyroid follicular tumors. <i>Endocrine Connections</i> , 2021 , 10, 309-315	3.5	2
355	Histological alterations following fine-needle aspiration for parathyroid adenoma: Incidence and diagnostic problems. <i>Pathology International</i> , 2021 , 71, 400-405	1.8	1
354	Federated Learning for Thyroid Ultrasound Image Analysis to Protect Personal Information: Validation Study in a Real Health Care Environment. <i>JMIR Medical Informatics</i> , 2021 , 9, e25869	3.6	6
353	High Prevalence of DICER1 Mutations and Low Frequency of Gene Fusions in Pediatric Follicular-Patterned Tumors of the Thyroid. <i>Endocrine Pathology</i> , 2021 , 32, 336-346	4.2	4
352	Preoperative detection of the TERT promoter mutations in papillary thyroid carcinomas. <i>Clinical Endocrinology</i> , 2021 , 95, 790-799	3.4	0
351	Exploratory Analysis to Predict Optimal Tumor Burden for Starting Lenvatinib in Patients With Radioiodine-Refractory Differentiated Thyroid Cancer. <i>Frontiers in Oncology</i> , 2021 , 11, 638123	5.3	0
350	Marked Decrease Over Time in Conversion Surgery After Active Surveillance of Low-Risk Papillary Thyroid Microcarcinoma. <i>Thyroid</i> , 2021 , 31, 217-223	6.2	6
349	Indications and Strategy for Active Surveillance of Adult Low-Risk Papillary Thyroid Microcarcinoma: Consensus Statements from the Japan Association of Endocrine Surgery Task Force on Management for Papillary Thyroid Microcarcinoma. <i>Thyroid</i> , 2021 , 31, 183-192	6.2	48
348	Thyroid Cancer Active Surveillance Program Retention and Adherence in Japan. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2021 , 147, 77-84	3.9	7

347	Fine-needle aspiration of parathyroid adenomas: Indications as a diagnostic approach. <i>Diagnostic Cytopathology</i> , 2021 , 49, 70-76	1.4	6
346	Neutrophil-to-Lymphocyte Ratio as a Mortality Predictor in Radioactive Iodine-Refractory Differentiated Thyroid Carcinoma following Lenvatinib Therapy. <i>Clinical Thyroidology</i> , 2021 , 33, 29-31	0.1	
345	Sorafenib and Lenvatinib Treatment for Metastasis/Recurrence of Radioactive Iodine-refractory Differentiated Thyroid Carcinoma. <i>In Vivo</i> , 2021 , 35, 1057-1064	2.3	2
344	Prognostic significance of vascular invasion and cell-proliferation activity in widely invasive follicular carcinoma of the thyroid. <i>Endocrine Journal</i> , 2021 , 68, 881-888	2.9	1
343	Papillary Carcinoma Observation 2021 , 199-203.e1		
342	Clinical significance and cytological detection of tracheal puncture following thyroid fine-needle aspiration: A retrospective study. <i>Diagnostic Cytopathology</i> , 2021 , 49, 1116-1121	1.4	1
341	Poorly Differentiated Thyroid Carcinoma Coexisting with Graves Disease Involving T3 Thyrotoxicosis due to Increased D1 and D2 Activities. <i>Thyroid</i> , 2021 , 31, 1592-1596	6.2	1
340	Evaluation of E-Cadherin and E-Catenin Immunoreactivity for Determining Undifferentiated Cells in Anaplastic Thyroid Carcinoma. <i>Pathobiology</i> , 2021 , 88, 351-358	3.6	0
339	Prognostic Significance of Neutrophil-to-Lymphocyte Ratio in Differentiated Thyroid Carcinoma Having Distant Metastasis: A Comparison With Thyroglobulin-doubling Rate and Tumor Volume-doubling Rate. <i>In Vivo</i> , 2021 , 35, 1125-1132	2.3	2
338	Papillary Thyroid Carcinoma with Honeycomb-Like Growth: Clinicopathological Characteristics and Diagnostic Significance as a Novel Variant. <i>Pathobiology</i> , 2021 , 1-9	3.6	
337	Routine Chest Computed Tomography at Presentation Does Not Identify Distant Metastasis in cT1aN0 Papillary Thyroid Carcinoma. <i>Thyroid</i> , 2020 , 30, 1620-1624	6.2	7
336	Re: "A Prospective Mixed-Methods Study of Decision-Making on Surgery or Active Surveillance for Low-Risk Papillary Thyroid Cancer" by Sawka. <i>Thyroid</i> , 2020 , 30, 1541-1542	6.2	
335	Case report: exceptionally rapid growth character of hobnail variant of papillary thyroid carcinoma: a report of four cases. <i>Endocrine Journal</i> , 2020 , 67, 1047-1053	2.9	4
334	Identification of Recurrent TERT Promoter Mutations in Intrathyroid Thymic Carcinomas. <i>Endocrine Pathology</i> , 2020 , 31, 274-282	4.2	3
333	Clinicopathological features of primary thyroid Burkitt lymphoma: a systematic review and meta-analysis. <i>Diagnostic Pathology</i> , 2020 , 15, 13	3	0
332	Older Age Significantly Affects Mortality of Patients with Papillary Thyroid Carcinoma Only When They Have High-Risk Features. <i>World Journal of Surgery</i> , 2020 , 44, 1885-1891	3.3	4
331	Papillary thyroid carcinomas are highly obscured by inflammatory hypoechoic regions caused by subacute thyroiditis: a longitudinal evaluation of 710 patients using ultrasonography. <i>Endocrine Journal</i> , 2020 , 67, 569-574	2.9	4
330	Advances in the Diagnosis and Management of Papillary Thyroid Microcarcinoma 2020 , 1-10		

329	Thyroid Tubercle of Zuckerkandl May Not Arise from the Ultimobranchial Body: Results from Histological Analysis. <i>Pathobiology</i> , 2020 , 87, 193-197	3.6	1
328	Prevalence and diagnostic significance of noninvasive follicular thyroid neoplasm with papillary-like nuclear features among tumors previously diagnosed as follicular adenoma: a single-institutional study in Japan. <i>Endocrine Journal</i> , 2020 , 67, 1071-1075	2.9	7
327	The Japanese reporting system for thyroid aspiration cytology 2019 (JRSTAC2019). <i>Gland Surgery</i> , 2020 , 9, 1653-1662	2.2	2
326	Ultrasound findings of noninvasive follicular thyroid neoplasm with papillary-like nuclear features compared with those of follicular variant of papillary carcinoma and encapsulated papillary carcinoma: a single-institution study in Japan. <i>Endocrine Journal</i> , 2020 , 67, 1147-1155	2.9	0
325	Diagnostic significance of apical membranous and cytoplasmic dot-like CD26 expression in encapsulated follicular variant of papillary thyroid carcinoma: a useful marker for capsular invasion. <i>Endocrine Journal</i> , 2020 , 67, 1207-1214	2.9	
324	Is Less More? A Microsimulation Model Comparing Cost-effectiveness of the Revised American Thyroid Association® 2015 to 2009 Guidelines for the Management of Patients With Thyroid Nodules and Differentiated Thyroid Cancer. <i>Annals of Surgery</i> , 2020 , 271, 765-773	7.8	11
323	CA19-9 May Be a Prognostic Factor for Medullary Thyroid Carcinoma. <i>Clinical Thyroidology</i> , 2020 , 32, 24-26	0.1	1
322	Active surveillance of low-risk papillary thyroid microcarcinomas in Japan and other countries: a review. <i>Expert Review of Endocrinology and Metabolism</i> , 2020 , 15, 5-12	4.1	6
321	Hashimoto® Thyroiditis with Increased IgG4-Positive Plasma Cells: Using Thyroid-Specific Diagnostic Criteria May Identify Early Phase IgG4 Thyroiditis. <i>Thyroid</i> , 2020 , 30, 251-261	6.2	7
320	Subset analysis of the Japanese risk classification guidelines for papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2020 , 67, 275-282	2.9	1
319	Can Prophylactic Central Neck Dissection for Papillary Thyroid Microcarcinoma Be Avoided to Decrease Thyroid Cancer Overtreatment?. <i>Clinical Thyroidology</i> , 2020 , 32, 477-479	0.1	
318	Tumor Volume Kinetic Analyses Might Explain Excellent Prognoses in Young Patients with Papillary Thyroid Carcinoma. <i>Journal of Thyroid Research</i> , 2020 , 2020, 4652767	2.6	2
317	Serum IgMs to Thyroglobulin Are Protective Against Autoimmune Thyroid Disease After Subacute Thyroiditis. <i>Clinical Thyroidology</i> , 2020 , 32, 321-323	0.1	
316	Active surveillance of low-risk papillary thyroid microcarcinomas. <i>Gland Surgery</i> , 2020 , 9, 1663-1673	2.2	6
315	Selective prophylactic lateral node dissection improves the ipsilateral lateral node recurrence-free survival: A retrospective single-center cohort study. <i>Annals of Medicine and Surgery</i> , 2020 , 57, 190-195	2	
314	Kinetic analysis of the growth rate of sporadic and hereditary medullary thyroid carcinoma: comparing the postoperative calcitonin-doubling rate with the hypothetical preoperative tumor volume-doubling rate. <i>Thyroid Research</i> , 2020 , 13, 13	2.4	0
313	Quality of Life in Patients with Low-Risk Papillary Thyroid Microcarcinoma: Active Surveillance Versus Immediate Surgery. <i>Endocrine Practice</i> , 2020 , 26, 1451-1457	3.2	4
312	Needle Tract Implantation Following Fine-Needle Aspiration of Thyroid Cancer. <i>World Journal of Surgery</i> , 2020 , 44, 378-384	3.3	7

311	Graves Disease and mental disorders. <i>Journal of Clinical and Translational Endocrinology</i> , 2020 , 19, 1002074	0.74	9
310	Risk of malignancy and clinical outcomes of cyst fluid only nodules in the thyroid based on ultrasound and aspiration cytology. <i>Diagnostic Cytopathology</i> , 2020 , 48, 30-34	1.4	9
309	Prognostic Value of Extranodal Tumor Extension in Papillary Thyroid Carcinoma: Proposal for Upstaging of Cases with Extranodal Tumor Extension. <i>World Journal of Surgery</i> , 2020 , 44, 638-643	3.3	6
308	Subclassification of Tumor Extension and Nodal Metastasis in Papillary Thyroid Cancer to Improve Prognostic Accuracy of the Eighth Edition of the Tumor-Node-Metastasis Classification. <i>World Journal of Surgery</i> , 2020 , 44, 336-345	3.3	4
307	Association of Patient Age With Progression of Low-risk Papillary Thyroid Carcinoma Under Active Surveillance: A Systematic Review and Meta-analysis. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2020 , 146, 552-560	3.9	19
306	Increased Cardiovascular Risks in Patients with TSH-Suppressed Differentiated Thyroid Cancer: A 26-Year Nationwide Swedish Cohort Study. <i>Clinical Thyroidology</i> , 2020 , 32, 178-180	0.1	0
305	Conservative Surveillance Management of Low-Risk Papillary Thyroid Microcarcinoma. <i>Endocrinology and Metabolism Clinics of North America</i> , 2019 , 48, 215-226	5.5	23
304	Active Surveillance Versus Immediate Surgery: Questionnaire Survey on the Current Treatment Strategy for Adult Patients with Low-Risk Papillary Thyroid Microcarcinoma in Japan. <i>Thyroid</i> , 2019 , 29, 1563-1571	6.2	25
303	Flow cytometric, gene rearrangement, and karyotypic analyses of 110 cases of primary thyroid lymphoma: a single-institutional experience in Japan. <i>Endocrine Journal</i> , 2019 , 66, 1083-1091	2.9	4
302	Cribiform-Morular Variant of Papillary Thyroid Carcinoma Shows High Ki-67 Labeling Indices, despite Its Excellent Prognosis. <i>Pathobiology</i> , 2019 , 86, 248-253	3.6	7
301	Patient Experience of Thyroid Cancer Active Surveillance in Japan. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2019 , 145, 363-370	3.9	52
300	The association between thyroid hormone balance and thyroid volume in patients with Hashimoto thyroiditis. <i>Endocrine Journal</i> , 2019 , 66, 763-768	2.9	2
299	Management of Papillary Microcarcinoma of the Thyroid with A Short Column (Management of Tumors Diagnosed as Follicular Neoplasm on Cytology) 2019 , 83-90		
298	Exploratory analysis of prognostic factors for lenvatinib in radioiodine-refractory differentiated thyroid cancer. <i>Head and Neck</i> , 2019 , 41, 3023-3032	4.2	20
297	Important cytological findings for distinction between follicular variant and conventional papillary thyroid carcinoma, including noninvasive follicular thyroid tumors with papillary-like nuclear features. <i>Endocrine Journal</i> , 2019 , 66, 475-483	2.9	6
296	A Novel Diagnostic Method for Thyroid Follicular Tumors Based on Immunofluorescence Analysis of p53-Binding Protein 1 Expression: Detection of Genomic Instability. <i>Thyroid</i> , 2019 , 29, 657-665	6.2	5
295	Type 1 and type 2 iodothyronine deiodinases in the thyroid gland of patients with huge goitrous Hashimoto thyroiditis. <i>Endocrine</i> , 2019 , 64, 584-590	4	5
294	Serum Thyroid Hormone Balance in Levothyroxine Monotherapy-Treated Patients with Atrophic Thyroid After Radioiodine Treatment for Graves Disease. <i>Thyroid</i> , 2019 , 29, 1364-1370	6.2	2

293	Control of Lung Metastases and Colon Polyposis with Lenvatinib Therapy in a Patient with Cribriform-Morular Variant of Papillary Thyroid Carcinoma and an Gene Mutation: A Case Study. <i>Thyroid</i> , 2019 , 29, 1511-1517	6.2	4
292	mRNA Expression as a Novel Prognostic Marker in Papillary Thyroid Carcinomas. <i>Thyroid</i> , 2019 , 29, 1105-1114	6.1	19
291	Ultrasound image analysis using deep learning algorithm for the diagnosis of thyroid nodules. <i>Medicine (United States)</i> , 2019 , 98, e15133	1.8	33
290	Thyroid Lymphoepithelial Cysts Mimicking Calcified or Solid Nodules on Ultrasonography. <i>Ultrasound International Open</i> , 2019 , 5, E60-E64	2.1	0
289	Thyroid function related symptoms during levothyroxine monotherapy in athyreotic patients. <i>Endocrine Journal</i> , 2019 , 66, 953-960	2.9	4
288	Moderate Frequency of Anti-Thyroglobulin Antibodies in the Early Phase of Subacute Thyroiditis. <i>European Thyroid Journal</i> , 2019 , 8, 268-272	4.2	16
287	Kinetic Analysis of Growth Activity in Enlarging Papillary Thyroid Microcarcinomas. <i>Thyroid</i> , 2019 , 29, 1765-1773	6.2	18
286	CD44v8-10 and CD44s Are Age-dependently Expressed in Primary Cultured Papillary Thyroid Carcinoma Cells and Are Associated with Cell Proliferation. <i>Kobe Journal of Medical Sciences</i> , 2019 , 65, E1-E9	0.6	2
285	Acoustic Breathiness Index for the Japanese-Speaking Population: Validation Study and Exploration of Affecting Factors. <i>Journal of Speech, Language, and Hearing Research</i> , 2019 , 62, 2617-2631	2.8	4
284	Active Surveillance as First-Line Management of Papillary Microcarcinoma. <i>Annual Review of Medicine</i> , 2019 , 70, 369-379	17.4	20
283	Diagnostic clues indicating tall cell variants of papillary thyroid carcinoma in fine needle aspiration. <i>Diagnostic Cytopathology</i> , 2019 , 47, 452-457	1.4	6
282	Natural history of papillary thyroid microcarcinoma: Kinetic analyses on tumor volume during active surveillance and before presentation. <i>Surgery</i> , 2019 , 165, 25-30	3.6	40
281	Appropriateness of the revised Japanese guidelines Qisk classification for the prognosis of papillary thyroid carcinoma: a retrospective analysis of 5,845 papillary thyroid carcinoma patients. <i>Endocrine Journal</i> , 2019 , 66, 127-134	2.9	9
280	Spontaneous Deceleration and Acceleration of Growth Rate in Medullary Thyroid Carcinomas Suggested by Changes in Calcitonin Doubling Times Over Long-Term Surveillance. <i>World Journal of Surgery</i> , 2019 , 43, 504-512	3.3	5
279	Optimal needle size for thyroid fine needle aspiration cytology. <i>Endocrine Journal</i> , 2019 , 66, 143-147	2.9	8
278	The Acoustic Voice Quality Index Version 03.01 for the Japanese-speaking Population. <i>Journal of Voice</i> , 2019 , 33, 125.e1-125.e12	1.9	20
277	Excellent Prognosis of Central Lymph Node Recurrence-Free Survival for cN0M0 Papillary Thyroid Carcinoma Patients Who Underwent Routine Prophylactic Central Node Dissection. <i>World Journal of Surgery</i> , 2018 , 42, 2462-2468	3.3	7
276	Overall Survival of Papillary Thyroid Carcinoma Patients: A Single-Institution Long-Term Follow-Up of 5897 Patients. <i>World Journal of Surgery</i> , 2018 , 42, 615-622	3.3	79

275	Prognostic value of the 8 edition of the tumor-node-metastasis classification for patients with papillary thyroid carcinoma: a single-institution study at a high-volume center in Japan. <i>Endocrine Journal</i> , 2018 , 65, 707-716	2.9	5
274	Identification of Cytological Features Distinguishing Mucosa-Associated Lymphoid Tissue Lymphoma from Reactive Lymphoid Proliferation Using Thyroid Liquid-Based Cytology. <i>Acta Cytologica</i> , 2018 , 62, 93-98	3	7
273	Trends in the Implementation of Active Surveillance for Low-Risk Papillary Thyroid Microcarcinomas at Kuma Hospital: Gradual Increase and Heterogeneity in the Acceptance of This New Management Option. <i>Thyroid</i> , 2018 , 28, 488-495	6.2	48
272	Insights into the Management of Papillary Microcarcinoma of the Thyroid. <i>Thyroid</i> , 2018 , 28, 23-31	6.2	98
271	Sialic Acid-Binding Immunoglobulin-Like Lectin1 as a Novel Predictive Biomarker for Relapse in Graves Disease: A Multicenter Study. <i>Thyroid</i> , 2018 , 28, 50-59	6.2	4
270	Serum calcitonin reference values for calcium stimulation tests by electrochemiluminescence immunoassay in Japanese men with non-medullary thyroid carcinoma. <i>Surgery Today</i> , 2018 , 48, 223-228	3	6
269	Static Prognostic Factors and Appropriate Surgical Designs for Patients with Medullary Thyroid Carcinoma: The Second Report from a Single-Institution Study in Japan. <i>World Journal of Surgery</i> , 2018 , 42, 3954-3966	3.3	16
268	Re-evaluation of MIB-1 immunostaining for diagnosing hyalinizing trabecular tumour of the thyroid: semi-automated techniques with manual antigen retrieval are more accurate than fully automated techniques. <i>Endocrine Journal</i> , 2018 , 65, 239-244	2.9	8
267	Derivation of thyroid lymphoepithelial cysts from follicular cells. <i>Endocrine Journal</i> , 2018 , 65, 579-586	2.9	3
266	Spontaneous Slowing and Regressing of Tumor Growth in Childhood/Adolescent Papillary Thyroid Carcinomas Suggested by the Postoperative Thyroglobulin-Doubling Time. <i>Journal of Thyroid Research</i> , 2018 , 2018, 6470251	2.6	4
265	Characteristics and natural course of hypoechoic thyroid lesions diagnosed as possible thyroid lymphomas by fine needle aspiration cytology. <i>Thyroid Research</i> , 2018 , 11, 8	2.4	2
264	Comment on: "Derivation of thyroid lymphoepithelial cysts from follicular cells". <i>Endocrine Journal</i> , 2018 , 65, 877-878	2.9	
263	Calcifications in Thyroid Tumors on Ultrasonography: Calcification Types and Relationship with Histopathological Type. <i>Ultrasound International Open</i> , 2018 , 4, E45-E51	2.1	13
262	Thyroid sclerosing mucoepidermoid carcinoma with eosinophilia distinct from the salivary type. <i>Endocrine Journal</i> , 2018 , 65, 427-436	2.9	8
261	Management of Low-Risk Papillary Thyroid Carcinoma and Papillary Microcarcinoma: The Japanese Experience 2018 , 121-129		
260	Long-Term Follow-Up of a Patient with Sporadic Nonautoimmune Hyperthyroidism Due to a Thyrotropin-Receptor Mutation (D619G). <i>AACE Clinical Case Reports</i> , 2018 , 4, 84-89	0.7	1
259	Estimation of the lifetime probability of disease progression of papillary microcarcinoma of the thyroid during active surveillance. <i>Surgery</i> , 2018 , 163, 48-52	3.6	95
258	Utility of monoclonal PAX8 antibody for distinguishing intrathyroid thymic carcinoma from follicular cell-derived thyroid carcinoma. <i>Endocrine Journal</i> , 2018 , 65, 1171-1175	2.9	3

257	Aggressive cribriform-morular variant of papillary thyroid carcinoma: Report of an unusual case with pulmonary metastasis displaying poorly differentiated features. <i>Pathology International</i> , 2018 , 68, 700-705	1.8	5
256	Thyroid Fine-Needle Aspiration and Smearing Techniques. <i>VideoEndocrinology</i> , 2018 , 5,	1.6	13
255	Response Letter to Editorials: active surveillance of low-risk papillary microcarcinoma of the thyroid. <i>Annals of Thyroid</i> , 2018 , 3, 14-14	0.5	
254	Thrombotic Microangiopathy with Severe Proteinuria Induced by Lenvatinib for Radioactive Iodine-Refractory Papillary Thyroid Carcinoma. <i>Case Reports in Oncology</i> , 2018 , 11, 735-741	1	9
253	International neuromonitoring study group guidelines 2018: Part II: Optimal recurrent laryngeal nerve management for invasive thyroid cancer-incorporation of surgical, laryngeal, and neural electrophysiologic data. <i>Laryngoscope</i> , 2018 , 128 Suppl 3, S18-S27	3.6	74
252	Impact of the modification of the diagnostic criteria in the 2017 Bethesda System for Reporting Thyroid Cytopathology: a report of a single institution in Japan. <i>Endocrine Journal</i> , 2018 , 65, 1193-1198	2.9	10
251	Phantom Nodules Detected by Ultrasound Examination of the Neck: The Possibility of Ectopic Cervical Thymic Tissue in Adults. <i>Ultrasound International Open</i> , 2018 , 4, E119-E123	2.1	1
250	Sonographic Evaluation of Nodules Newly Detected in the Neck After Thyroidectomy: Suture Granuloma Versus Recurrent Carcinoma. <i>Ultrasound International Open</i> , 2018 , 4, E124-E130	2.1	3
249	Re-Evaluation of 162 Malignant Thyroid Nodules that were Interpreted as Benign Based on Ultrasound Findings. <i>Ultrasound International Open</i> , 2018 , 4, E110-E116	2.1	
248	Reply to: "Influence of specialized clinic on initial treatment choice in localized thyroid cancer". <i>European Journal of Surgical Oncology</i> , 2018 , 44, 2001-2002	3.6	
247	Calcitonin measurement in fine-needle aspirate washout fluid by electrochemiluminescence immunoassay for thyroid tumors. <i>Thyroid Research</i> , 2018 , 11, 15	2.4	6
246	Immunohistochemical and Molecular Analyses Focusing on Mesenchymal Cells in Papillary Thyroid Carcinoma with Desmoid-Type Fibromatosis. <i>Pathobiology</i> , 2018 , 85, 300-303	3.6	9
245	Can Ultrasound Alone Predict Papillary Thyroid Carcinoma with Desmoid-Type Fibromatosis? A Retrospective Analysis of 13 Cases, Focusing on the Stromal Area. <i>Ultrasound International Open</i> , 2018 , 4, E39-E44	2.1	3
244	Prognostic value of the 8 tumor-node-metastasis classification for follicular carcinoma and poorly differentiated carcinoma of the thyroid in Japan. <i>Endocrine Journal</i> , 2018 , 65, 621-627	2.9	6
243	Successful treatment switch from lenvatinib to sorafenib in a patient with radioactive iodine-refractory differentiated thyroid cancer intolerant to lenvatinib due to severe proteinuria. <i>Auris Nasus Larynx</i> , 2018 , 45, 1249-1252	2.2	6
242	Warthin-like papillary thyroid carcinoma with immunoglobulin G4-positive plasma cells possibly related to Hashimoto's thyroiditis. <i>Endocrine Journal</i> , 2018 , 65, 175-180	2.9	3
241	Validation of the Acoustic Voice Quality Index in the Japanese Language. <i>Journal of Voice</i> , 2017 , 31, 260.e1-260.e9	2.9	9
240	Biochemical Markers Reflecting Thyroid Function in Athyreotic Patients on Levothyroxine Monotherapy. <i>Thyroid</i> , 2017 , 27, 484-490	6.2	46

239	TERT promoter mutations and Ki-67 labeling index as a prognostic marker of papillary thyroid carcinomas: combination of two independent factors. <i>Scientific Reports</i> , 2017 , 7, 41752	4.9	32
238	Minute Hyalinizing Trabecular Tumor Misinterpreted as a Blood Vessel on Ultrasonography: A Case Report. <i>Ultrasound International Open</i> , 2017 , 3, E42-E44	2.1	1
237	Frequent BRAF and Absence of TERT Promoter Mutations Characterize Sporadic Pediatric Papillary Thyroid Carcinomas in Japan. <i>Endocrine Pathology</i> , 2017 , 28, 103-111	4.2	26
236	The Effectiveness of Prophylactic Modified Neck Dissection for Reducing the Development of Lymph Node Recurrence of Papillary Thyroid Carcinoma. <i>World Journal of Surgery</i> , 2017 , 41, 2283-2289	3.3	25
235	"Nodule in Nodule" on Thyroid Ultrasonography: Possibility of Follicular Carcinoma Transformed from Benign Thyroid Tumor. <i>European Thyroid Journal</i> , 2017 , 6, 101-107	4.2	6
234	Surgical Methods and Experiences of Surgeons did not Significantly Affect the Recovery in Phonation Following Reconstruction of the Recurrent Laryngeal Nerve: Reply. <i>World Journal of Surgery</i> , 2017 , 41, 331-332	3.3	
233	Paediatric follicular thyroid carcinoma - indolent cancer with low prevalence of RAS mutations and absence of PAX8-PPARG fusion in a Japanese population. <i>Histopathology</i> , 2017 , 71, 760-768	7.3	15
232	TERT Promoter Mutations Were Not Found in Papillary Thyroid Microcarcinomas That Showed Disease Progression on Active Surveillance. <i>Thyroid</i> , 2017 , 27, 1206-1207	6.2	29
231	Role of immediate recurrent laryngeal nerve reconstruction in surgery for thyroid cancers with fixed vocal cords. <i>Head and Neck</i> , 2017 , 39, 427-431	4.2	6
230	Cytoplasmic Lipid Accumulation Characteristic of the Cribriform Variant of Papillary Thyroid Carcinoma. <i>Pathobiology</i> , 2017 , 84, 251-257	3.6	5
229	Comparative histopathological analysis of sporadic pediatric papillary thyroid carcinoma from Japan and Ukraine. <i>Endocrine Journal</i> , 2017 , 64, 977-993	2.9	9
228	Papillary thyroid carcinoma with desmoid-type fibromatosis: A clinical, pathological, and immunohistochemical study of 14 cases. <i>Endocrine Journal</i> , 2017 , 64, 1017-1023	2.9	15
227	The thyroid cancer policy model: A mathematical simulation model of papillary thyroid carcinoma in The U.S. population. <i>PLoS ONE</i> , 2017 , 12, e0177068	3.7	4
226	Comparative frequency of four different types of pregnancy-associated thyrotoxicosis in a single thyroid centre. <i>Thyroid Research</i> , 2017 , 10, 4	2.4	7
225	Fine-needle aspiration cytology for medullary thyroid carcinoma: a single institutional experience in Japan. <i>Endocrine Journal</i> , 2017 , 64, 1099-1104	2.9	19
224	Noninvasive follicular thyroid neoplasm with papillary-like nuclear features: a single-institutional experience in Japan. <i>Endocrine Journal</i> , 2017 , 64, 1149-1155	2.9	32
223	Preoperative diagnostic algorithm of primary thyroid lymphoma using ultrasound, aspiration cytology, and flow cytometry. <i>Endocrine Journal</i> , 2017 , 64, 859-865	2.9	24
222	Comparison of thyroglobulin and thyroid peroxidase antibodies measured by five different kits in autoimmune thyroid diseases. <i>Endocrine Journal</i> , 2017 , 64, 955-961	2.9	8

221	Reappraisal of "cyst fluid only" on thyroid fine-needle aspiration cytology. <i>Endocrine Journal</i> , 2017 , 64, 759-765	2.9	15
220	Letter to the Editor: Reply. <i>World Journal of Surgery</i> , 2017 , 41, 2645-2646	3.3	
219	Genotype Analyses in the Japanese and Belarusian Populations Reveal Independent Effects of rs965513 and rs1867277 but Do Not Support the Role of FOXE1 Polyalanine Tract Length in Conferring Risk for Papillary Thyroid Carcinoma. <i>Thyroid</i> , 2017 , 27, 224-235	6.2	13
218	Prognostic Significance of the Proportion of Tall Cell Components in Papillary Thyroid Carcinoma. <i>World Journal of Surgery</i> , 2017 , 41, 742-747	3.3	16
217	Comparison of the costs of active surveillance and immediate surgery in the management of low-risk papillary microcarcinoma of the thyroid. <i>Endocrine Journal</i> , 2017 , 64, 59-64	2.9	77
216	Chromophobe renal cell carcinoma-like thyroid carcinoma: A novel clinicopathologic entity possibly associated with tuberous sclerosis complex. <i>Endocrine Journal</i> , 2017 , 64, 843-850	2.9	8
215	Active Surveillance as the Initial Course of Action in Low-Risk Papillary Microcarcinoma 2017 , 135-141		1
214	Incidences of Unfavorable Events in the Management of Low-Risk Papillary Microcarcinoma of the Thyroid by Active Surveillance Versus Immediate Surgery. <i>Thyroid</i> , 2016 , 26, 150-5	6.2	200
213	Revisiting Low-Risk Thyroid Papillary Microcarcinomas Resected Without Observation: Was Immediate Surgery Necessary?. <i>World Journal of Surgery</i> , 2016 , 40, 523-8	3.3	44
212	Diagnosis and surgical indications of oxyphilic follicular tumors in Japan: Surgical specimens and cytology. <i>Endocrine Journal</i> , 2016 , 63, 977-982	2.9	10
211	Kinetic analyses of changes in serum TSH receptor antibody values after total thyroidectomy in patients with Graves Disease. <i>Endocrine Journal</i> , 2016 , 63, 179-85	2.9	7
210	Insights and clinical questions about the active surveillance of low-risk papillary thyroid microcarcinomas [Review]. <i>Endocrine Journal</i> , 2016 , 63, 323-8	2.9	30
209	Diagnostic value of GATA-3 in cytological identification of parathyroid tissues. <i>Endocrine Journal</i> , 2016 , 63, 621-6	2.9	13
208	Pathological characteristics of low-risk papillary thyroid microcarcinoma with progression during active surveillance. <i>Endocrine Journal</i> , 2016 , 63, 805-810	2.9	40
207	Reference values of serum calcitonin with calcium stimulation tests by electrochemiluminescence immunoassay before/after total thyroidectomy in Japanese patients with thyroid diseases other than medullary thyroid carcinoma. <i>Endocrine Journal</i> , 2016 , 63, 627-32	2.9	13
206	Calcitonin doubling time in medullary thyroid carcinoma after the detection of distant metastases keenly predicts patients' carcinoma death. <i>Endocrine Journal</i> , 2016 , 63, 663-7	2.9	12
205	Prognostic impact of Ki-67 labeling index in minimally invasive follicular thyroid carcinoma. <i>Endocrine Journal</i> , 2016 , 63, 913-917	2.9	7
204	Partial prediction of postpartum Graves' thyrotoxicosis by sensitive bioassay for thyroid-stimulating antibody measured in early pregnancy. <i>Endocrine Journal</i> , 2016 , 63, 929-932	2.9	3

203	Age- and Gender-Specific Risk of Thyroid Cancer in Patients With Familial Adenomatous Polyposis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, 4611-4617	5.6	27
202	Surgical Methods and Experiences of Surgeons did not Significantly Affect the Recovery in Phonation Following Reconstruction of the Recurrent Laryngeal Nerve. <i>World Journal of Surgery</i> , 2016 , 40, 2948-2955	3.3	13
201	Malignant melanoma arising in melanin-producing medullary thyroid carcinoma. <i>International Journal of Surgery Case Reports</i> , 2016 , 20, 118-22	0.8	4
200	Innervation of the cricothyroid muscle by the recurrent laryngeal nerve. <i>Head and Neck</i> , 2016 , 38 Suppl 1, E441-5	4.2	17
199	Clinical Trials of Active Surveillance of Papillary Microcarcinoma of the Thyroid. <i>World Journal of Surgery</i> , 2016 , 40, 516-22	3.3	148
198	Effects of Pregnancy on Papillary Microcarcinomas of the Thyroid Re-Evaluated in the Entire Patient Series at Kuma Hospital. <i>Thyroid</i> , 2016 , 26, 156-60	6.2	46
197	Immunohistochemical detection of NRAS(Q61R) protein in follicular-patterned thyroid tumors. <i>Human Pathology</i> , 2016 , 53, 51-7	3.7	15
196	A Patient with Papillary Thyroid Carcinoma and Biochemical Evidence of Disease at Follow-Up Visits and Increasing Serum Tg Values at the Follow-Up Assessments 2016 , 135-143		
195	2015 American Thyroid Association Guidelines for Thyroid Nodules and Differentiated Thyroid Cancer and Their Implementation in Various Care Settings. <i>Thyroid</i> , 2016 , 26, 319-21	6.2	25
194	A Clinical Framework to Facilitate Risk Stratification When Considering an Active Surveillance Alternative to Immediate Biopsy and Surgery in Papillary Microcarcinoma. <i>Thyroid</i> , 2016 , 26, 144-9	6.2	185
193	Tumor protrusion with intensive blood signals on ultrasonography is a strongly suggestive finding of follicular thyroid carcinoma. <i>Medical Ultrasonography</i> , 2016 , 18, 25-9	1.4	15
192	Intraoperative Neural Injury Management: Transection and Segmental Defects 2016 , 253-258		2
191	A Novel Germline Mutation of KEAP1 (R483H) Associated with a Non-Toxic Multinodular Goiter. <i>Frontiers in Endocrinology</i> , 2016 , 7, 131	5.7	10
190	Prospective study of vocal fold function after loss of the neuromonitoring signal in thyroid surgery: The International Neural Monitoring Study Group@POLT study. <i>Laryngoscope</i> , 2016 , 126, 1260-6	3.6	69
189	Differentiating between benign follicular nodules and follicular neoplasms in thyroid liquid-based cytology preparations. <i>Diagnostic Cytopathology</i> , 2016 , 44, 659-64	1.4	4
188	Impact of positional changes in neural monitoring endotracheal tube on amplitude and latency of electromyographic response in monitored thyroid surgery: Results from the Porcine Experiment. <i>Head and Neck</i> , 2016 , 38 Suppl 1, E1004-8	4.2	41
187	Periodic Granulocyte Count Measuring Is Useful for Detecting Asymptomatic Agranulocytosis in Antithyroid Drug-Treated Patients with Graves Disease. <i>European Thyroid Journal</i> , 2016 , 5, 253-260	4.2	10
186	Germline RET mutation carriers in Japanese patients with apparently sporadic medullary thyroid carcinoma: A single institution experience. <i>Auris Nasus Larynx</i> , 2016 , 43, 551-5	2.2	9

185	The Safety and Efficacy of Weekly Paclitaxel Administration for Anaplastic Thyroid Cancer Patients: A Nationwide Prospective Study. <i>Thyroid</i> , 2016 , 26, 1293-9	6.2	33
184	Right-sided aortic arch and aberrant left subclavian artery with or without a left nonrecurrent inferior laryngeal nerve. <i>Head and Neck</i> , 2016 , 38, E2508-11	4.2	13
183	Innervation of the cricothyroid muscle by extralaryngeal branches of the recurrent laryngeal nerve. <i>Laryngoscope</i> , 2016 , 126, 1157-62	3.6	14
182	Prospective randomized study on injury of the external branch of the superior laryngeal nerve during thyroidectomy comparing intraoperative nerve monitoring and a conventional technique. <i>Head and Neck</i> , 2015 , 37, 1456-60	4.2	30
181	Effect of the presence of remnant thyroid tissue on the serum thyroid hormone balance in thyroidectomized patients. <i>European Journal of Endocrinology</i> , 2015 , 173, 333-40	6.5	25
180	Metastatic carcinoma to the thyroid gland from renal cell carcinoma: role of ultrasonography in preoperative diagnosis. <i>Thyroid Research</i> , 2015 , 8, 4	2.4	21
179	Is an Increase in Thyroid Nodule Volume a Risk Factor for Malignancy?. <i>Thyroid</i> , 2015 , 25, 804-11	6.2	27
178	Cytological findings for the diagnosis of primary thyroid mucosa-associated lymphoid tissue lymphoma by fine needle aspiration. <i>Acta Cytologica</i> , 2015 , 59, 26-36	3	13
177	Nonoperative management of low-risk differentiated thyroid carcinoma. <i>Current Opinion in Oncology</i> , 2015 , 27, 15-20	4.2	56
176	Concept and design of a nationwide prospective feasibility/efficacy/safety study of weekly paclitaxel for patients with pathologically confirmed anaplastic thyroid cancer (ATCCJ-PTX-P2). <i>BMC Cancer</i> , 2015 , 15, 475	4.8	9
175	Characteristic sonographic features of cribriform papillary thyroid carcinoma for differentiation from other thyroid nodules. <i>Journal of Medical Ultrasonics (2001)</i> , 2015 , 42, 83-7	1.4	11
174	Thyroxine treatment may be useful for subclinical hypothyroidism in patients with female infertility. <i>Endocrine Journal</i> , 2015 , 62, 87-92	2.9	23
173	Diagnostic significance of PAX8 in thyroid squamous cell carcinoma. <i>Endocrine Journal</i> , 2015 , 62, 991-5	2.9	35
172	Occurrence of thyroxine tablet (Thyradin S(□)) - induced liver dysfunction in a patient with subclinical hypothyroidism. <i>Endocrine Journal</i> , 2015 , 62, 719-24	2.9	4
171	Graves Disease Patients with Persistent Hyperthyroidism and Diffuse Lymphoplasmacytic Infiltration in the Thyroid Show No Histopathological Compatibility with IgG4-Related Disease. <i>PLoS ONE</i> , 2015 , 10, e0134143	3.7	11
170	The common genetic variant rs944289 on chromosome 14q13.3 associates with risk of both malignant and benign thyroid tumors in the Japanese population. <i>Thyroid</i> , 2015 , 25, 333-40	6.2	33
169	Cytological characteristics of papillary thyroid carcinoma on LBC specimens, compared with conventional specimens. <i>Diagnostic Cytopathology</i> , 2015 , 43, 108-13	1.4	23
168	Rapid bioassay for detection of thyroid-stimulating antibodies using cyclic adenosine monophosphate-gated calcium channel and aequorin. <i>European Thyroid Journal</i> , 2015 , 4, 14-9	4.2	23

167	Airtight tracheocutaneostomy after window resection of the trachea for invasive papillary thyroid carcinoma: experience of 109 cases. <i>World Journal of Surgery</i> , 2014 , 38, 660-6	3.3	9
166	BRAF(V600E) mutation analysis in papillary thyroid carcinoma: is it useful for all patients?. <i>World Journal of Surgery</i> , 2014 , 38, 679-87	3.3	43
165	Patient age is significantly related to the progression of papillary microcarcinoma of the thyroid under observation. <i>Thyroid</i> , 2014 , 24, 27-34	6.2	472
164	Up-regulation of urinary-type plasminogen activator correlates with high-risk papillary thyroid carcinoma with BRAF(V600E) mutation and its possible molecular mechanism. <i>Pathology Research and Practice</i> , 2014 , 210, 733-8	3.4	2
163	Papillary thyroid microcarcinoma might progress during pregnancy. <i>Thyroid</i> , 2014 , 24, 840-4	6.2	58
162	Differentiation of postpartum Graves thyrotoxicosis from postpartum destructive thyrotoxicosis using antithyrotropin receptor antibodies and thyroid blood flow. <i>Thyroid</i> , 2014 , 24, 1027-31	6.2	47
161	Anaplastic thyroid carcinomas incidentally found on postoperative pathological examination. <i>World Journal of Surgery</i> , 2014 , 38, 2311-6	3.3	16
160	Outcome of vocal cord function after partial layer resection of the recurrent laryngeal nerve in patients with invasive papillary thyroid cancer. <i>Surgery</i> , 2014 , 155, 184-9	3.6	31
159	Introducing the reporting system for thyroid fine-needle aspiration cytology according to the new guidelines of the Japan Thyroid Association. <i>Endocrine Journal</i> , 2014 , 61, 539-52	2.9	38
158	Prognostic significance of young age in papillary thyroid carcinoma: analysis of 5,733 patients with 150 months median follow-up. <i>Endocrine Journal</i> , 2014 , 61, 491-7	2.9	17
157	Permanent hypoparathyroidism after completion total thyroidectomy as a second surgery: How do we avoid it?. <i>Endocrine Journal</i> , 2014 , 61, 403-8	2.9	24
156	Prognostic significance of patient age in minimally and widely invasive follicular thyroid carcinoma: investigation of three age groups. <i>Endocrine Journal</i> , 2014 , 61, 265-71	2.9	21
155	Prognostic values of clinical lymph node metastasis and macroscopic extrathyroid extension in papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2014 , 61, 745-50	2.9	29
154	Prognosis and prognostic factors of differentiated thyroid carcinoma after the appearance of metastasis refractory to radioactive iodine therapy. <i>Endocrine Journal</i> , 2014 , 61, 821-4	2.9	20
153	Changes in serum thyroglobulin antibody levels as a dynamic prognostic factor for early-phase recurrence of thyroglobulin antibody-positive papillary thyroid carcinoma after total thyroidectomy. <i>Endocrine Journal</i> , 2014 , 61, 961-5	2.9	20
152	Prognosis and growth activity depend on patient age in clinical and subclinical papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2014 , 61, 205-13	2.9	32
151	Static and dynamic prognostic factors of papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2014 , 61, 1145-51	2.9	18
150	Tandem Germline RET Mutations in a Family Pathogenetic for Multiple Endocrine Neoplasia 2B, Confirmed by a Natural Experiment. <i>European Thyroid Journal</i> , 2014 , 3, 272-7	4.2	2

149	Prevalence of thyrotropin receptor germline mutations and clinical courses in 89 hyperthyroid patients with diffuse goiter and negative anti-thyrotropin receptor antibodies. <i>Thyroid</i> , 2014 , 24, 789-95	6.2	10
148	Cytologic findings and differential diagnoses of primary thyroid MALT lymphoma with striking plasma cell differentiation and amyloid deposition. <i>Diagnostic Cytopathology</i> , 2014 , 42, 73-7	1.4	12
147	Evaluation of "Benign" in Thyroid Bethesda System. <i>The Journal of the Japanese Society of Clinical Cytology</i> , 2014 , 53, 257-263	0	1
146	Proposed algorithm for cytological diagnosis of thyroid follicular lesions and the reporting system. <i>The Journal of the Japanese Society of Clinical Cytology</i> , 2014 , 53, 264-270	0	0
145	Functional characterization of the novel BRAF complex mutation, BRAF(V600delinsYM) , identified in papillary thyroid carcinoma. <i>International Journal of Cancer</i> , 2013 , 132, 738-43	7.5	16
144	External branch of the superior laryngeal nerve monitoring during thyroid and parathyroid surgery: International Neural Monitoring Study Group standards guideline statement. <i>Laryngoscope</i> , 2013 , 123 Suppl 4, S1-14	3.6	218
143	Papillary thyroid carcinoma with honeycomb-like multiple small cysts: characteristic features on ultrasonography. <i>European Thyroid Journal</i> , 2013 , 2, 270-4	4.2	7
142	Evaluation of cytologically benign solitary thyroid nodules by ultrasonography: a retrospective analysis of 1877 cases. <i>Auris Nasus Larynx</i> , 2013 , 40, 308-11	2.2	13
141	Attenuation of postmenopausal bone loss in patients with transient hypoparathyroidism after total thyroidectomy. <i>World Journal of Surgery</i> , 2013 , 37, 2860-5	3.3	17
140	Voice quality after surgical treatment for thyroid cancer. <i>Thyroid</i> , 2013 , 23, 847-53	6.2	35
139	Ki-67 labeling index is a predictor of postoperative persistent disease and cancer growth and a prognostic indicator in papillary thyroid carcinoma. <i>European Thyroid Journal</i> , 2013 , 2, 57-64	4.2	26
138	Analysis of 754 cases of antithyroid drug-induced agranulocytosis over 30 years in Japan. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 4776-83	5.6	95
137	Intrathyroidal epithelial thymoma/carcinoma showing thymus-like differentiation; comparison with thymic lymphoepithelioma-like carcinoma and a possibility of development from a multipotential stem cell. <i>Apmis</i> , 2013 , 121, 523-30	3.4	18
136	Immunoglobulin G4 thyroiditis in a Graves disease patient with a large goiter developing hypothyroidism. <i>Thyroid</i> , 2013 , 23, 1496-7	6.2	16
135	Initial treatment with 15 mg of prednisolone daily is sufficient for most patients with subacute thyroiditis in Japan. <i>Thyroid</i> , 2013 , 23, 269-72	6.2	35
134	Prognostic significance of changes in serum thyroglobulin antibody levels of pre- and post-total thyroidectomy in thyroglobulin antibody-positive papillary thyroid carcinoma patients. <i>Endocrine Journal</i> , 2013 , 60, 871-6	2.9	39
133	Prognostic factors of minimally invasive follicular thyroid carcinoma: extensive vascular invasion significantly affects patient prognosis. <i>Endocrine Journal</i> , 2013 , 60, 637-42	2.9	69
132	Distant metastasis at diagnosis and large tumor size are significant prognostic factors of widely invasive follicular thyroid carcinoma. <i>Endocrine Journal</i> , 2013 , 60, 829-33	2.9	23

131	Relationship of biochemically persistent disease and thyroglobulin-doubling time to age at surgery in patients with papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2013 , 60, 415-421	2.9	34
130	PSYCHOLOGICAL CHARACTERISTICS OF THE NEO-FFI AND THE TREE DRAWING TEST IN PATIENTS WITH THYROID DISEASE. <i>Psychologia</i> , 2013 , 56, 138-153	0.2	2
129	Tumor size is the strongest predictor of microscopic lymph node metastasis and lymph node recurrence of N0 papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2013 , 60, 113-7	2.9	51
128	Intrathyroid epithelial thymoma (ITET) and carcinoma showing thymus-like differentiation (CASTLE): CD5-positive neoplasms mimicking squamous cell carcinoma of the thyroid. <i>Histology and Histopathology</i> , 2013 , 28, 543-56	1.4	27
127	Cytological findings of primary thyroid MALT lymphoma with extreme plasma cell differentiation in fine needle aspiration specimens. <i>The Journal of the Japanese Society of Clinical Cytology</i> , 2013 , 52, 507-517	0.1	2
126	Findings and significance of mountain range-like clusters derived from follicular colonization for the diagnosis of MALT lymphoma. <i>The Journal of the Japanese Society of Clinical Cytology</i> , 2013 , 52, 428-436	0.3	2
125	Relationship of biochemically persistent disease and thyroglobulin-doubling time to age at surgery in patients with papillary thyroid carcinoma. <i>Endocrine Journal</i> , 2013 , 60, 415-21	2.9	25
124	Laryngeal approach to the recurrent laryngeal nerve involved by thyroid cancer at the ligament of Berry. <i>Surgery</i> , 2012 , 152, 57-60	3.6	12
123	Prognosis of low-risk papillary thyroid carcinoma patients: its relationship with the size of primary tumors. <i>Endocrine Journal</i> , 2012 , 59, 119-25	2.9	43
122	IgG4-related disease of the thyroid glands. <i>Endocrine Journal</i> , 2012 , 59, 273-81	2.9	70
121	Prognosis and prognostic factors of patients with papillary thyroid carcinoma requiring resection of recurrent laryngeal nerve due to carcinoma extension. <i>Endocrine Journal</i> , 2012 , 59, 247-52	2.9	9
120	Investigation of the prognosis of patients with papillary thyroid carcinoma by tumor size. <i>Endocrine Journal</i> , 2012 , 59, 457-64	2.9	28
119	Relationship between prognosis of papillary thyroid carcinoma patient and age: a retrospective single-institution study. <i>Endocrine Journal</i> , 2012 , 59, 399-405	2.9	40
118	Prognostic factors for recurrence of papillary thyroid carcinoma in the lymph nodes, lung, and bone: analysis of 5,768 patients with average 10-year follow-up. <i>World Journal of Surgery</i> , 2012 , 36, 1274-8	3.3	162
117	Prognostic factors and treatment outcomes for anaplastic thyroid carcinoma: ATC Research Consortium of Japan cohort study of 677 patients. <i>World Journal of Surgery</i> , 2012 , 36, 1247-54	3.3	170
116	TSH-suppressive doses of levothyroxine are required to achieve preoperative native serum triiodothyronine levels in patients who have undergone total thyroidectomy. <i>European Journal of Endocrinology</i> , 2012 , 167, 373-8	6.5	80
115	Distinct histopathological features of Hashimoto's thyroiditis with respect to IgG4-related disease. <i>Modern Pathology</i> , 2012 , 25, 1086-97	9.8	71
114	Histopathological diagnoses of "accessory" thyroid nodules diagnosed as benign by fine-needle aspiration cytology and ultrasonography. <i>Thyroid</i> , 2012 , 22, 299-303	6.2	5

113	Undetectable serum thyroglobulin levels in patients with medullary thyroid carcinoma after total thyroidectomy without radioiodine ablation. <i>Thyroid</i> , 2012 , 22, 680-2	6.2	11
112	Clinical significance and prognostic impact of subcutaneous or intrasternal muscular recurrence of papillary thyroid carcinoma. <i>Journal of Thyroid Research</i> , 2012 , 2012, 819797	2.6	5
111	Prognosis and prognostic factors of papillary thyroid carcinoma in patients under 20 years. <i>Endocrine Journal</i> , 2012 , 59, 539-45	2.9	34
110	The FOXE1 and NKX2-1 loci are associated with susceptibility to papillary thyroid carcinoma in the Japanese population. <i>Journal of Medical Genetics</i> , 2011 , 48, 645-8	5.8	68
109	Prognostic impact of serum thyroglobulin doubling-time under thyrotropin suppression in patients with papillary thyroid carcinoma who underwent total thyroidectomy. <i>Thyroid</i> , 2011 , 21, 707-16	6.2	183
108	Role of ultrasonography in patients with cytologically follicular thyroid tumor. <i>Auris Nasus Larynx</i> , 2011 , 38, 508-11	2.2	16
107	Serum calcitonin levels with calcium loading tests before and after total thyroidectomy in patients with thyroid diseases other than medullary thyroid carcinoma. <i>Endocrine Journal</i> , 2011 , 58, 217-21	2.9	19
106	Our experience of treatment of cribriform morular variant of papillary thyroid carcinoma; difference in clinicopathological features of FAP-associated and sporadic patients. <i>Endocrine Journal</i> , 2011 , 58, 685-9	2.9	50
105	Prognostic significance of carcinoma extension from primary lesions and metastatic nodes in papillary thyroid carcinoma: appropriateness of three subdivisions of extension. <i>Endocrine Journal</i> , 2011 , 58, 961-7	2.9	7
104	Electrophysiologic recurrent laryngeal nerve monitoring during thyroid and parathyroid surgery: international standards guideline statement. <i>Laryngoscope</i> , 2011 , 121 Suppl 1, S1-16	3.6	636
103	Significance of prophylactic modified radical neck dissection for patients with low-risk papillary thyroid carcinoma measuring 1.1-3.0 cm: first report of a trial at Kuma Hospital. <i>Surgery Today</i> , 2011 , 41, 1486-91	3	19
102	Final pathology findings after immediate or delayed surgery in patients with cytologically benign or follicular thyroid nodules. <i>World Journal of Surgery</i> , 2011 , 35, 558-62	3.3	13
101	Prognosis of patients with papillary thyroid carcinoma showing postoperative recurrence to the central neck. <i>World Journal of Surgery</i> , 2011 , 35, 767-72	3.3	39
100	Prognostic factors of papillary thyroid carcinoma vary according to sex and patient age. <i>World Journal of Surgery</i> , 2011 , 35, 2684-90	3.3	18
99	The thyroid function of Graves disease patients is aggravated by depressive personality during antithyroid drug treatment. <i>BioPsychoSocial Medicine</i> , 2011 , 5, 9	2.8	10
98	Loss of cellular polarity/cohesiveness in the invasive front of papillary thyroid carcinoma, a novel predictor for lymph node metastasis; possible morphological indicator of epithelial mesenchymal transition. <i>Journal of Clinical Pathology</i> , 2011 , 64, 325-9	3.9	36
97	Tumor thrombus of thyroid malignancies in veins: importance of detection by ultrasonography. <i>Thyroid</i> , 2011 , 21, 527-31	6.2	26
96	Type 1 and type 2 iodothyronine deiodinases in the thyroid gland of patients with 3,5,3,5-tetraiodothyronine-predominant Graves disease. <i>European Journal of Endocrinology</i> , 2011 , 164, 95-100	6.5	30

95	Induction chemotherapy with weekly paclitaxel administration for anaplastic thyroid carcinoma. <i>Thyroid</i> , 2010 , 20, 7-14	6.2	105
94	Subclinical nonautoimmune hyperthyroidism in a family segregates with a thyrotropin receptor mutation with weakly increased constitutive activity. <i>Thyroid</i> , 2010 , 20, 1307-14	6.2	9
93	Distinct clinical, serological, and sonographic characteristics of hashimoto's thyroiditis based with and without IgG4-positive plasma cells. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, 1309-17	5.6	124
92	Distribution of IgG4- and/or IgG-positive plasma cells in Hashimoto's thyroiditis: an immunohistochemical study. <i>Pathobiology</i> , 2010 , 77, 267-72	3.6	23
91	Prognosis and prognostic factors of patients with papillary carcinoma showing distant metastasis at surgery (M1 patients) in Japan. <i>Endocrine Journal</i> , 2010 , 57, 523-31	2.9	35
90	Thyroid ultrasonography. <i>World Journal of Surgery</i> , 2010 , 34, 1171-80	3.3	19
89	An observational trial for papillary thyroid microcarcinoma in Japanese patients. <i>World Journal of Surgery</i> , 2010 , 34, 28-35	3.3	523
88	Excellent prognosis of patients with solitary T1N0M0 papillary thyroid carcinoma who underwent thyroidectomy and elective lymph node dissection without radioiodine therapy. <i>World Journal of Surgery</i> , 2010 , 34, 1285-90	3.3	86
87	Establishment of an intraoperative staging system (iStage) by improving UICC TNM classification system for papillary thyroid carcinoma. <i>World Journal of Surgery</i> , 2010 , 34, 2570-80	3.3	58
86	Clinical outcomes of patients with papillary thyroid carcinoma after the detection of distant recurrence. <i>World Journal of Surgery</i> , 2010 , 34, 2333-7	3.3	29
85	Prognostic significance of ki-67 labeling index in papillary thyroid carcinoma. <i>World Journal of Surgery</i> , 2010 , 34, 3015-21	3.3	50
84	Prognostic impact of extrathyroid extension and clinical lymph node metastasis in papillary thyroid carcinoma depend on carcinoma size. <i>World Journal of Surgery</i> , 2010 , 34, 3007-14	3.3	32
83	Cribiform-morular variant of papillary thyroid carcinoma--cytological and immunocytochemical findings of 18 cases. <i>Diagnostic Cytopathology</i> , 2010 , 38, 890-6	1.4	44
82	Thyroid function in early pregnancy in Japanese healthy women: relation to urinary iodine excretion, emesis, and fetal and child development. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 1683-8	5.6	66
81	Biological behavior and prognosis of familial papillary thyroid carcinoma. <i>Surgery</i> , 2009 , 145, 100-5	3.6	80
80	Improvement in phonation after reconstruction of the recurrent laryngeal nerve in patients with thyroid cancer invading the nerve. <i>Surgery</i> , 2009 , 146, 1056-62	3.6	84
79	Local prognosis of patients with papillary thyroid carcinoma who were intra-operatively diagnosed as having minimal invasion of the trachea: a 17-year experience in a single institute. <i>Asian Journal of Surgery</i> , 2009 , 32, 102-8	1.6	18
78	Cytologic findings of primary thyroid MALT lymphoma with extreme plasma cell differentiation: FNA cytology of two cases. <i>Diagnostic Cytopathology</i> , 2009 , 37, 815-9	1.4	16

77	Alternative surgical strategies and favorable outcomes in patients with medullary thyroid carcinoma in Japan: experience of a single institution. <i>World Journal of Surgery</i> , 2009 , 33, 58-66	3.3	41
76	Clinicopathologic characteristics and prognosis of diffuse sclerosing variant of papillary thyroid carcinoma in Japan: an 18-year experience at a single institution. <i>World Journal of Surgery</i> , 2009 , 33, 958-62	3.3	40
75	Methimazole-induced agranulocytosis in patients with Graves Disease is more frequent with an initial dose of 30 mg daily than with 15 mg daily. <i>Thyroid</i> , 2009 , 19, 559-63	6.2	82
74	Loss of cellular polarity/cohesiveness in the invasive front of papillary thyroid carcinoma and periostin expression. <i>Cancer Letters</i> , 2009 , 281, 188-95	9.9	33
73	BRAF mutation in papillary thyroid carcinoma in a Japanese population: its lack of correlation with high-risk clinicopathological features and disease-free survival of patients. <i>Endocrine Journal</i> , 2009 , 56, 89-97	2.9	194
72	Prognosis of patients with papillary thyroid carcinoma having clinically apparent metastasis to the lateral compartment. <i>Endocrine Journal</i> , 2009 , 56, 759-66	2.9	116
71	Prognostic factors and therapeutic strategies for differentiated carcinomas of the thyroid. <i>Endocrine Journal</i> , 2009 , 56, 177-92	2.9	114
70	Prevalence of TSH receptor and Gsalpha mutations in 45 autonomously functioning thyroid nodules in Japan. <i>Endocrine Journal</i> , 2009 , 56, 791-8	2.9	28
69	Subclassification of non-solid-type papillary thyroid carcinoma identification of high-risk group in common type. <i>Cancer Science</i> , 2008 , 99, 1908-15	6.9	39
68	Serial changes in liver function tests in patients with thyrotoxicosis induced by Graves Disease and painless thyroiditis. <i>Thyroid</i> , 2008 , 18, 283-7	6.2	56
67	Evaluation of a new rapid and fully automated electrochemiluminescence immunoassay for thyrotropin receptor autoantibodies. <i>Thyroid</i> , 2008 , 18, 1157-64	6.2	35
66	3,5,3-Triiodothyronine thyrotoxicosis due to increased conversion of administered levothyroxine in patients with massive metastatic follicular thyroid carcinoma. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 2239-42	5.6	35
65	Papillary carcinoma obscured by complication with subacute thyroiditis: sequential ultrasonographic and histopathological findings in five cases. <i>Thyroid</i> , 2008 , 18, 1221-5	6.2	28
64	Prevalence and biological behaviour of variants of papillary thyroid carcinoma: experience at a single institute. <i>Pathology</i> , 2008 , 40, 617-22	1.6	43
63	Clinical characteristics of 852 patients with subacute thyroiditis before treatment. <i>Internal Medicine</i> , 2008 , 47, 725-9	1.1	117
62	Prevalence and prognostic significance of poor differentiation and tall cell variant in papillary carcinoma in Japan. <i>World Journal of Surgery</i> , 2008 , 32, 1535-43; discussion 1544-5	3.3	81
61	Biological behavior and prognosis of encapsulated papillary carcinoma of the thyroid: experience of a Japanese hospital for thyroid care. <i>World Journal of Surgery</i> , 2008 , 32, 1789-94	3.3	21
60	Ultrasonographic evaluation of thyroid nodules in 900 patients: comparison among ultrasonographic, cytological, and histological findings. <i>Thyroid</i> , 2007 , 17, 1269-76	6.2	79

59	Quantitative measurement of thyroid blood flow for differentiation of painless thyroiditis from Graves disease. <i>Clinical Endocrinology</i> , 2007 , 67, 41-5	3.4	81
58	Lack of association between BRAF V600E mutation and mitogen-activated protein kinase activation in papillary thyroid carcinoma. <i>Pathology International</i> , 2007 , 57, 12-20	1.8	28
57	Risk factors contributing to a poor prognosis of papillary thyroid carcinoma: validity of UICC/AJCC TNM classification and stage grouping. <i>World Journal of Surgery</i> , 2007 , 31, 838-48	3.3	105
56	Lateral and mediastinal lymph node dissection in differentiated thyroid carcinoma: indications, benefits, and risks. <i>World Journal of Surgery</i> , 2007 , 31, 905-15	3.3	79
55	Extranodal tumor extension to adjacent organs predicts a worse cause-specific survival in patients with papillary thyroid carcinoma. <i>World Journal of Surgery</i> , 2007 , 31, 1194-201	3.3	48
54	Prognosis and prognostic factors of follicular carcinoma in Japan: importance of postoperative pathological examination. <i>World Journal of Surgery</i> , 2007 , 31, 1417-24	3.3	73
53	Prognosis of patients with benign thyroid diseases accompanied by incidental papillary carcinoma undetectable on preoperative imaging tests. <i>World Journal of Surgery</i> , 2007 , 31, 1672-6	3.3	39
52	Lateral mobilization of the recurrent laryngeal nerve to facilitate tracheal surgery in patients with thyroid cancer invading the trachea near Berry ligament. <i>World Journal of Surgery</i> , 2007 , 31, 2081-4	3.3	16
51	Risk factors for recurrence to the lymph node in papillary thyroid carcinoma patients without preoperatively detectable lateral node metastasis: validity of prophylactic modified radical neck dissection. <i>World Journal of Surgery</i> , 2007 , 31, 2085-91	3.3	197
50	Results of a phase I clinical study using dendritic cell vaccinations for thyroid cancer. <i>Thyroid</i> , 2007 , 17, 53-8	6.2	19
49	A therapeutic strategy for incidentally detected papillary microcarcinoma of the thyroid. <i>Nature Clinical Practice Endocrinology and Metabolism</i> , 2007 , 3, 240-8		108
48	Quantitative measurement of anti-TSH receptor antibodies is useful for the evaluation of treatment effect. <i>Thyroid</i> , 2007 , 17, 85-6	6.2	3
47	Effect of levo-thyroxine replacement on non-high-density lipoprotein cholesterol in hypothyroid patients. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007 , 92, 608-11	5.6	50
46	Clinicopathologic significance of intrathyroidal epithelial thymoma/carcinoma showing thymus-like differentiation: a collaborative study with Member Institutes of The Japanese Society of Thyroid Surgery. <i>American Journal of Clinical Pathology</i> , 2007 , 127, 230-6	1.9	92
45	A novel thyrotropin receptor germline mutation (Asp617Tyr) causing hereditary hyperthyroidism. <i>Endocrine Journal</i> , 2007 , 54, 927-34	2.9	29
44	Comparison of methimazole and propylthiouracil in patients with hyperthyroidism caused by Graves disease. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007 , 92, 2157-62	5.6	194
43	Diagnosis of medullary thyroid carcinoma by calcitonin measurement in fine-needle aspiration biopsy specimens. <i>Thyroid</i> , 2007 , 17, 635-8	6.2	80
42	Prognostic significance of extrathyroid extension of papillary thyroid carcinoma: massive but not minimal extension affects the relapse-free survival. <i>World Journal of Surgery</i> , 2006 , 30, 780-6	3.3	205

41	Mutational analysis of the APC gene in cribriform-morula variant of papillary thyroid carcinoma. <i>World Journal of Surgery</i> , 2006 , 30, 775-9	3.3	42
40	Clinical significance of lymph node metastasis of thyroid papillary carcinoma located in one lobe. <i>World Journal of Surgery</i> , 2006 , 30, 1821-8	3.3	151
39	Transient vocal cord paralysis after fine-needle aspiration biopsy of thyroid tumor. <i>Thyroid</i> , 2006 , 16, 697-9	6.2	30
38	Usefulness of ultrasonography for diagnosis of malignant lymphoma of the thyroid. <i>Thyroid</i> , 2006 , 16, 983-7	6.2	53
37	Needle tract implantation of follicular neoplasm after fine-needle aspiration biopsy: report of a case. <i>Thyroid</i> , 2006 , 16, 1059-62	6.2	27
36	Sporadic congenital hyperthyroidism due to a germline mutation in the thyrotropin receptor gene (Leu 512 Gln) in a Japanese patient. <i>Endocrine Journal</i> , 2006 , 53, 735-40	2.9	38
35	Four cases of Graves disease which developed after painful Hashimoto thyroiditis. <i>Internal Medicine</i> , 2006 , 45, 385-9	1.1	29
34	A case with diffuse sclerosing variant of papillary carcinoma of the thyroid: characteristic features on ultrasonography. <i>Journal of Medical Ultrasonics (2001)</i> , 2006 , 33, 159-61	1.4	11
33	Ratio of serum free triiodothyronine to free thyroxine in Graves hyperthyroidism and thyrotoxicosis caused by painless thyroiditis. <i>Endocrine Journal</i> , 2005 , 52, 537-42	2.9	43
32	Recovery of parathyroid function after total thyroidectomy: long-term follow-up study. <i>ANZ Journal of Surgery</i> , 2005 , 75, 532-6	1	57
31	Usefulness of thyroglobulin measurement in fine-needle aspiration biopsy specimens for diagnosing cervical lymph node metastasis in patients with papillary thyroid cancer. <i>World Journal of Surgery</i> , 2005 , 29, 483-5	3.3	143
30	Needle tract implantation of papillary thyroid carcinoma after fine-needle aspiration biopsy. <i>World Journal of Surgery</i> , 2005 , 29, 1544-9	3.3	42
29	Stretching exercises to reduce symptoms of postoperative neck discomfort after thyroid surgery: prospective randomized study. <i>World Journal of Surgery</i> , 2005 , 29, 775-9	3.3	27
28	Ultrasonographically and anatomopathologically detectable node metastases in the lateral compartment as indicators of worse relapse-free survival in patients with papillary thyroid carcinoma. <i>World Journal of Surgery</i> , 2005 , 29, 917-20	3.3	148
27	Ill-defined edge on ultrasonographic examination can be a marker of aggressive characteristic of papillary thyroid microcarcinoma. <i>World Journal of Surgery</i> , 2005 , 29, 1007-11; discussion 1011-2	3.3	55
26	Diagnosis of follicular carcinoma of the thyroid: role of sonography in preoperative diagnosis of follicular nodules. <i>Journal of Medical Ultrasonics (2001)</i> , 2005 , 32, 153-8	1.4	23
25	Morules in cribriform-morular variant of papillary thyroid carcinoma: Immunohistochemical characteristics and distinction from squamous metaplasia. <i>Apmis</i> , 2004 , 112, 275-82	3.4	41
24	Preoperative ultrasonographic examination for lymph node metastasis: usefulness when designing lymph node dissection for papillary microcarcinoma of the thyroid. <i>World Journal of Surgery</i> , 2004 , 28, 498-501	3.3	171

23	Cribriform-morular variant of papillary thyroid carcinoma: clue to early detection of familial adenomatous polyposis-associated colon cancer. <i>World Journal of Surgery</i> , 2004 , 28, 886-9	3.3	83
22	Papillary microcarcinoma of the thyroid: how should it be treated?. <i>World Journal of Surgery</i> , 2004 , 28, 1115-21	3.3	191
21	Disturbed metabolism of remnant lipoproteins in patients with subclinical hypothyroidism. <i>American Journal of Medicine</i> , 2004 , 117, 696-9	2.4	24
20	Cribriform-morular variant of papillary thyroid carcinoma. Report of a case showing morules with peculiar nuclear clearing. <i>Acta Cytologica</i> , 2004 , 48, 431-6	3	22
19	Transient hyperthyroidism after withdrawal of antithyroid drugs in patients with Graves Disease. <i>Endocrine Journal</i> , 2004 , 51, 213-7	2.9	5
18	Changes in serum TSH receptor antibody (TRAb) values in patients with Graves Disease after total or subtotal thyroidectomy. <i>Endocrine Journal</i> , 2003 , 50, 595-601	2.9	29
17	Thyroid nodular lesion: analysis of cancer risk based on Kuma Hospital experience. <i>Pathology International</i> , 2003 , 53, 579-83	1.8	11
16	The relationship of psychological factors to the prognosis of hyperthyroidism in antithyroid drug-treated patients with Graves Disease. <i>Clinical Endocrinology</i> , 2003 , 58, 550-5	3.4	39
15	Serum concentrations of remnant-like particles in hypothyroid patients before and after thyroxine replacement. <i>Clinical Endocrinology</i> , 2003 , 58, 621-6	3.4	29
14	Cribriform-morular variant of papillary thyroid carcinoma: a pathological and molecular genetic study with evidence of frequent somatic mutations in exon 3 of the beta-catenin gene. <i>Journal of Pathology</i> , 2003 , 199, 58-67	9.4	112
13	An observation trial without surgical treatment in patients with papillary microcarcinoma of the thyroid. <i>Thyroid</i> , 2003 , 13, 381-7	6.2	582
12	Prospective trial of unilateral surgery for nonhereditary medullary thyroid carcinoma in patients without germline RET mutations. <i>World Journal of Surgery</i> , 2002 , 26, 1023-8	3.3	68
11	Recovery of parathyroid function after total thyroidectomy. <i>Surgery Today</i> , 2000 , 30, 333-8	3	31
10	Unilateral surgery supported by germline RET oncogene mutation analysis in patients with sporadic medullary thyroid carcinoma. <i>World Journal of Surgery</i> , 2000 , 24, 1367-72	3.3	34
9	Gene rearrangement of immunoglobulin as a marker of thyroid lymphoma. <i>World Journal of Surgery</i> , 1998 , 22, 558-61	3.3	15
8	The role of ansa-to-recurrent-laryngeal nerve anastomosis in operations for thyroid cancer. <i>The European Journal of Surgery</i> , 1998 , 164, 927-33		63
7	Thyroid cancer detected by ultrasound-guided fine-needle aspiration biopsy. <i>World Journal of Surgery</i> , 1996 , 20, 848-53; discussion 853	3.3	96
6	Accurate and simple method of diagnosing thyroid nodules the modified technique of ultrasound-guided fine needle aspiration biopsy. <i>Thyroid</i> , 1995 , 5, 141-5	6.2	129

5	Ansa-recurrent nerve anastomosis for vocal cord paralysis due to mediastinal lesions. <i>Annals of Thoracic Surgery</i> , 1994 , 57, 1020-1	2.7	15
4	Clinical aspects of primary thyroid lymphoma: diagnosis and treatment based on our experience of 119 cases. <i>Thyroid</i> , 1993 , 3, 93-9	6.2	209
3	Evaluation of surgical results and prediction of prognosis in patients with medullary thyroid carcinoma by analysis of serum calcitonin levels. <i>World Journal of Surgery</i> , 1988 , 12, 610-5	3.3	25
2	Intrathyroidal epithelial thymoma: an entity distinct from squamous cell carcinoma of the thyroid. <i>World Journal of Surgery</i> , 1985 , 9, 128-35	3.3	154
1	Relation of doubling time of plasma calcitonin levels to prognosis and recurrence of medullary thyroid carcinoma. <i>Annals of Surgery</i> , 1984 , 199, 461-6	7.8	88