

Erik Hedström

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7201138/publications.pdf>

Version: 2024-02-01

58
papers

1,713
citations

393982

19
h-index

288905

40
g-index

60
all docs

60
docs citations

60
times ranked

2233
citing authors

#	ARTICLE	IF	CITATIONS
1	Comparison of 2D and 4D Flow MRI in Neonates Without General Anesthesia. <i>Journal of Magnetic Resonance Imaging</i> , 2023, 57, 71-82.	1.9	4
2	Super-Resolution Cine Image Enhancement for Fetal Cardiac Magnetic Resonance Imaging. <i>Journal of Magnetic Resonance Imaging</i> , 2022, 56, 223-231.	1.9	10
3	Pulse Wave Velocity Measurements by Magnetic Resonance Imaging in Neonates and Adolescents: Methodological Aspects and Their Clinical Implications. <i>Pediatric Cardiology</i> , 2022, , 1.	0.6	3
4	Atrioventricular plane displacement versus mitral and tricuspid annular plane systolic excursion: A comparison between cardiac magnetic resonance and M-mode echocardiography. <i>Clinical Physiology and Functional Imaging</i> , 2021, 41, 262-270.	0.5	5
5	Utility of Fetal Cardiovascular Magnetic Resonance for Prenatal Diagnosis of Complex Congenital Heart Defects. <i>JAMA Network Open</i> , 2021, 4, e213538.	2.8	28
6	Hydraulic force is a novel mechanism of diastolic function that may contribute to decreased diastolic filling in HFpEF and facilitate filling in HFrEF. <i>Journal of Applied Physiology</i> , 2021, 130, 993-1000.	1.2	2
7	Free-breathing fetal cardiac MRI with doppler ultrasound gating, compressed sensing, and motion compensation. <i>Journal of Magnetic Resonance Imaging</i> , 2020, 51, 260-272.	1.9	25
8	Changes in left and right ventricular longitudinal function after pulmonary valve replacement in patients with Tetralogy of Fallot. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2020, 318, H345-H353.	1.5	6
9	Risk assessment in PAH using quantitative CMR tricuspid regurgitation: relation to heart catheterization. <i>ESC Heart Failure</i> , 2020, 7, 1653-1663.	1.4	4
10	Cardiovascular effects of severe late-onset preeclampsia are reversed within six months postpartum. <i>Pregnancy Hypertension</i> , 2020, 19, 18-24.	0.6	8
11	Alpha-1 microglobulin as a potential therapeutic candidate for treatment of hypertension and oxidative stress in the STOX1 preeclampsia mouse model. <i>Scientific Reports</i> , 2019, 9, 8561.	1.6	19
12	A new vessel segmentation algorithm for robust blood flow quantification from two-dimensional phase-contrast magnetic resonance images. <i>Clinical Physiology and Functional Imaging</i> , 2019, 39, 327-338.	0.5	15
13	Fetal iGRASP cine CMR assisting in prenatal diagnosis of complicated cardiac malformation with impact on delivery planning. <i>Clinical Physiology and Functional Imaging</i> , 2019, 39, 231-235.	0.5	9
14	Quantification of blood flow in the fetus with cardiovascular magnetic resonance imaging using Doppler ultrasound gating: validation against metric optimized gating. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2019, 21, 74.	1.6	19
15	Independent validation of metric optimized gating for fetal cardiovascular phase-contrast flow imaging. <i>Magnetic Resonance in Medicine</i> , 2019, 81, 495-503.	1.9	11
16	Altered biventricular hemodynamic forces in patients with repaired tetralogy of Fallot and right ventricular volume overload because of pulmonary regurgitation. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2018, 315, H1691-H1702.	1.5	24
17	Dynamic fetal cardiovascular magnetic resonance imaging using Doppler ultrasound gating. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2018, 20, 17.	1.6	55
18	Automatic T2* determination for quantification of iron load in heart and liver: a comparison between automatic inline Maximum Likelihood Estimate and the truncation and offset methods. <i>Clinical Physiology and Functional Imaging</i> , 2017, 37, 299-304.	0.5	4

#	ARTICLE	IF	CITATIONS
19	Self-gated fetal cardiac MRI with tiny golden angle iGRASP: A feasibility study. <i>Journal of Magnetic Resonance Imaging</i> , 2017, 46, 207-217.	1.9	45
20	The effect of initial teaching on evaluation of left ventricular volumes by cardiovascular magnetic resonance imaging: comparison between complete and intermediate beginners and experienced observers. <i>BMC Medical Imaging</i> , 2017, 17, 33.	1.4	5
21	Longitudinal shortening remains the principal component of left ventricular pumping in patients with chronic myocardial infarction even when the absolute atrioventricular plane displacement is decreased. <i>BMC Cardiovascular Disorders</i> , 2017, 17, 208.	0.7	15
22	Required temporal resolution for accurate thoracic aortic pulse wave velocity measurements by phase-contrast magnetic resonance imaging and comparison with clinical standard applanation tonometry. <i>BMC Cardiovascular Disorders</i> , 2016, 16, 110.	0.7	15
23	Validation of T1 and T2 algorithms for quantitative MRI: performance by a vendor-independent software. <i>BMC Medical Imaging</i> , 2016, 16, 46.	1.4	12
24	Validation of a new t_2^* algorithm and its uncertainty value for cardiac and liver iron load determination from MRI magnitude images. <i>Magnetic Resonance in Medicine</i> , 2016, 75, 1717-1729.	1.9	9
25	Using a modified 3D-printer for mapping the magnetic field of RF coils designed for fetal and neonatal imaging. <i>Journal of Magnetic Resonance</i> , 2016, 269, 146-151.	1.2	4
26	Validation and Development of a New Automatic Algorithm for Time-Resolved Segmentation of the Left Ventricle in Magnetic Resonance Imaging. <i>BioMed Research International</i> , 2015, 2015, 1-12.	0.9	33
27	Diagnostic performance of the Selvester QRS scoring system in relation to clinical ECG assessment of patients with lateral myocardial infarction using cardiac magnetic resonance as reference standard. <i>Journal of Electrocardiology</i> , 2015, 48, 750-757.	0.4	2
28	A new validated T_2^* analysis method with certainty estimates for cardiac and liver iron load determination. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2015, 17, P52.	1.6	2
29	Ruptured Aneurysm of the Sinus of Valsalva. <i>Journal of the American College of Cardiology</i> , 2012, 59, 538.	1.2	3
30	Quantification of Absolute Myocardial Perfusion in Patients With Coronary Artery Disease. <i>Journal of the American College of Cardiology</i> , 2012, 60, 1546-1555.	1.2	206
31	Noncompaction of the Myocardium. <i>Journal of the American College of Cardiology</i> , 2011, 58, e25.	1.2	3
32	Measurements of wound edge microvascular blood flow during negative pressure wound therapy using thermodiffusion and transcutaneous and invasive laser Doppler velocimetry. <i>Wound Repair and Regeneration</i> , 2011, 19, 727-733.	1.5	46
33	The influence of different sizes and types of wound fillers on wound contraction and tissue pressure during negative pressure wound therapy. <i>International Wound Journal</i> , 2011, 8, 336-342.	1.3	30
34	The evaluation of an electrocardiographic myocardial ischemia acuteness score to predict the amount of myocardial salvage achieved by early percutaneous coronary intervention. <i>Journal of Electrocardiology</i> , 2011, 44, 525-532.	0.4	16
35	An automatic method for quantification of myocardium at risk from myocardial perfusion SPECT in patients with acute coronary occlusion. <i>Journal of Nuclear Cardiology</i> , 2010, 17, 831-840.	1.4	9
36	Cardiovascular magnetic resonance of the myocardium at risk in acute reperfused myocardial infarction: comparison of T2-weighted imaging versus the circumferential endocardial extent of late gadolinium enhancement with transmural projection. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2010, 12, 18.	1.6	42

#	ARTICLE	IF	CITATIONS
37	Effects of gadolinium contrast agent on aortic blood flow and myocardial strain measurements by phase-contrast cardiovascular magnetic resonance. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2010, 12, 70.	1.6	8
38	Biochemical markers of inflammatory response and their relation to myocardial injury. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2010, 12, .	1.6	0
39	Consideration of the Impact of Reperfusion Therapy on the Quantitative Relationship between the Selvester QRS Score and Infarct Size by Cardiac MRI. <i>Annals of Noninvasive Electrocardiology</i> , 2010, 15, 238-244.	0.5	14
40	Rapid Initial Reduction of Hyperenhanced Myocardium After Reperfused First Myocardial Infarction Suggests Recovery of the Peri-Infarction Zone. <i>Circulation: Cardiovascular Imaging</i> , 2009, 2, 47-55.	1.3	113
41	Location of myocardium at risk in patients with first-time ST-elevation infarction: comparison among single photon emission computed tomography, magnetic resonance imaging, and electrocardiography. <i>Journal of Electrocardiology</i> , 2009, 42, 198-203.	0.4	13
42	The Dipolar ElectroCARDioTOpographic (DECARTO)â€“like method for graphic presentation of location and extent of area at risk estimated from ST-segment deviations in patients with acute myocardial infarction. <i>Journal of Electrocardiology</i> , 2009, 42, 172-180.	0.4	18
43	Age and gender specific normal values of left ventricular mass, volume and function for gradient echo magnetic resonance imaging: a cross sectional study. <i>BMC Medical Imaging</i> , 2009, 9, 2.	1.4	169
44	Infarct evolution in man studied in patients with first-time coronary occlusion in comparison to different species - implications for assessment of myocardial salvage. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2009, 11, 38.	1.6	95
45	Myocardium at Risk After Acute Infarction in Humans on Cardiac Magnetic Resonance. <i>JACC: Cardiovascular Imaging</i> , 2009, 2, 569-576.	2.3	184
46	Spline-Based Cardiac Motion Tracking Using Velocity-Encoded Magnetic Resonance Imaging. <i>IEEE Transactions on Medical Imaging</i> , 2008, 27, 1045-1053.	5.4	16
47	Peak CKMB and cTnT accurately estimates myocardial infarct size after reperfusion. <i>Scandinavian Cardiovascular Journal</i> , 2007, 41, 44-50.	0.4	41
48	The endocardial extent of reperfused first-time myocardial infarction is more predictive of pathologic Q waves than is infarct transmural: a magnetic resonance imaging study. <i>Clinical Physiology and Functional Imaging</i> , 2007, 27, 101-108.	0.5	23
49	Physiological determinants of the variation in left ventricular mass from early adolescence to late adulthood in healthy subjects. <i>Clinical Physiology and Functional Imaging</i> , 2007, 27, 254-262.	0.5	20
50	Left ventricular mass by 12-lead electrocardiogram in healthy subjects: comparison to cardiac magnetic resonance imaging. <i>Journal of Electrocardiology</i> , 2006, 39, 67-72.	0.4	33
51	Importance of perfusion in myocardial viability studies using delayed contrast-enhanced magnetic resonance imaging. <i>Journal of Magnetic Resonance Imaging</i> , 2006, 24, 77-83.	1.9	9
52	Infarct transmural and adjacent segmental function as determinants of wall thickening in revascularized chronic ischemic heart disease. <i>Clinical Physiology and Functional Imaging</i> , 2005, 25, 209-214.	0.5	6
53	Physiological determinants of the variation in left ventricular mass from early adolescence to late adulthood in healthy subjects. <i>Clinical Physiology and Functional Imaging</i> , 2005, 25, 332-339.	0.5	15
54	Size and transmural extent of first-time reperfused myocardial infarction assessed by cardiac magnetic resonance can be estimated by 12-lead electrocardiogram. <i>American Heart Journal</i> , 2005, 150, 920.e1-920.e9.	1.2	49

#	ARTICLE	IF	CITATIONS
55	Semi-automatic quantification of myocardial infarction from delayed contrast enhanced magnetic resonance imaging. <i>Scandinavian Cardiovascular Journal</i> , 2005, 39, 267-275.	0.4	86
56	Determination of the left ventricular long-axis orientation from a single short-axis MR image: relation to BMI and age. <i>Clinical Physiology and Functional Imaging</i> , 2004, 24, 310-315.	0.5	26
57	Myocardial SPECT perfusion defect size compared to infarct size by delayed gadolinium-enhanced magnetic resonance imaging in patients with acute or chronic infarction. <i>Clinical Physiology and Functional Imaging</i> , 2004, 24, 380-386.	0.5	16
58	A method for assembling a collaborative research team from multiple disciplines and academic centers to study the relationships between ECG estimation and MRI measurement of myocardial infarct size. <i>Journal of Electrocardiology</i> , 2001, 34, 1-6.	0.4	11