

Gregory B Martin

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7202382/gregory-b-martin-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

193
papers

20,563
citations

74
h-index

142
g-index

205
ext. papers

23,040
ext. citations

9.3
avg, IF

6.59
L-index

#	Paper	IF	Citations
193	Integrative Proteomic and Phosphoproteomic Analyses of Pattern- and Effector-Triggered Immunity in Tomato.. <i>Frontiers in Plant Science</i> , 2021 , 12, 768693	6.2	2
192	Spelling Changes and Fluorescent Tagging With Prime Editing Vectors for Plants. <i>Frontiers in Genome Editing</i> , 2021 , 3, 617553	2.5	9
191	WRKY22 and WRKY25 transcription factors are positive regulators of defense responses in <i>Nicotiana benthamiana</i> . <i>Plant Molecular Biology</i> , 2021 , 105, 65-82	4.6	8
190	Tomato Wall-Associated Kinase SlWak1 Depends on Fls2/Fls3 to Promote Apoplastic Immune Responses to. <i>Plant Physiology</i> , 2020 , 183, 1869-1882	6.6	20
189	Ptr1 evolved convergently with RPS2 and Mr5 to mediate recognition of AvrRpt2 in diverse solanaceous species. <i>Plant Journal</i> , 2020 , 103, 1433-1445	6.9	9
188	Molecular Characterization of Differences between the Tomato Immune Receptors Flagellin Sensing 3 and Flagellin Sensing 2. <i>Plant Physiology</i> , 2020 , 183, 1825-1837	6.6	5
187	Generation and Molecular Characterization of CRISPR/Cas9-Induced Mutations in 63 Immunity-Associated Genes in Tomato Reveals Specificity and a Range of Gene Modifications. <i>Frontiers in Plant Science</i> , 2020 , 11, 10	6.2	20
186	Genome of <i>Solanum pimpinellifolium</i> provides insights into structural variants during tomato breeding. <i>Nature Communications</i> , 2020 , 11, 5817	17.4	24
185	Plant Genome Editing Database (PGED): A Call for Submission of Information about Genome-Edited Plant Mutants. <i>Molecular Plant</i> , 2019 , 12, 127-129	14.4	11
184	The tomato Pto gene confers resistance to <i>Pseudomonas floricida</i> , an emergent plant pathogen with just nine type III effectors. <i>Plant Pathology</i> , 2019 , 68, 977-984	2.8	2
183	PP2C phosphatase Pic1 negatively regulates the phosphorylation status of Pti1b kinase, a regulator of flagellin-triggered immunity in tomato. <i>Biochemical Journal</i> , 2019 , 476, 1621-1635	3.8	6
182	Transcriptome-based identification and validation of reference genes for plant-bacteria interaction studies using <i>Nicotiana benthamiana</i> . <i>Scientific Reports</i> , 2019 , 9, 1632	4.9	18
181	Natural variation for unusual host responses and flagellin-mediated immunity against <i>Pseudomonas syringae</i> in genetically diverse tomato accessions. <i>New Phytologist</i> , 2019 , 223, 447-461	9.8	12
180	Mai1 Protein Acts Between Host Recognition of Pathogen Effectors and Mitogen-Activated Protein Kinase Signaling. <i>Molecular Plant-Microbe Interactions</i> , 2019 , 32, 1496-1507	3.6	7
179	The Locus of Confers Resistance to Race 1 Strains of pv. and to by Recognizing the Type III Effectors AvrRpt2 and RipBN. <i>Molecular Plant-Microbe Interactions</i> , 2019 , 32, 949-960	3.6	17
178	Virus-induced gene silencing database for phenomics and functional genomics in. <i>Plant Direct</i> , 2018 , 2, e00055	3.3	11
177	The Bacterial Effector AvrPto Targets the Regulatory Coreceptor SOBIR1 and Suppresses Defense Signaling Mediated by the Receptor-Like Protein Cf-4. <i>Molecular Plant-Microbe Interactions</i> , 2018 , 31, 75-85	3.6	10

176	Pseudomonas syringae pv. tomato Strains from New York Exhibit Virulence Attributes Intermediate Between Typical Race 0 and Race 1 Strains. <i>Plant Disease</i> , 2017 , 101, 1442-1448	1.5	5
175	A Subset of Ubiquitin-Conjugating Enzymes Is Essential for Plant Immunity. <i>Plant Physiology</i> , 2017 , 173, 1371-1390	6.6	32
174	Generation of a Collection of Mutant Tomato Lines Using Pooled CRISPR Libraries. <i>Plant Physiology</i> , 2017 , 174, 2023-2037	6.6	55
173	The Tomato Kinase Pti1 Contributes to Production of Reactive Oxygen Species in Response to Two Flagellin-Derived Peptides and Promotes Resistance to Pseudomonas syringae Infection. <i>Molecular Plant-Microbe Interactions</i> , 2017 , 30, 725-738	3.6	14
172	Use of RNA-seq data to identify and validate RT-qPCR reference genes for studying the tomato-Pseudomonas pathosystem. <i>Scientific Reports</i> , 2017 , 7, 44905	4.9	39
171	Detecting the interaction of peptide ligands with plant membrane receptors. <i>Current Protocols in Plant Biology</i> , 2017 , 2, 240-269	2.8	0
170	Ser360 and Ser364 in the Kinase Domain of Tomato SLMAPKKK Are Critical for Programmed Cell Death Associated with Plant Immunity. <i>Plant Pathology Journal</i> , 2017 , 33, 163-169	2.5	3
169	Tomato receptor FLAGELLIN-SENSING 3 binds flgII-28 and activates the plant immune system. <i>Nature Plants</i> , 2016 , 2, 16128	11.5	109
168	High-throughput CRISPR Vector Construction and Characterization of DNA Modifications by Generation of Tomato Hairy Roots. <i>Journal of Visualized Experiments</i> , 2016 ,	1.6	17
167	A novel method of transcriptome interpretation reveals a quantitative suppressive effect on tomato immune signaling by two domains in a single pathogen effector protein. <i>BMC Genomics</i> , 2016 , 17, 229	4.5	5
166	Natural Variation in Tomato Reveals Differences in the Recognition of AvrPto and AvrPtoB Effectors from Pseudomonas syringae. <i>Molecular Plant</i> , 2016 , 9, 639-649	14.4	10
165	Detecting N-myristoylation and S-acylation of host and pathogen proteins in plants using click chemistry. <i>Plant Methods</i> , 2016 , 12, 38	5.8	15
164	iTAK: A Program for Genome-wide Prediction and Classification of Plant Transcription Factors, Transcriptional Regulators, and Protein Kinases. <i>Molecular Plant</i> , 2016 , 9, 1667-1670	14.4	352
163	Greasy tactics in the plant-pathogen molecular arms race. <i>Journal of Experimental Botany</i> , 2015 , 66, 1607-16		16
162	Acquisition of Iron Is Required for Growth of Salmonella spp. in Tomato Fruit. <i>Applied and Environmental Microbiology</i> , 2015 , 81, 3663-70	4.8	8
161	Functional genomics of tomato for the study of plant immunity. <i>Briefings in Functional Genomics</i> , 2015 , 14, 291-301	4.9	12
160	Five Xanthomonas type III effectors suppress cell death induced by components of immunity-associated MAP kinase cascades. <i>Plant Signaling and Behavior</i> , 2015 , 10, e1064573	2.5	15
159	Complete Genome Sequence of a Tomato-Infecting Tomato Mottle Mosaic Virus in New York. <i>Genome Announcements</i> , 2015 , 3,		7

158	Identification of a Candidate Gene in <i>Solanum habrochaites</i> for Resistance to a Race 1 Strain of <i>Pseudomonas syringae</i> pv. tomato. <i>Plant Genome</i> , 2015 , 8, eplantgenome2015.02.0006	4.4	9
157	<i>Pseudomonas syringae</i> pv. tomato DC3000 Type III Secretion Effector Polymutants Reveal an Interplay between HopAD1 and AvrPtoB. <i>Cell Host and Microbe</i> , 2015 , 17, 752-62	23.4	66
156	The SGN VIGS tool: user-friendly software to design virus-induced gene silencing (VIGS) constructs for functional genomics. <i>Molecular Plant</i> , 2015 , 8, 486-8	14.4	86
155	Comparative genomics and phylogenetic discordance of cultivated tomato and close wild relatives. <i>PeerJ</i> , 2015 , 3, e793	3.1	18
154	Natural variation for responsiveness to flg22, flgII-28, and csp22 and <i>Pseudomonas syringae</i> pv. tomato in heirloom tomatoes. <i>PLoS ONE</i> , 2014 , 9, e106119	3.7	25
153	Transcriptomic analysis reveals tomato genes whose expression is induced specifically during effector-triggered immunity and identifies the Epk1 protein kinase which is required for the host response to three bacterial effector proteins. <i>Genome Biology</i> , 2014 , 15, 492	18.3	52
152	Pto kinase binds two domains of AvrPtoB and its proximity to the effector E3 ligase determines if it evades degradation and activates plant immunity. <i>PLoS Pathogens</i> , 2014 , 10, e1004227	7.6	44
151	Analysis of wild-species introgressions in tomato inbreds uncovers ancestral origins. <i>BMC Plant Biology</i> , 2014 , 14, 287	5.3	19
150	Transcriptomics-based screen for genes induced by flagellin and repressed by pathogen effectors identifies a cell wall-associated kinase involved in plant immunity. <i>Genome Biology</i> , 2013 , 14, R139	18.3	92
149	Salmonella colonization activates the plant immune system and benefits from association with plant pathogenic bacteria. <i>Environmental Microbiology</i> , 2013 , 15, 2418-30	5.2	44
148	Thymoquinone causes multiple effects, including cell death, on dividing plant cells. <i>Comptes Rendus - Biologies</i> , 2013 , 336, 546-56	1.4	3
147	Two leucines in the N-terminal MAPK-docking site of tomato SIMKK2 are critical for interaction with a downstream MAPK to elicit programmed cell death associated with plant immunity. <i>FEBS Letters</i> , 2013 , 587, 1460-5	3.8	8
146	Allelic variation in two distinct <i>Pseudomonas syringae</i> flagellin epitopes modulates the strength of plant immune responses but not bacterial motility. <i>New Phytologist</i> , 2013 , 200, 847-860	9.8	91
145	The tomato Fni3 lysine-63-specific ubiquitin-conjugating enzyme and suv ubiquitin E2 variant positively regulate plant immunity. <i>Plant Cell</i> , 2013 , 25, 3615-31	11.6	39
144	The tomato calcium sensor Cbl10 and its interacting protein kinase Cipk6 define a signaling pathway in plant immunity. <i>Plant Cell</i> , 2013 , 25, 2748-64	11.6	84
143	Nonhost resistance of tomato to the bean pathogen <i>Pseudomonas syringae</i> pv. <i>syringae</i> B728a is due to a defective E3 ubiquitin ligase domain in <i>avrptobb728a</i> . <i>Molecular Plant-Microbe Interactions</i> , 2013 , 26, 387-97	3.6	10
142	Plant programmed cell death caused by an autoactive form of Prf is suppressed by co-expression of the Prf LRR domain. <i>Molecular Plant</i> , 2012 , 5, 1058-67	14.4	15
141	Molecular Mechanisms Involved in the Interaction Between Tomato and <i>Pseudomonas syringae</i> pv. tomato 2012 , 187-209		

140	A draft genome sequence of <i>Nicotiana benthamiana</i> to enhance molecular plant-microbe biology research. <i>Molecular Plant-Microbe Interactions</i> , 2012 , 25, 1523-30	3.6	311
139	A tomato LysM receptor-like kinase promotes immunity and its kinase activity is inhibited by AvrPtoB. <i>Plant Journal</i> , 2012 , 69, 92-103	6.9	90
138	Type III secretion and effectors shape the survival and growth pattern of <i>Pseudomonas syringae</i> on leaf surfaces. <i>Plant Physiology</i> , 2012 , 158, 1803-18	6.6	48
137	The Bsubunit of the SnRK1 complex is phosphorylated by the plant cell death suppressor Adi3. <i>Plant Physiology</i> , 2012 , 159, 1277-90	6.6	30
136	Structural analysis of <i>Pseudomonas syringae</i> AvrPtoB bound to host BAK1 reveals two similar kinase-interacting domains in a type III Effector. <i>Cell Host and Microbe</i> , 2011 , 10, 616-26	23.4	99
135	Effector-triggered immunity mediated by the Pto kinase. <i>Trends in Plant Science</i> , 2011 , 16, 132-40	13.1	88
134	Genetic disassembly and combinatorial reassembly identify a minimal functional repertoire of type III effectors in <i>Pseudomonas syringae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 2975-80	11.5	152
133	Tomato 14-3-3 protein TFT7 interacts with a MAP kinase kinase to regulate immunity-associated programmed cell death mediated by diverse disease resistance proteins. <i>Journal of Biological Chemistry</i> , 2011 , 286, 14129-36	5.4	58
132	Two virulence determinants of type III effector AvrPto are functionally conserved in diverse <i>Pseudomonas syringae</i> pathovars. <i>New Phytologist</i> , 2010 , 187, 969-982	9.8	17
131	Phosphorylation of the <i>Pseudomonas syringae</i> effector AvrPto is required for FLS2/BAK1-independent virulence activity and recognition by tobacco. <i>Plant Journal</i> , 2010 , 61, 16-24	6.9	32
130	A secreted effector protein (SNE1) from <i>Phytophthora infestans</i> is a broadly acting suppressor of programmed cell death. <i>Plant Journal</i> , 2010 , 62, 357-66	6.9	92
129	Tomato 14-3-3 protein 7 positively regulates immunity-associated programmed cell death by enhancing protein abundance and signaling ability of MAPKKK {alpha}. <i>Plant Cell</i> , 2010 , 22, 260-72	11.6	110
128	Endosome-associated CRT1 functions early in resistance gene-mediated defense signaling in <i>Arabidopsis</i> and tobacco. <i>Plant Cell</i> , 2010 , 22, 918-36	11.6	51
127	Identification of <i>Nicotiana benthamiana</i> genes involved in pathogen-associated molecular pattern-triggered immunity. <i>Molecular Plant-Microbe Interactions</i> , 2010 , 23, 715-26	3.6	62
126	Methods to study PAMP-triggered immunity using tomato and <i>Nicotiana benthamiana</i> . <i>Molecular Plant-Microbe Interactions</i> , 2010 , 23, 991-9	3.6	117
125	The T-loop extension of the tomato protein kinase AvrPto-dependent Pto-interacting protein 3 (Adi3) directs nuclear localization for suppression of plant cell death. <i>Journal of Biological Chemistry</i> , 2010 , 285, 17584-94	5.4	26
124	Deletions in the repertoire of <i>Pseudomonas syringae</i> pv. tomato DC3000 type III secretion effector genes reveal functional overlap among effectors. <i>PLoS Pathogens</i> , 2009 , 5, e1000388	7.6	192
123	Crystal structure of the complex between <i>Pseudomonas</i> effector AvrPtoB and the tomato Pto kinase reveals both a shared and a unique interface compared with AvrPto-Pto. <i>Plant Cell</i> , 2009 , 21, 1846-59	11.6	67

122	Advances in experimental methods for the elucidation of <i>Pseudomonas syringae</i> effector function with a focus on AvrPtoB. <i>Molecular Plant Pathology</i> , 2009 , 10, 777-93	5.7	18
121	Virus-induced gene silencing (VIGS) in <i>Nicotiana benthamiana</i> and tomato. <i>Journal of Visualized Experiments</i> , 2009 ,	1.6	71
120	<i>Xanthomonas</i> T3S Effector XopN Suppresses PAMP-Triggered Immunity and Interacts with a Tomato Atypical Receptor-Like Kinase and TFT1. <i>Plant Cell</i> , 2009 , 21, 1305-23	11.6	133
119	A draft genome sequence of <i>Pseudomonas syringae</i> pv. tomato T1 reveals a type III effector repertoire significantly divergent from that of <i>Pseudomonas syringae</i> pv. tomato DC3000. <i>Molecular Plant-Microbe Interactions</i> , 2009 , 22, 52-62	3.6	109
118	Assay for pathogen-associated molecular pattern (PAMP)-triggered immunity (PTI) in plants. <i>Journal of Visualized Experiments</i> , 2009 ,	1.6	8
117	Bacterial effectors target the common signaling partner BAK1 to disrupt multiple MAMP receptor-signaling complexes and impede plant immunity. <i>Cell Host and Microbe</i> , 2008 , 4, 17-27	23.4	410
116	A bacterial E3 ubiquitin ligase targets a host protein kinase to disrupt plant immunity. <i>Nature</i> , 2007 , 448, 370-4	50.4	248
115	An NB-LRR protein required for HR signalling mediated by both extra- and intracellular resistance proteins. <i>Plant Journal</i> , 2007 , 50, 14-28	6.9	137
114	A <i>Pseudomonas syringae</i> pv. tomato DC3000 mutant lacking the type III effector HopQ1-1 is able to cause disease in the model plant <i>Nicotiana benthamiana</i> . <i>Plant Journal</i> , 2007 , 51, 32-46	6.9	205
113	The N-terminal region of <i>Pseudomonas</i> type III effector AvrPtoB elicits Pto-dependent immunity and has two distinct virulence determinants. <i>Plant Journal</i> , 2007 , 52, 595-614	6.9	69
112	DspA/E, a type III effector of <i>Erwinia amylovora</i> , is required for early rapid growth in <i>Nicotiana benthamiana</i> and causes NbSGT1-dependent cell death. <i>Molecular Plant Pathology</i> , 2007 , 8, 255-65	5.7	31
111	Aconitase plays a role in regulating resistance to oxidative stress and cell death in <i>Arabidopsis</i> and <i>Nicotiana benthamiana</i> . <i>Plant Molecular Biology</i> , 2007 , 63, 273-87	4.6	108
110	<i>Pseudomonas syringae</i> type III effector AvrPtoB is phosphorylated in plant cells on serine 258, promoting its virulence activity. <i>Journal of Biological Chemistry</i> , 2007 , 282, 30737-44	5.4	31
109	Manipulation of plant programmed cell death pathways during plant-pathogen interactions. <i>Plant Signaling and Behavior</i> , 2007 , 2, 188-9	2.5	10
108	Identification and characterization of plant genes involved in <i>Agrobacterium</i> -mediated plant transformation by virus-induced gene silencing. <i>Molecular Plant-Microbe Interactions</i> , 2007 , 20, 41-52	3.6	70
107	Pto- and Prf-mediated recognition of AvrPto and AvrPtoB restricts the ability of diverse <i>Pseudomonas syringae</i> pathovars to infect tomato. <i>Molecular Plant-Microbe Interactions</i> , 2007 , 20, 806-15	3.6	56
106	Host-mediated phosphorylation of type III effector AvrPto promotes <i>Pseudomonas</i> virulence and avirulence in tomato. <i>Plant Cell</i> , 2006 , 18, 502-14	11.6	58
105	Type III effector AvrPtoB requires intrinsic E3 ubiquitin ligase activity to suppress plant cell death and immunity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006 , 103, 2851-6	11.5	178

104	Diverse AvrPtoB homologs from several <i>Pseudomonas syringae</i> pathovars elicit Pto-dependent resistance and have similar virulence activities. <i>Applied and Environmental Microbiology</i> , 2006 , 72, 702-124.8	55
103	A bacterial inhibitor of host programmed cell death defenses is an E3 ubiquitin ligase. <i>Science</i> , 2006 , 311, 222-6	33.3 273
102	Comparative genomics of host-specific virulence in <i>Pseudomonas syringae</i> . <i>Genetics</i> , 2006 , 174, 1041-564	117
101	Specific bacterial suppressors of MAMP signaling upstream of MAPKKK in Arabidopsis innate immunity. <i>Cell</i> , 2006 , 125, 563-75	56.2 341
100	Whole-genome expression profiling defines the HrpL regulon of <i>Pseudomonas syringae</i> pv. tomato DC3000, allows de novo reconstruction of the Hrp cis element, and identifies novel coregulated genes. <i>Molecular Plant-Microbe Interactions</i> , 2006 , 19, 1167-79	3.6 93
99	A novel link between tomato GRAS genes, plant disease resistance and mechanical stress response. <i>Molecular Plant Pathology</i> , 2006 , 7, 593-604	5.7 70
98	Bacterial elicitation and evasion of plant innate immunity. <i>Nature Reviews Molecular Cell Biology</i> , 2006 , 7, 601-11	48.7 315
97	Adi3 is a Pdk1-interacting AGC kinase that negatively regulates plant cell death. <i>EMBO Journal</i> , 2006 , 25, 255-65	13 68
96	An avrPto/avrPtoB mutant of <i>Pseudomonas syringae</i> pv. tomato DC3000 does not elicit Pto-mediated resistance and is less virulent on tomato. <i>Molecular Plant-Microbe Interactions</i> , 2005 , 18, 43-51	3.6 109
95	AvrPtoB: a bacterial type III effector that both elicits and suppresses programmed cell death associated with plant immunity. <i>FEMS Microbiology Letters</i> , 2005 , 245, 1-8	2.9 56
94	<i>Pseudomonas syringae</i> pv. tomato type III effectors AvrPto and AvrPtoB promote ethylene-dependent cell death in tomato. <i>Plant Journal</i> , 2005 , 44, 139-54	6.9 93
93	Role of mitogen-activated protein kinases in plant immunity. <i>Current Opinion in Plant Biology</i> , 2005 , 8, 541-7	9.9 237
92	Calmodulin-like proteins from Arabidopsis and tomato are involved in host defense against <i>Pseudomonas syringae</i> pv. tomato. <i>Plant Molecular Biology</i> , 2005 , 58, 887-897	4.6 94
91	Transcriptome and selected metabolite analyses reveal multiple points of ethylene control during tomato fruit development. <i>Plant Cell</i> , 2005 , 17, 2954-65	11.6 400
90	Gene profiling of a compatible interaction between <i>Phytophthora infestans</i> and <i>Solanum tuberosum</i> suggests a role for carbonic anhydrase. <i>Molecular Plant-Microbe Interactions</i> , 2005 , 18, 913-223.6	132
89	Suppression of pathogen-inducible NO synthase (iNOS) activity in tomato increases susceptibility to <i>Pseudomonas syringae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 8239-44	11.5 17
88	PeerGAD: a peer-review-based and community-centric web application for viewing and annotating prokaryotic genome sequences. <i>Nucleic Acids Research</i> , 2004 , 32, 3124-35	20.1 12
87	Identification of MAPKs and their possible MAPK kinase activators involved in the Pto-mediated defense response of tomato. <i>Journal of Biological Chemistry</i> , 2004 , 279, 49229-35	5.4 90

86	Silencing of subfamily I of protein phosphatase 2A catalytic subunits results in activation of plant defense responses and localized cell death. <i>Plant Journal</i> , 2004 , 38, 563-77	6.9	98
85	Applications and advantages of virus-induced gene silencing for gene function studies in plants. <i>Plant Journal</i> , 2004 , 39, 734-46	6.9	530
84	Comprehensive EST analysis of tomato and comparative genomics of fruit ripening. <i>Plant Journal</i> , 2004 , 40, 47-59	6.9	196
83	MAPKKKalpha is a positive regulator of cell death associated with both plant immunity and disease. <i>EMBO Journal</i> , 2004 , 23, 3072-82	13	262
82	The solution structure of type III effector protein AvrPto reveals conformational and dynamic features important for plant pathogenesis. <i>Structure</i> , 2004 , 12, 1257-68	5.2	43
81	Strategies used by bacterial pathogens to suppress plant defenses. <i>Current Opinion in Plant Biology</i> , 2004 , 7, 356-64	9.9	180
80	Strategies used by bacterial pathogens to suppress plant defenses. <i>Current Opinion in Plant Biology</i> , 2004 , 7, 356-356	9.9	
79	Identification and expression profiling of tomato genes differentially regulated during a resistance response to <i>Xanthomonas campestris</i> pv. <i>vesicatoria</i> . <i>Molecular Plant-Microbe Interactions</i> , 2004 , 17, 1212-22	3.6	47
78	Molecular Mechanisms Involved in Bacterial Speck Disease Resistance of Tomato. <i>Plant Pathology Journal</i> , 2004 , 20, 7-12	2.5	15
77	Overexpression of the disease resistance gene Pto in tomato induces gene expression changes similar to immune responses in human and fruitfly. <i>Plant Physiology</i> , 2003 , 132, 1901-12	6.6	44
76	<i>Pseudomonas</i> type III effector AvrPtoB induces plant disease susceptibility by inhibition of host programmed cell death. <i>EMBO Journal</i> , 2003 , 22, 60-9	13	316
75	Partial resistance of tomato to <i>Phytophthora infestans</i> is not dependent upon ethylene, jasmonic acid, or salicylic acid signaling pathways. <i>Molecular Plant-Microbe Interactions</i> , 2003 , 16, 141-8	3.6	54
74	Understanding the functions of plant disease resistance proteins. <i>Annual Review of Plant Biology</i> , 2003 , 54, 23-61	30.7	731
73	Two MAPK cascades, NPR1, and TGA transcription factors play a role in Pto-mediated disease resistance in tomato. <i>Plant Journal</i> , 2003 , 36, 905-17	6.9	263
72	Molecular basis of Pto-mediated resistance to bacterial speck disease in tomato. <i>Annual Review of Phytopathology</i> , 2003 , 41, 215-43	10.8	252
71	The complete genome sequence of the Arabidopsis and tomato pathogen <i>Pseudomonas syringae</i> pv. <i>tomato</i> DC3000. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 10181-6	11.5	695
70	The tomato transcription factor Pti4 regulates defense-related gene expression via GCC box and non-GCC box cis elements. <i>Plant Cell</i> , 2003 , 15, 3033-50	11.6	204
69	Location and activity of members of a family of virPphA homologues in pathovars of <i>Pseudomonas syringae</i> and <i>P. savastanoi</i> . <i>Molecular Plant Pathology</i> , 2002 , 3, 205-16	5.7	37

68	Comprehensive transcript profiling of Pto- and Prf-mediated host defense responses to infection by <i>Pseudomonas syringae</i> pv. tomato. <i>Plant Journal</i> , 2002 , 32, 299-315	6.9	122
67	¹ H, ¹⁵ N and ¹³ C chemical shift assignments of the structured core of the pseudomonas effector protein AvrPto. <i>Journal of Biomolecular NMR</i> , 2002 , 23, 247-8	3	3
66	The tobacco salicylic acid-binding protein 3 (SABP3) is the chloroplast carbonic anhydrase, which exhibits antioxidant activity and plays a role in the hypersensitive defense response. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 11640-5	11.5	285
65	Genomewide identification of <i>Pseudomonas syringae</i> pv. tomato DC3000 promoters controlled by the HrpL alternative sigma factor. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 2275-80	11.5	256
64	Tomato transcription factors pti4, pti5, and pti6 activate defense responses when expressed in Arabidopsis. <i>Plant Cell</i> , 2002 , 14, 817-31	11.6	297
63	Deductions about the number, organization, and evolution of genes in the tomato genome based on analysis of a large expressed sequence tag collection and selective genomic sequencing. <i>Plant Cell</i> , 2002 , 14, 1441-56	11.6	259
62	Two distinct <i>Pseudomonas</i> effector proteins interact with the Pto kinase and activate plant immunity. <i>Cell</i> , 2002 , 109, 589-98	56.2	233
61	Innate immunity in plants. <i>Current Opinion in Immunology</i> , 2001 , 13, 55-62	7.8	109
60	Arabidopsis genome sequence as a tool for functional genomics in tomato. <i>Genome Biology</i> , 2001 , 2, REVIEWS1003	18.3	17
59	Ancient origin of pathogen recognition specificity conferred by the tomato disease resistance gene Pto. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2001 , 98, 2059-64	11.5	61
58	The major site of the pti1 kinase phosphorylated by the pto kinase is located in the activation domain and is required for pto-pti1 physical interaction. <i>FEBS Journal</i> , 2000 , 267, 171-8		31
57	Signal recognition and transduction mediated by the tomato Pto kinase: a paradigm of innate immunity in plants. <i>Microbes and Infection</i> , 2000 , 2, 1591-7	9.3	27
56	Thr38 and Ser198 are Pto autophosphorylation sites required for the AvrPto-Pto-mediated hypersensitive response. <i>EMBO Journal</i> , 2000 , 19, 2257-69	13	90
55	High-resolution linkage analysis and physical characterization of the EIX-responding locus in tomato. <i>Theoretical and Applied Genetics</i> , 2000 , 100, 184-189	6	34
54	AvrPto-dependent Pto-interacting proteins and AvrPto-interacting proteins in tomato. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000 , 97, 8836-40	11.5	108
53	The <i>Pseudomonas</i> AvrPto Protein Is Differentially Recognized by Tomato and Tobacco and Is Localized to the Plant Plasma Membrane. <i>Plant Cell</i> , 2000 , 12, 2323	11.6	1
52	Pti4 Is Induced by Ethylene and Salicylic Acid, and Its Product Is Phosphorylated by the Pto Kinase. <i>Plant Cell</i> , 2000 , 12, 771	11.6	7
51	Protein kinases in the plant defense response. <i>Advances in Botanical Research</i> , 2000 , 32, 379-404	2.2	13

50	The pseudomonas AvrPto protein is differentially recognized by tomato and tobacco and is localized to the plant plasma membrane. <i>Plant Cell</i> , 2000 , 12, 2323-2338	11.6	144
49	Pti4 is induced by ethylene and salicylic acid, and its product is phosphorylated by the Pto kinase. <i>Plant Cell</i> , 2000 , 12, 771-86	11.6	253
48	Overexpression of Pto activates defense responses and confers broad resistance. <i>Plant Cell</i> , 1999 , 11, 15-29	11.6	229
47	<i>Pseudomonas syringae</i> pv tomato induces the expression of tomato EREBP-like genes pti4 and pti5 independent of ethylene, salicylate and jasmonate. <i>Plant Journal</i> , 1999 , 20, 475-83	6.9	85
46	Rapid transcript accumulation of pathogenesis-related genes during an incompatible interaction in bacterial speck disease-resistant tomato plants. <i>Plant Molecular Biology</i> , 1999 , 40, 455-65	4.6	50
45	Functional analysis of plant disease resistance genes and their downstream effectors. <i>Current Opinion in Plant Biology</i> , 1999 , 2, 273-9	9.9	148
44	Overexpression of Pto Activates Defense Responses and Confers Broad Resistance. <i>Plant Cell</i> , 1999 , 11, 15	11.6	7
43	Biochemical characterization of the Pto protein kinase conferring speck disease resistance in tomato. <i>Current Plant Science and Biotechnology in Agriculture</i> , 1999 , 541-544		
42	Gene discovery for crop improvement. <i>Current Opinion in Biotechnology</i> , 1998 , 9, 220-6	11.4	17
41	Recognition specificity for the bacterial avirulence protein AvrPto is determined by Thr-204 in the activation loop of the tomato Pto kinase. <i>Molecular Cell</i> , 1998 , 2, 241-5	17.6	111
40	Molecular mechanisms involved in bacterial speck disease resistance of tomato. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 1998 , 353, 1455-1461	5.8	16
39	A nitrilase-like protein interacts with GCC box DNA-binding proteins involved in ethylene and defense responses. <i>Plant Physiology</i> , 1998 , 118, 867-74	6.6	46
38	Biochemical properties of two protein kinases involved in disease resistance signaling in tomato. <i>Journal of Biological Chemistry</i> , 1998 , 273, 15860-5	5.4	34
37	The myristylation motif of Pto is not required for disease resistance. <i>Molecular Plant-Microbe Interactions</i> , 1998 , 11, 572-6	3.6	36
36	Alleles of Pto and Fen occur in bacterial speck-susceptible and fenthion-insensitive tomato cultivars and encode active protein kinases. <i>Plant Cell</i> , 1997 , 9, 61-73	11.6	67
35	Alleles of Pto and Fen Occur in Bacterial Speck-Susceptible and Fenthion-Insensitive Tomato Cultivars and Encode Active Protein Kinases. <i>Plant Cell</i> , 1997 , 9, 61	11.6	10
34	The Pto kinase conferring resistance to tomato bacterial speck disease interacts with proteins that bind a cis-element of pathogenesis-related genes. <i>EMBO Journal</i> , 1997 , 16, 3207-18	13	374
33	Initiation of Plant Disease Resistance by Physical Interaction of AvrPto and Pto Kinase. <i>Science</i> , 1996 , 274, 2060-3	33.3	575

32	The Pto kinase mediates a signaling pathway leading to the oxidative burst in tomato. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996 , 93, 13393-7	11.5	74
31	Molecular Cloning of Plant Disease Resistance Genes 1996 , 1-32		4
30	The Pto bacterial resistance gene and the Fen insecticide sensitivity gene encode functional protein kinases with serine/threonine specificity. <i>Plant Physiology</i> , 1995 , 108, 1735-9	6.6	88
29	Expression of the Tomato Pto Gene in Tobacco Enhances Resistance to <i>Pseudomonas syringae</i> pv <i>tabaci</i> Expressing <i>avrPto</i> . <i>Plant Cell</i> , 1995 , 7, 1529-1536	11.6	115
28	The tomato gene <i>Pti1</i> encodes a serine/threonine kinase that is phosphorylated by Pto and is involved in the hypersensitive response. <i>Cell</i> , 1995 , 83, 925-35	56.2	346
27	Expression of the Tomato Pto Gene in Tobacco Enhances Resistance to <i>Pseudomonas syringae</i> pv <i>tabaci</i> Expressing <i>avrPto</i> . <i>Plant Cell</i> , 1995 , 7, 1529	11.6	21
26	The disease-resistance gene <i>Pto</i> and the fenthion-sensitivity gene <i>fen</i> encode closely related functional protein kinases. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995 , 92, 4181-4	11.5	50
25	Chromosome landing: a paradigm for map-based gene cloning in plants with large genomes. <i>Trends in Genetics</i> , 1995 , 11, 63-8	8.5	350
24	Molecular genetic analysis of the ripening-inhibitor and non-ripening loci of tomato: a first step in genetic map-based cloning of fruit ripening genes. <i>Molecular Genetics and Genomics</i> , 1995 , 248, 195-206		43
23	A member of the tomato Pto gene family confers sensitivity to fenthion resulting in rapid cell death. <i>Plant Cell</i> , 1994 , 6, 1543-52	11.6	176
22	Map-based cloning in crop plants: tomato as a model system II. Isolation and characterization of a set of overlapping yeast artificial chromosomes encompassing the jointless locus. <i>Molecular Genetics and Genomics</i> , 1994 , 244, 613-21		27
21	Chromosome 2-specific DNA clones from flow-sorted chromosomes of tomato. <i>Molecular Genetics and Genomics</i> , 1994 , 242, 551-8		63
20	Analysis of the molecular basis of <i>Pseudomonas syringae</i> pv. tomato resistance in tomato. <i>Euphytica</i> , 1994 , 79, 187-193	2.1	3
19	A Member of the Tomato Pto Gene Family Confers Sensitivity to Fenthion Resulting in Rapid Cell Death. <i>Plant Cell</i> , 1994 , 6, 1543	11.6	17
18	Construction of plant yeast artificial chromosome libraries 1994 , 383-399		
17	Map-based cloning of a protein kinase gene conferring disease resistance in tomato. <i>Science</i> , 1993 , 262, 1432-6	33.3	1242
16	Pre-germination genotypic screening using PCR amplification of half-seeds. <i>Theoretical and Applied Genetics</i> , 1993 , 86, 694-8	6	86
15	High-Resolution Linkage Analysis and Physical Characterization of the Pto Bacterial Resistance Locus in Tomato. <i>Molecular Plant-Microbe Interactions</i> , 1993 , 6, 26	3.6	56

14	Construction of a yeast artificial chromosome library of tomato and identification of cloned segments linked to two disease resistance loci. <i>Molecular Genetics and Genomics</i> , 1992 , 233, 25-32		101
13	High density molecular linkage maps of the tomato and potato genomes. <i>Genetics</i> , 1992 , 132, 1141-60	4	1160
12	Rapid identification of markers linked to a <i>Pseudomonas</i> resistance gene in tomato by using random primers and near-isogenic lines. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991 , 88, 2336-40	11.5	383
11	<i>Bradyrhizobium japonicum</i> ntrBC/glnAandnifA/glnAMutants: Further Evidence that Separate Regulatory Pathways Govern glnII Expression in Free-Living and Symbiotic Cells. <i>Molecular Plant-Microbe Interactions</i> , 1991 , 4, 254	3.6	4
10	<i>Bradyrhizobium japonicum</i> glnB, a putative nitrogen-regulatory gene, is regulated by NtrC at tandem promoters. <i>Journal of Bacteriology</i> , 1989 , 171, 5638-45	3.5	56
9	Role of the <i>Bradyrhizobium japonicum</i> ntrC gene product in differential regulation of the glutamine synthetase II gene (glnII). <i>Journal of Bacteriology</i> , 1988 , 170, 5452-9	3.5	57
8	Differential transcription of the two glutamine synthetase genes of <i>Bradyrhizobium japonicum</i> . <i>Journal of Bacteriology</i> , 1987 , 169, 5861-6	3.5	57
7	Landraces of <i>Phaseolus vulgaris</i> (Fabaceae) in Northern Malawi. I. Regional variation. <i>Economic Botany</i> , 1987 , 41, 190-203	1.7	41
6	Landraces of <i>Phaseolus vulgaris</i> (Fabaceae) in Northern Malawi. II. Generation and maintenance of variability. <i>Economic Botany</i> , 1987 , 41, 204-215	1.7	34
5	Ptr1 evolved convergently with RPS2 and Mr5 to mediate recognition of AvrRpt2 in diverse solanaceous species		2
4	Generation and molecular characterization of CRISPR/Cas9-induced mutations in 63 immunity-associated genes in tomato reveals specificity and a range of gene modifications		1
3	A <i>Solanum lycopersicoides</i> reference genome facilitates biological discovery in tomato		3
2	Genome of <i>Solanum pimpinellifolium</i> provides insights into structural variants during tomato breeding		1
1	Spelling changes and fluorescent tagging with prime editing vectors for plants		6