

Martin G Schmid

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7207518/martin-g-schmid-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

82

papers

3,252

citations

33

h-index

55

g-index

83

ext. papers

3,481

ext. citations

3.3

avg, IF

5.58

L-index

#	Paper	IF	Citations
82	Natural product isolation--how to get from biological material to pure compounds. <i>Natural Product Reports</i> , 2013 , 30, 525-45	15.1	232
81	Chiral separation principles in capillary electrophoresis. <i>Journal of Chromatography A</i> , 1997 , 792, 179-225	4.5	222
80	Chiral separation by capillary electromigration techniques. <i>Journal of Chromatography A</i> , 2008 , 1204, 140-56	4.5	212
79	Chiral separation by chromatographic and electromigration techniques. A review. <i>Biopharmaceutics and Drug Disposition</i> , 2001 , 22, 291-336	1.7	185
78	Recent progress in chiral separation principles in capillary electrophoresis. <i>Electrophoresis</i> , 2000 , 21, 4112-35	3.6	162
77	Recent advances in chiral separation principles in capillary electrophoresis and capillary electrochromatography. <i>Electrophoresis</i> , 2004 , 25, 3981-96	3.6	109
76	Advances in chiral separation using capillary electromigration techniques. <i>Electrophoresis</i> , 2007 , 28, 114-36	3.6	95
75	Chiral separation of amino acids by ligand-exchange capillary electrochromatography using continuous beds. <i>Electrophoresis</i> , 2000 , 21, 3141-4	3.6	93
74	A new easy-to-prepare homogeneous continuous electrochromatographic bed for enantiomer recognition. <i>Electrophoresis</i> , 2000 , 21, 3116-25	3.6	86
73	Chiral separation principles in chromatographic and electromigration techniques. <i>Molecular Biotechnology</i> , 2006 , 32, 159-80	3	80
72	Chiral separation of new cathinone- and amphetamine-related designer drugs by gas chromatography-mass spectrometry using trifluoroacetyl-L-prolyl chloride as chiral derivatization reagent. <i>Journal of Chromatography A</i> , 2012 , 1269, 352-9	4.5	69
71	Enantioseparation by chromatographic and electromigration techniques using ligand-exchange as chiral separation principle. <i>Analytical and Bioanalytical Chemistry</i> , 2011 , 400, 2305-16	4.4	67
70	Capillary zone electrophoretic separation of the enantiomers of dipeptides based on host-guest complexation with a chiral crown ether. <i>Journal of Chromatography A</i> , 1995 , 709, 81-88	4.5	64
69	Application of ligand-exchange capillary electrophoresis to the chiral separation of alpha-hydroxy acids and beta-blockers. <i>Journal of Chromatography A</i> , 2000 , 875, 307-14	4.5	61
68	Chiral ligand-exchange capillary electrophoresis. <i>Journal of Proteomics</i> , 2001 , 48, 143-54		61
67	Enantioseparation of hydroxy acids on easy-to-prepare continuous beds for capillary electrochromatography. <i>Electrophoresis</i> , 2001 , 22, 2616-9	3.6	58
66	Chiral separation of cathinone derivatives used as recreational drugs by cyclodextrin-modified capillary electrophoresis. <i>Electrophoresis</i> , 2012 , 33, 1624-30	3.6	57

65	Chiral separation of alpha-amino acids by ligand-exchange capillary electrophoresis using N-(2-hydroxy-octyl)-L-4-hydroxyproline as a selector. <i>Electrophoresis</i> , 1998 , 19, 2109-12	3.6	55
64	Chiral separations of cathinone and amphetamine-derivatives: Comparative study between capillary electrochromatography, supercritical fluid chromatography and three liquid chromatographic modes. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016 , 121, 232-243	3.5	53
63	Chiral separation of amino acids and glyceryl dipeptides by chiral ligand-exchange capillary electrophoresis comparing Cu(II), Co(II), Ni(II) and Zn(II) complexes of three different sugar acids. <i>Journal of Chromatography A</i> , 2008 , 1204, 210-8	4.5	47
62	Enantiomeric separation of new cathinone derivatives designer drugs by capillary electrochromatography using a chiral stationary phase, based on amylose tris(5-chloro-2-methylphenylcarbamate). <i>Electrophoresis</i> , 2014 , 35, 3242-9	3.6	44
61	Chiral separation of cathinone and amphetamine derivatives by HPLC/UV using sulfated β -cyclodextrin as chiral mobile phase additive. <i>Chirality</i> , 2014 , 26, 411-8	2.1	44
60	Chiral separation of cathinone derivatives used as recreational drugs by HPLC-UV using a CHIRALPAK \square AS-H column as stationary phase. <i>Chirality</i> , 2012 , 24, 486-92	2.1	41
59	New particle-loaded monoliths for chiral capillary electrochromatographic separation. <i>Electrophoresis</i> , 2004 , 25, 3195-203	3.6	39
58	Capillary electrophoretic chiral resolution of vicinal diols by complexation with borate and cyclodextrin: Comparative studies on different cyclodextrin derivatives. <i>Chirality</i> , 1997 , 9, 153-156	2.1	37
57	Fast chiral separation by ligand-exchange HPLC using a dynamically coated monolithic column. <i>Journal of Separation Science</i> , 2006 , 29, 1470-5	3.4	37
56	Determination of the relative percentage distribution of THCA and (Δ^9)-THC in herbal cannabis seized in Austria - Impact of different storage temperatures on stability. <i>Forensic Science International</i> , 2015 , 254, 167-71	2.6	35
55	Analysis and characterization of the novel psychoactive drug 4-chloromethcathinone (clephedrone). <i>Forensic Science International</i> , 2014 , 244, e56-9	2.6	35
54	Enantioseparation by ligand-exchange using particle-loaded monoliths: capillary-LC versus capillary electrochromatography. <i>Journal of Proteomics</i> , 2007 , 70, 77-85		34
53	Chiral separation of sympathomimetics by ligand exchange capillary electrophoresis. <i>Electrophoresis</i> , 1999 , 20, 2458-61	3.6	34
52	Cellulose tris-(3,5-dimethylphenylcarbamate)-based chiral stationary phase for the enantioseparation of drugs in supercritical fluid chromatography: comparison with HPLC. <i>Journal of Separation Science</i> , 2018 , 41, 1471-1478	3.4	33
51	Enantioseparation of benzofurans and other novel psychoactive compounds by CE and sulfobutylether β -cyclodextrin as chiral selector added to the BGE. <i>Electrophoresis</i> , 2014 , 35, 2793-9	3.6	33
50	Chiral separation of sympathomimetics and beta-blockers by ligand-exchange CE using Cu(II) complexes of L-tartaric acid and L-threonine as chiral selectors. <i>Electrophoresis</i> , 2007 , 28, 2675-82	3.6	33
49	Enantioseparation of dipeptides by capillary electrochromatography on a teicoplanin aglycone chiral stationary phase. <i>Journal of Chromatography A</i> , 2003 , 990, 83-90	4.5	32
48	Enantioseparation of amino acids and drugs by CEC, pressure supported CEC, and micro-hPLC using a teicoplanin aglycone stationary phase. <i>Journal of Separation Science</i> , 2002 , 25, 1297-1302	3.4	31

47	Indirect chiral separation of new recreational drugs by gas chromatography-mass spectrometry using trifluoroacetyl-L-prolyl chloride as chiral derivatization reagent. <i>Chirality</i> , 2015 , 27, 211-5	2.1	30
46	A Metabolomic Approach Applied to a Liquid Chromatography Coupled to High-Resolution Tandem Mass Spectrometry Method (HPLC-ESI-HRMS/MS): Towards the Comprehensive Evaluation of the Chemical Composition of Cannabis Medicinal Extracts. <i>Phytochemical Analysis</i> , 2018 , 29, 144-155	3.4	30
45	Enantioselective sequential-injection chemiluminescence immunoassays for 3,3',5-triiodothyronine (T3) and thyroxine (T4). <i>Analytica Chimica Acta</i> , 2002 , 463, 5-14	6.6	29
44	Enantioseparation of amino acids, alpha-hydroxy acids, and dipeptides by ligand-exchange CEC using silica-based chiral stationary phases. <i>Electrophoresis</i> , 2009 , 30, 2897-904	3.6	26
43	Indirect chiral separation of 8 novel amphetamine derivatives as potential new psychoactive compounds by GC-MS and HPLC. <i>Science and Justice - Journal of the Forensic Science Society</i> , 2017 , 57, 6-12	2	25
42	Stereoselective interaction of drug enantiomers with human serum transferrin in capillary zone electrophoresis (II). <i>Electrophoresis</i> , 1998 , 19, 282-7	3.6	25
41	Enantioselective separation of biologically active basic compounds in ultra-performance supercritical fluid chromatography. <i>Analytica Chimica Acta</i> , 2016 , 932, 98-105	6.6	25
40	Development of an enantioseparation method for novel psychoactive drugs by HPLC using a Lux Cellulose-2 column in polar organic phase mode. <i>Forensic Science International</i> , 2017 , 270, 232-240	2.6	24
39	Development of enantioselective chemiluminescence flow- and sequential-injection immunoassays for alpha-amino acids. <i>Journal of Proteomics</i> , 2002 , 53, 1-14		24
38	Separation of enantiomers and positional isomers of novel psychoactive substances in solid samples by chromatographic and electrophoretic techniques - A selective review. <i>Journal of Chromatography A</i> , 2020 , 1624, 461256	4.5	23
37	Enantioseparation of glyceryl-dipeptides by CEC using particle-loaded monoliths prepared by ring-opening metathesis polymerization (ROMP). <i>Journal of Proteomics</i> , 2006 , 69, 67-77		22
36	Chiral resolution of diastereomeric di- and tripeptides on a teicoplanin aglycone phase by capillary electrochromatography. <i>Electrophoresis</i> , 2003 , 24, 2543-9	3.6	21
35	Enantiorecognition of triiodothyronine and thyroxine enantiomers using different chiral selectors by HPLC and micro-HPLC. <i>Journal of Proteomics</i> , 2008 , 70, 1254-60		20
34	Influence of structure and chirality of the selector on the chiral recognition of amino acids using ligand-exchange capillary electrophoresis. <i>Electrophoresis</i> , 2002 , 23, 3006-12	3.6	19
33	Analysis of a new drug of abuse: cathinone derivative 1-(3,4-dimethoxyphenyl)-2-(ethylamino)pentan-1-one. <i>Journal of Separation Science</i> , 2015 , 38, 825-8	3.4	18
32	Enantioseparation of dansyl amino acids by HPLC on a monolithic column dynamically coated with a vancomycin derivative. <i>Biomedical Chromatography</i> , 2010 , 24, 1213-9	1.7	18
31	COMPARATIVE STUDY OF THE CHIRAL RESOLUTION OF β BLOCKERS ON CELLULOSE TRIS (3,5-DIMETHYL-PHENYL-CARBAMATE) PHASES IN NORMAL AND REVERSED PHASE MODES. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2001 , 24, 2493-2504	1.3	18
30	Separation of enantiomers of new psychoactive substances by high-performance liquid chromatography. <i>Journal of Separation Science</i> , 2018 , 41, 1274-1286	3.4	17

29	Role of the charge in continuous beds in the chiral separation of hydroxy acids by ligand-exchange capillary electrochromatography. <i>Electrophoresis</i> , 2003 , 24, 2983-5	3.6	16
28	Enantioseparation of dipeptides and tripeptides by micro-HPLC comparing teicoplanin and teicoplanin aglycone as chiral selectors. <i>Journal of Proteomics</i> , 2004 , 61, 1-10		16
27	Chiral separation of halogenated amino acids by ligand-exchange capillary electrophoresis. <i>Electrophoresis</i> , 2005 , 26, 3878-83	3.6	15
26	Chiral separation of cathinone derivatives using β -cyclodextrin-assisted capillary electrophoresis-Comparison of four different β -cyclodextrin derivatives used as chiral selectors. <i>Electrophoresis</i> , 2019 , 40, 1787-1794	3.6	14
25	Chiral Resolution of Dipeptides by Ligand Exchange Chromatography on Chemically Bonded Chiral Phases. <i>Journal of Liquid Chromatography and Related Technologies</i> , 1996 , 19, 2933-2942	1.3	13
24	Enantioselective potential of teicoplanin- and vancomycin-based superficially porous particles-packed columns for supercritical fluid chromatography. <i>Journal of Chromatography A</i> , 2020 , 1612, 460687	4.5	13
23	Enantiomeric separation of Novel Psychoactive Substances by capillary electrophoresis using (+)-18-crown-6-tetracarboxylic acid as chiral selector. <i>Chirality</i> , 2018 , 30, 1019	2.1	12
22	Enantioseparation of amino acids and alpha-hydroxy acids on ligand-exchange continuous beds by capillary electrochromatography. <i>Electrophoresis</i> , 2010 , 31, 1517-20	3.6	12
21	Determination of metabolites of 5-hydroxymethylfurfural in human urine after oral application. <i>Journal of Separation Science</i> , 2013 , 36, 670-6	3.4	10
20	Determination of the chiral status of different novel psychoactive substance classes by capillary electrophoresis and β -cyclodextrin derivatives. <i>Chirality</i> , 2020 , 32, 1191-1207	2.1	10
19	Enantioseparation performance of superficially porous particle vancomycin-based chiral stationary phases in supercritical fluid chromatography and high performance liquid chromatography; applicability for psychoactive substances. <i>Journal of Chromatography A</i> , 2021 , 1637, 461846	4.5	10
18	Development and validation of an HPLC method to determine metabolites of 5-hydroxymethylfurfural (5-HMF). <i>Journal of Separation Science</i> , 2012 , 35, 2567-74	3.4	9
17	Micro-HPLC-UV analysis of cocaine and its adulterants in illicit cocaine samples seized by Austrian police from 2012 to 2017. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2018 , 41, 6-13	1.3	8
16	Enantioselective high-performance liquid chromatography of therapeutically relevant aminoalcohols as their fluorescent 1-naphthyl isocyanate derivatives. <i>Biomedical Chromatography</i> , 2001 , 15, 212-6	1.7	8
15	Successful use of a novel Lux [®] i-Amylose-1 chiral column for enantioseparation of "legal highs" by HPLC. <i>Chirality</i> , 2020 , 32, 42-52	2.1	8
14	A Simple HPLC-UV Approach for Rapid Enantioseparation of Cathinones, Pyrovalerones and Other Novel Psychoactive Substances on a 2.5- μ m Cellulose Tris-(3,5-dimethylphenyl-carbamate) Column. <i>Chromatographia</i> , 2020 , 83, 321-329	2.1	7
13	Chiral separation of the β -sympathomimetic fenoterol by HPLC and capillary zone electrophoresis for pharmacokinetic studies. <i>Biomedical Chromatography</i> , 2010 , 24, 1125-9	1.7	7
12	Enantioselective separation of Novel Psychoactive Substances using a Lux [®] AMP 3 β h column and HPLC-UV. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2020 , 179, 112967	3.5	7

11	Determination of cathinone and cathine in Khat plant material by LC-MS/MS: Fresh vs. dried leaves. <i>Forensic Science International</i> , 2021 , 319, 110658	2.6	6
10	Chiral Resolution of RR, SS-Hydrobenzoin by Liquid Chromatography Using Borate-Cyclodextrin Complexation. <i>Journal of High Resolution Chromatography</i> , 1998 , 21, 414-416		5
9	Test purchase of new synthetic tryptamines via the Internet: Identity check by GC-MS and separation by HPLC. <i>Journal of Applied Pharmaceutical Science</i> , 028-034	2	4
8	Advantages and Pitfalls of Capillary Electrophoresis of Pharmaceutical Compounds and Their Enantiomers in Complex Samples: Comparison of Hydrodynamically Opened and Closed Systems. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	4
7	Chiral ligand-exchange capillary electrophoresis and capillary electrochromatography. <i>Methods in Molecular Biology</i> , 2004 , 243, 375-86	1.4	3
6	Use of chiral crown ethers in capillary electrophoresis. <i>Methods in Molecular Biology</i> , 2004 , 243, 317-21	1.4	3
5	Chiral Separation by Ligand Exchange 155-179		3
4	Application of chiral ligand-exchange stationary phases in capillary electrochromatography. <i>Methods in Molecular Biology</i> , 2013 , 970, 443-55	1.4	2
3	CHIRAL SEPARATION OF SEVERAL THIOBARBITURATES ON A CELLULOSE TRIS(4-METHYLBENZOATE) PHASE UNDER REVERSED-PHASE CONDITIONS. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2001 , 24, 69-77	1.3	2
2	Recent progress in chiral separation principles in capillary electrophoresis 2000 , 21, 4112		1
1	A simple and isocratic protein-based high performance liquid chromatography method for the enantioseparation of amphetamine derivatives. <i>Journal of Chromatography Open</i> , 2021 , 1, 100013		