

Jia-Rou Liu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7211610/publications.pdf>

Version: 2024-02-01

24
papers

307
citations

840119

11
h-index

887659

17
g-index

24
all docs

24
docs citations

24
times ranked

608
citing authors

#	ARTICLE	IF	CITATIONS
1	Effectiveness and Safety of Direct Oral Anticoagulants in an Asian Population with Atrial Fibrillation Undergoing Dialysis: A Population-Based Cohort Study and Meta-Analysis. <i>Cardiovascular Drugs and Therapy</i> , 2021, 35, 975-986.	1.3	20
2	Oral Anticoagulants and Antiplatelet Agents in Patients With Atrial Fibrillation and Concomitant Critical Limb Ischemia: A Nationwide Cohort Study. <i>Canadian Journal of Cardiology</i> , 2021, 37, 113-121.	0.8	3
3	Efficacy and Safety of Ticagrelor vs. Clopidogrel in East Asian Patients with Acute Myocardial Infarction: A Nationwide Cohort Study. <i>Clinical Pharmacology and Therapeutics</i> , 2021, 109, 443-451.	2.3	6
4	Non-vitamin K antagonist oral anticoagulants and warfarin in atrial fibrillation patients with concomitant peripheral artery disease. <i>European Heart Journal - Cardiovascular Pharmacotherapy</i> , 2021, 7, 50-58.	1.4	18
5	Gonadotropin-releasing hormone antagonist associated with lower cardiovascular risk compared with gonadotropin-releasing hormone agonist in prostate cancer: A nationwide cohort and in vitro study. <i>Prostate</i> , 2021, 81, 902-912.	1.2	10
6	Clinical Outcomes in Atrial Fibrillation Patients With a History of Cancer Treated With Non-Vitamin K Antagonist Oral Anticoagulants. <i>Stroke</i> , 2021, 52, 3132-3141.	1.0	20
7	Outcomes after ticagrelor versus clopidogrel treatment in end-stage renal disease patients with acute myocardial infarction: a nationwide cohort study. <i>Scientific Reports</i> , 2021, 11, 20826.	1.6	4
8	Major adverse cardiovascular and limb events in patients with diabetes and concomitant peripheral artery disease treated with sodium glucose cotransporter 2 inhibitor versus dipeptidyl peptidase-4 inhibitor. <i>Cardiovascular Diabetology</i> , 2020, 19, 160.	2.7	20
9	Impact of bleeding during dual antiplatelet therapy in patients with coronary artery disease. <i>Scientific Reports</i> , 2020, 10, 21345.	1.6	3
10	Pioglitazone Is Associated with Lower Major Adverse Cardiovascular and Cerebrovascular Events than DPP4-Inhibitors in Diabetic Patients with End-Stage Renal Disease: A Taiwan Nationwide Cohort Study, 2006-2016. <i>Journal of Clinical Medicine</i> , 2020, 9, 3578.	1.0	3
11	Pioglitazone Reduces Mortality and Adverse Events in Patients With Type 2 Diabetes and With Advanced Chronic Kidney Disease: National Cohort Study. <i>Diabetes Care</i> , 2020, 43, e152-e153.	4.3	11
12	Effectiveness, safety, and major adverse limb events in atrial fibrillation patients with concomitant diabetes mellitus treated with non-vitamin K antagonist oral anticoagulants. <i>Cardiovascular Diabetology</i> , 2020, 19, 63.	2.7	26
13	Direct Oral Anticoagulants in Atrial Fibrillation Patients With Concomitant Hyperthyroidism. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, 2893-2904.	1.8	7
14	Association of maternal ethnicity and urbanicity on severe pediatric disease: a nationwide cohort study. <i>BMC Pediatrics</i> , 2019, 19, 514.	0.7	2
15	End-stage renal disease patients using angiotensin-converting enzyme inhibitors and angiotensin receptor blockers may reduce the risk of mortality: a Taiwanese Nationwide cohort study. <i>Internal Medicine Journal</i> , 2018, 48, 1123-1132.	0.5	4
16	Suicide death rates in patients with cardiovascular diseases - A 15-year nationwide cohort study in Taiwan. <i>Journal of Affective Disorders</i> , 2018, 238, 187-193.	2.0	20
17	Dipeptidyl peptidase-4 inhibitor decreases the risk of atrial fibrillation in patients with type 2 diabetes: a nationwide cohort study in Taiwan. <i>Cardiovascular Diabetology</i> , 2017, 16, 159.	2.7	44
18	A comparison between angiotensin converting enzyme inhibitors and angiotensin receptor blockers on end stage renal disease and major adverse cardiovascular events in diabetic patients: a population-based dynamic cohort study in Taiwan. <i>Cardiovascular Diabetology</i> , 2016, 15, 56.	2.7	13

#	ARTICLE	IF	CITATIONS
19	Dialysis Patients with Implanted Drug-Eluting Stents Have Lower Major Cardiac Events and Mortality than Those with Implanted Bare-Metal Stents: A Taiwanese Nationwide Cohort Study. PLoS ONE, 2016, 11, e0146343.	1.1	10
20	Young Male Patients with Atrial Fibrillation and CHA2DS2-VASc Score of 1 May Not Need Anticoagulants: A Nationwide Population-Based Study. PLoS ONE, 2016, 11, e0151485.	1.1	17
21	Simvastatin+ezetimibe combination therapy is associated with a lower rate of major adverse cardiac events in type 2 diabetics than high potency statins alone: A population-based dynamic cohort study. International Journal of Cardiology, 2015, 190, 20-25.	0.8	22
22	A Robust GWSS Method to Simultaneously Detect Rare and Common Variants for Complex Disease. PLoS ONE, 2015, 10, e0120873.	1.1	0
23	Hyperthyroid and Hypothyroid Status Was Strongly Associated with Gout and Weakly Associated with Hyperuricaemia. PLoS ONE, 2014, 9, e114579.	1.1	23
24	A Robust Rerank Approach for Feature Selection and Its Application to Pooling-Based GWA Studies. Computational and Mathematical Methods in Medicine, 2013, 2013, 1-11.	0.7	1