

Laurent Brard

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7212123/publications.pdf>

Version: 2024-02-01

54
papers

1,809
citations

318942

23
h-index

312153

41
g-index

56
all docs

56
docs citations

56
times ranked

3065
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Identification of Somatic Mitochondrial DNA Mutations, Heteroplasmy, and Increased Levels of Catenanes in Tumor Specimens Obtained from Three Endometrial Cancer Patients. <i>Life</i> , 2022, 12, 562. | 1.1 | 2 |
| 2 | Assessment of the diagnostic and prognostic relevance of ACAT1 and CE levels in plasma, peritoneal fluid and tumor tissue of epithelial ovarian cancer patients - a pilot study. <i>BMC Cancer</i> , 2022, 22, 387. | 1.1 | 6 |
| 3 | Assessment of peritoneal microbial features and tumor marker levels as potential diagnostic tools for ovarian cancer. <i>PLoS ONE</i> , 2020, 15, e0227707. | 1.1 | 28 |
| 4 | Assessment of acyl-CoA cholesterol acyltransferase (ACAT-1) role in ovarian cancer progression—An in vitro study. <i>PLoS ONE</i> , 2020, 15, e0228024. | 1.1 | 31 |
| 5 | Rural—urban differences in surgical treatment, regional lymph node examination, and survival in endometrial cancer patients. <i>Cancer Causes and Control</i> , 2018, 29, 221-232. | 0.8 | 13 |
| 6 | Utility and Generalizability of Multistate, Population-Based Cancer Registry Data for Rural Cancer Surveillance Research in the United States. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2018, 27, 1252-1260. | 1.1 | 18 |
| 7 | Rural—Urban Differences in Cancer Incidence and Trends in the United States. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2018, 27, 1265-1274. | 1.1 | 264 |
| 8 | Evaluation of the cytotoxicity of the Bithionol - cisplatin combination in a panel of human ovarian cancer cell lines. <i>BMC Cancer</i> , 2017, 17, 49. | 1.1 | 19 |
| 9 | Evaluation of the cytotoxicity of the Bithionol-paclitaxel combination in a panel of human ovarian cancer cell lines. <i>PLoS ONE</i> , 2017, 12, e0185111. | 1.1 | 9 |
| 10 | Assessment of the antitumor potential of Bithionol in vivo using a xenograft model of ovarian cancer. <i>Anti-Cancer Drugs</i> , 2016, 27, 547-559. | 0.7 | 6 |
| 11 | Expression Profiling of Primary and Metastatic Ovarian Tumors Reveals Differences Indicative of Aggressive Disease. <i>PLoS ONE</i> , 2014, 9, e94476. | 1.1 | 66 |
| 12 | Bithionol inhibits ovarian cancer cell growth In Vitro- studies on mechanism(s) of action. <i>BMC Cancer</i> , 2014, 14, 61. | 1.1 | 38 |
| 13 | PT19c, Another Nonhypercalcemic Vitamin D2 Derivative, Demonstrates Antitumor Efficacy in Epithelial Ovarian and Endometrial Cancer Models. <i>Genes and Cancer</i> , 2013, 4, 524-534. | 0.6 | 11 |
| 14 | Identification of Ovarian Cancer Metastatic miRNAs. <i>PLoS ONE</i> , 2013, 8, e58226. | 1.1 | 78 |
| 15 | Anti-angiogenic activity of cranberry proanthocyanidins and cytotoxic properties in ovarian cancer cells. <i>International Journal of Oncology</i> , 2012, 40, 227-35. | 1.4 | 34 |
| 16 | Purified cranberry proanthocyanidines (PAC-1A) cause pro-apoptotic signaling, ROS generation, cyclophosphamide retention and cytotoxicity in high-risk neuroblastoma cells. <i>International Journal of Oncology</i> , 2012, 40, 99-108. | 1.4 | 21 |
| 17 | WNT7A Regulates Tumor Growth and Progression in Ovarian Cancer through the WNT/ β -Catenin Pathway. <i>Molecular Cancer Research</i> , 2012, 10, 469-482. | 1.5 | 159 |
| 18 | Tetrathiomolybdate sensitizes ovarian cancer cells to anticancer drugs doxorubicin, fenretinide, 5-fluorouracil and mitomycin C. <i>BMC Cancer</i> , 2012, 12, 147. | 1.1 | 30 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Cytotoxic Properties of Adamantyl Isothiocyanate and Potential <i>In vivo</i> Metabolite Adamantyl-N-Acetylcystein in Gynecological Cancer Cells. <i>Chemical Biology and Drug Design</i> , 2012, 79, 92-103. | 1.5 | 2 |
| 20 | 7 Methyl indole ethyl isothiocyanate causes ROS mediated apoptosis and cell cycle arrest in endometrial cancer cells. <i>Gynecologic Oncology</i> , 2012, 126, 252-258. | 0.6 | 9 |
| 21 | Efficacy of a Non-Hypercalcemic Vitamin-D2 Derived Anti-Cancer Agent (MT19c) and Inhibition of Fatty Acid Synthesis in an Ovarian Cancer Xenograft Model. <i>PLoS ONE</i> , 2012, 7, e34443. | 1.1 | 16 |
| 22 | Antitumor activity of nifurtimox is enhanced with tetrathiomolybdate in medulloblastoma. <i>International Journal of Oncology</i> , 2011, 38, 1329-41. | 1.4 | 18 |
| 23 | Organometallic Iron(III)-Salophene Exerts Cytotoxic Properties in Neuroblastoma Cells via MAPK Activation and ROS Generation. <i>PLoS ONE</i> , 2011, 6, e19049. | 1.1 | 17 |
| 24 | A Phase 1 Study of Nifurtimox in Patients With Relapsed/Refractory Neuroblastoma. <i>Journal of Pediatric Hematology/Oncology</i> , 2011, 33, 25-30. | 0.3 | 41 |
| 25 | T090137 Inhibits Cisplatin-Induced Apoptosis in Ovarian Cancer Cells. <i>International Journal of Gynecological Cancer</i> , 2011, 21, 1350-1356. | 1.2 | 8 |
| 26 | Tetrathiomolybdate induces doxorubicin sensitivity in resistant tumor cell lines. <i>Gynecologic Oncology</i> , 2011, 122, 183-189. | 0.6 | 23 |
| 27 | Evaluation of the first Ergocalciferol-derived, non hypercalcemic anti-cancer agent MT19c in ovarian cancer SKOV-3 cell lines. <i>Gynecologic Oncology</i> , 2011, 123, 370-378. | 0.6 | 11 |
| 28 | A coumarin derivative (RKS262) inhibits cell-cycle progression, causes pro-apoptotic signaling and cytotoxicity in ovarian cancer cells. <i>Investigational New Drugs</i> , 2011, 29, 63-72. | 1.2 | 49 |
| 29 | Integrated genomics of ovarian xenograft tumor progression and chemotherapy response. <i>BMC Cancer</i> , 2011, 11, 308. | 1.1 | 10 |
| 30 | Oral RKS262 reduces tumor burden in a neuroblastoma xenograft animal model and mediates cytotoxicity through SAPK/JNK and ROS activation in vitro. <i>Cancer Biology and Therapy</i> , 2011, 11, 1036-1045. | 1.5 | 8 |
| 31 | Effect of a Vitamin D3 derivative (B3CD) with postulated anti-cancer activity in an ovarian cancer animal model. <i>Investigational New Drugs</i> , 2010, 28, 543-553. | 1.2 | 17 |
| 32 | Lipophilic aroylhydrazone chelator HNTMB and its multiple effects on ovarian cancer cells. <i>BMC Cancer</i> , 2010, 10, 72. | 1.1 | 13 |
| 33 | Chemotherapeutic Effect of Calcidiol Derivative B3CD in a Neuroblastoma Xenograft Model. <i>Chemical Biology and Drug Design</i> , 2010, 76, 164-173. | 1.5 | 7 |
| 34 | Apoptotic and chemotherapeutic properties of iron(III)-salophene in an ovarian cancer animal model. <i>Drug Design, Development and Therapy</i> , 2009, 3, 17. | 2.0 | 9 |
| 35 | Is There a Taxane-Free Interval That Predicts Response to Taxanes as a Later-Line Treatment of Recurrent Ovarian or Primary Peritoneal Cancer?. <i>International Journal of Gynecological Cancer</i> , 2009, 19, 343-347. | 1.2 | 4 |
| 36 | Cranberry proanthocyanidins are cytotoxic to human cancer cells and sensitize platinum-resistant ovarian cancer cells to paraplatin. <i>Phytotherapy Research</i> , 2009, 23, 1066-1074. | 2.8 | 56 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Nifurtimox Induces Apoptosis of Neuroblastoma Cells In Vitro and In Vivo. <i>Journal of Pediatric Hematology/Oncology</i> , 2009, 31, 187-193. | 0.3 | 36 |
| 38 | Induction of cytotoxicity, apoptosis and cell cycle arrest by 1-t-butyl carbamoyl, 7-methyl-indole-3-ethyl isothiocyanate (NB7M) in nervous system cancer cells. <i>Drug Design, Development and Therapy</i> , 2009, 2, 61-9. | 2.0 | 7 |
| 39 | Isothiocyanate NB7M causes selective cytotoxicity, pro-apoptotic signalling and cell-cycle regression in ovarian cancer cells. <i>British Journal of Cancer</i> , 2008, 99, 1823-1831. | 2.9 | 13 |
| 40 | Isolated sentinel lymph node dissection with conservative management in patients with squamous cell carcinoma of the vulva: A prospective trial. <i>Gynecologic Oncology</i> , 2008, 109, 65-70. | 0.6 | 62 |
| 41 | A novel indole ethyl isothiocyanate (7Me-IEITC) with anti-proliferative and pro-apoptotic effects on platinum-resistant human ovarian cancer cells. <i>Gynecologic Oncology</i> , 2008, 109, 240-249. | 0.6 | 19 |
| 42 | Synthesis of Bicyclic Aryl Thiazolines with Selective Anti-Proliferative Effects on Human Cancer Cell Lines. <i>Letters in Organic Chemistry</i> , 2008, 5, 103-109. | 0.2 | 3 |
| 43 | For Women Receiving Chemotherapy for Clinically Apparent Early Ovarian Cancer, Is There a Benefit to Surgical Staging?. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 2008, 31, 39-42. | 0.6 | 11 |
| 44 | Iron(III)-Salophene: An Organometallic Compound with Selective Cytotoxic and Anti-Proliferative Properties in Platinum-Resistant Ovarian Cancer Cells. <i>PLoS ONE</i> , 2008, 3, e2303. | 1.1 | 50 |
| 45 | Effect of indole ethyl isothiocyanates on proliferation, apoptosis, and MAPK signaling in neuroblastoma cell lines. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2007, 17, 5846-5852. | 1.0 | 27 |
| 46 | Anti-proliferative and Pro-apoptotic Properties of 3-Bromoacetoxy Calcidiol in High-risk Neuroblastoma. <i>Chemical Biology and Drug Design</i> , 2007, 70, 302-310. | 1.5 | 21 |
| 47 | Does the platinum-free interval predict the incidence or severity of hypersensitivity reactions to carboplatin? The experience from Women and Infants' Hospital. <i>Gynecologic Oncology</i> , 2007, 105, 81-83. | 0.6 | 55 |
| 48 | Inguinal Sentinel Node Dissection versus Standard Inguinal Node Dissection in Patients With Vulvar Cancer: A Comparison of the Size of Metastasis Detected in Inguinal Lymph Nodes. <i>Obstetrical and Gynecological Survey</i> , 2006, 61, 518-519. | 0.2 | 2 |
| 49 | Inhibition of angiogenesis by vitamin D-binding protein: Characterization of anti-endothelial activity of DBP-maf. <i>Angiogenesis</i> , 2006, 8, 349-360. | 3.7 | 36 |
| 50 | Iron chelators deferoxamine and diethylenetriamine pentaacetic acid induce apoptosis in ovarian carcinoma. <i>Gynecologic Oncology</i> , 2006, 100, 116-127. | 0.6 | 53 |
| 51 | Inguinal sentinel node dissection versus standard inguinal node dissection in patients with vulvar cancer: A comparison of the size of metastasis detected in inguinal lymph nodes. <i>Gynecologic Oncology</i> , 2006, 101, 24-27. | 0.6 | 27 |
| 52 | The effect of total parenteral nutrition on the survival of terminally ill ovarian cancer patients. <i>Gynecologic Oncology</i> , 2006, 103, 176-180. | 0.6 | 87 |
| 53 | Phenethyl isothiocyanate (PEITC) inhibits growth of ovarian cancer cells by inducing apoptosis: Role of caspase and MAPK activation. <i>Gynecologic Oncology</i> , 2006, 103, 261-270. | 0.6 | 115 |
| 54 | Benzyl isothiocyanate (BITC) induces apoptosis in ovarian cancer cells in vitro. <i>Journal of Experimental Therapeutics and Oncology</i> , 2006, 5, 287-300. | 0.5 | 34 |