

Mariana Araújo Vieira do Carmo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7214074/publications.pdf>

Version: 2024-02-01

35
papers

2,000
citations

304743

22
h-index

361022

35
g-index

35
all docs

35
docs citations

35
times ranked

2350
citing authors

#	ARTICLE	IF	CITATIONS
1	Purple tea (<i>Camellia sinensis</i> var. <i>assamica</i>) leaves as a potential functional ingredient: From extraction of phenolic compounds to cell-based antioxidant/biological activities. <i>Food and Chemical Toxicology</i> , 2022, 159, 112668.	3.6	9
2	Metabolomics, sensory evaluation, and enzymatic hydrolysis reveal the effect of storage on the critical astringency-active components of crude Pu-erh tea. <i>Journal of Food Composition and Analysis</i> , 2022, 107, 104387.	3.9	13
3	From the forest to the plate – Hemicelluloses, galactoglucomannan, glucuronoxylan, and phenolic-rich extracts from unconventional sources as functional food ingredients. <i>Food Chemistry</i> , 2022, 381, 132284.	8.2	19
4	Black tea kombucha: Physicochemical, microbiological and comprehensive phenolic profile changes during fermentation, and antimalarial activity. <i>Food Chemistry</i> , 2022, 384, 132515.	8.2	27
5	Polyphenols of jaboticaba [<i>Myrciaria jaboticaba</i> (Vell.) O.Berg] seeds incorporated in a yogurt model exert antioxidant activity and modulate gut microbiota of 1,2-dimethylhydrazine-induced colon cancer in rats. <i>Food Chemistry</i> , 2021, 334, 127565.	8.2	50
6	Effects of epigallocatechin gallate, epigallocatechin and epicatechin gallate on the chemical and cell-based antioxidant activity, sensory properties, and cytotoxicity of a catechin-free model beverage. <i>Food Chemistry</i> , 2021, 339, 128060.	8.2	64
7	Effects of microwave heating on the chemical composition and bioactivity of orange juice-milk beverages. <i>Food Chemistry</i> , 2021, 345, 128746.	8.2	28
8	Ellagitannins from jaboticaba (<i>Myrciaria jaboticaba</i>) seeds attenuated inflammation, oxidative stress, aberrant crypt foci, and modulated gut microbiota in rats with 1,2 dimethyl hydrazine-induced colon carcinogenesis. <i>Food and Chemical Toxicology</i> , 2021, 154, 112287.	3.6	13
9	Extraction optimization of bioactive compounds from ora-pro-nobis (<i>Pereskia aculeata</i> Miller) leaves and their in vitro antioxidant and antihemolytic activities. <i>Food Chemistry</i> , 2021, 361, 130078.	8.2	14
10	Selina-1,3,7(11)-trien-8-one and Oxidoselina-1,3,7(11)-trien-8-one from <i>Eugenia uniflora</i> Leaf Essential Oil and Their Cytotoxic Effects on Human Cell Lines. <i>Molecules</i> , 2021, 26, 740.	3.8	4
11	Antioxidant/pro-oxidant and antiproliferative activities of phenolic-rich foods and extracts: A cell-based point of view. <i>Advances in Food and Nutrition Research</i> , 2021, 98, 253-280.	3.0	12
12	Untargeted and targeted metabolomics reveal the chemical characteristic of pu-erh tea (<i>Camellia</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	8.2	77
13	From byproduct to a functional ingredient: Camu-camu (<i>Myrciaria dubia</i>) seed extract as an antioxidant agent in a yogurt model. <i>Journal of Dairy Science</i> , 2020, 103, 1131-1140.	3.4	44
14	Camu-camu seed (<i>Myrciaria dubia</i>) – From side stream to an antioxidant, antihyperglycemic, antiproliferative, antimicrobial, antihemolytic, anti-inflammatory, and antihypertensive ingredient. <i>Food Chemistry</i> , 2020, 310, 125909.	8.2	56
15	Optimizing the extraction of bioactive compounds from pu-erh tea (<i>Camellia sinensis</i> var. <i>assamica</i>) and evaluation of antioxidant, cytotoxic, antimicrobial, antihemolytic, and inhibition of α -amylase and α -glucosidase activities. <i>Food Research International</i> , 2020, 137, 109430.	6.2	26
16	Response surface optimization of phenolic compounds extraction from camu-camu (<i>Myrciaria</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5 2358-2367.	3.1	11
17	A new analytical concept based on chemistry and toxicology for herbal extracts analysis: From phenolic composition to bioactivity. <i>Food Research International</i> , 2020, 132, 109090.	6.2	23
18	Response surface optimization of phenolic compounds from jaboticaba (<i>Myrciaria cauliflora</i> [Mart.]) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5 assessments. <i>Food and Chemical Toxicology</i> , 2020, 142, 111439.	3.6	32

#	ARTICLE	IF	CITATIONS
19	Camu-camu (<i>Myrciaria dubia</i>) seeds as a novel source of bioactive compounds with promising antimalarial and antischistosomicidal properties. <i>Food Research International</i> , 2020, 136, 109334.	6.2	13
20	Hydroalcoholic <i>Myrciaria dubia</i> (camu-camu) seed extracts prevent chromosome damage and act as antioxidant and cytotoxic agents. <i>Food Research International</i> , 2019, 125, 108551.	6.2	24
21	Chemistry and Biological Activities of Processed <i>Camellia sinensis</i> Teas: A Comprehensive Review. <i>Comprehensive Reviews in Food Science and Food Safety</i> , 2019, 18, 1474-1495.	11.7	283
22	Antioxidants-rich ice cream containing herbal extracts and fructooligosaccharides: manufacture, functional and sensory properties. <i>Food Chemistry</i> , 2019, 298, 125098.	8.2	29
23	Multivariate effects of Chinese keemun black tea grades (<i>Camellia sinensis</i> var. <i>sinensis</i>) on the phenolic composition, antioxidant, antihemolytic and cytotoxic/cytoprotection activities. <i>Food Research International</i> , 2019, 125, 108516.	6.2	52
24	Effect of <i>Pereskia aculeata</i> Mill. in vitro and in overweight humans: A randomized controlled trial. <i>Journal of Food Biochemistry</i> , 2019, 43, e12903.	2.9	12
25	Red Chicory (<i>Cichorium intybus</i>) Extract Rich in Anthocyanins: Chemical Stability, Antioxidant Activity, and Antiproliferative Activity <i>In Vitro</i> . <i>Journal of Food Science</i> , 2019, 84, 990-1001.	3.1	39
26	Flaxleaf Fleabane Leaves (<i>Conyza bonariensis</i>), A New Functional Nonconventional Edible Plant?. <i>Journal of Food Science</i> , 2019, 84, 3473-3482.	3.1	13
27	<i>Sclerotinia Sclerotiorum</i> (White Mold): Cytotoxic, Mutagenic, and Antimalarial Effects <i>In Vivo</i> and <i>In Vitro</i> . <i>Journal of Food Science</i> , 2019, 84, 3866-3875.	3.1	10
28	Hibiscus <i>sabdariffa</i> anthocyanins-rich extract: Chemical stability, in vitro antioxidant and antiproliferative activities. <i>Food and Chemical Toxicology</i> , 2018, 113, 187-197.	3.6	92
29	Optimized <i>Camellia sinensis</i> var. <i>sinensis</i> , <i>Ilex paraguariensis</i> , and <i>Aspalathus linearis</i> blend presents high antioxidant and antiproliferative activities in a beverage model. <i>Food Chemistry</i> , 2018, 254, 348-358.	8.2	58
30	Chemical study, antioxidant, anti-hypertensive, and cytotoxic/cytoprotective activities of <i>Centaurea cyanus</i> L. petals aqueous extract. <i>Food and Chemical Toxicology</i> , 2018, 118, 439-453.	3.6	68
31	In vitro antioxidant and antihypertensive compounds from camu-camu (<i>Myrciaria dubia</i> McVaugh), <i>Tj ETQq1 1 0.784314 rgBT /Overl</i> 479-490.	3.6	64
32	High-throughput assay comparison and standardization for metal chelating capacity screening: A proposal and application. <i>Food Chemistry</i> , 2017, 214, 515-522.	8.2	146
33	Comparison between Folin-Ciocalteu and Prussian Blue Assays to Estimate The Total Phenolic Content of Juices and Teas Using 96-Well Microplates. <i>Journal of Food Science</i> , 2015, 80, C2397-403.	3.1	132
34	Observations on the use of statistical methods in Food Science and Technology. <i>Food Research International</i> , 2014, 55, 137-149.	6.2	392
35	Analytical Strategy Coupled with Response Surface Methodology To Maximize the Extraction of Antioxidants from Ternary Mixtures of Green, Yellow, and Red Teas (<i>Camellia sinensis</i> var.) <i>Tj ETQq1 1 0.784314 rgBT /Overl</i>	3.1	10