

# Kyu Hun Choi

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/7226807/kyu-hun-choi-publications-by-citations.pdf>

**Version:** 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

134  
papers

2,498  
citations

26  
h-index

43  
g-index

138  
ext. papers

3,022  
ext. citations

3.8  
avg, IF

4.2  
L-index

#	Paper	IF	Citations
134	Serum C-Reactive Protein: A Predictor of Mortality in Continuous Ambulatory Peritoneal Dialysis Patients. <i>Peritoneal Dialysis International</i> , <b>1998</b> , 18, 387-394	2.8	128
133	Changing prevalence of glomerular diseases in Korean adults: a review of 20 years of experience. <i>Nephrology Dialysis Transplantation</i> , <b>2009</b> , 24, 2406-10	4.3	119
132	Changes in Causative Organisms and Their Antimicrobial Susceptibilities in Capd Peritonitis: A Single Center's Experience over one Decade. <i>Peritoneal Dialysis International</i> , <b>2004</b> , 24, 424-432	2.8	111
131	Red blood cell distribution width is an independent predictor of mortality in acute kidney injury patients treated with continuous renal replacement therapy. <i>Nephrology Dialysis Transplantation</i> , <b>2012</b> , 27, 589-94	4.3	104
130	KNOW-CKD (KoreaN cohort study for Outcome in patients With Chronic Kidney Disease): design and methods. <i>BMC Nephrology</i> , <b>2014</b> , 15, 80	2.7	102
129	Decreased circulating C3 levels and mesangial C3 deposition predict renal outcome in patients with IgA nephropathy. <i>PLoS ONE</i> , <b>2012</b> , 7, e40495	3.7	73
128	Reduced residual renal function is a risk of peritonitis in continuous ambulatory peritoneal dialysis patients. <i>Nephrology Dialysis Transplantation</i> , <b>2007</b> , 22, 2653-8	4.3	72
127	Obesity, Metabolic Abnormality, and Progression of CKD. <i>American Journal of Kidney Diseases</i> , <b>2018</b> , 72, 400-410	7.4	67
126	Clinical features and outcomes of IgA nephropathy with nephrotic syndrome. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , <b>2012</b> , 7, 427-36	6.9	67
125	Improving Outcome of Capd: Twenty-Five Years Experience in a Single Korean Center. <i>Peritoneal Dialysis International</i> , <b>2007</b> , 27, 432-440	2.8	61
124	An analysis of 4,514 cases of renal biopsy in Korea. <i>Yonsei Medical Journal</i> , <b>2001</b> , 42, 247-54	3	61
123	Plasma levels of soluble receptor for advanced glycation end products (sRAGE) and proinflammatory ligand for RAGE (EN-RAGE) are associated with carotid atherosclerosis in patients with peritoneal dialysis. <i>Atherosclerosis</i> , <b>2012</b> , 220, 208-14	3.1	50
122	Live donor renal allograft in end-stage renal failure patients from immunoglobulin A nephropathy. <i>Transplantation</i> , <b>2001</b> , 71, 233-8	1.8	46
121	Effect of losartan and amlodipine on proteinuria and transforming growth factor-beta1 in patients with IgA nephropathy. <i>Nephrology Dialysis Transplantation</i> , <b>2003</b> , 18, 1115-21	4.3	45
120	Using the Oxford classification of IgA nephropathy to predict long-term outcomes of Henoch-Schönlein purpura nephritis in adults. <i>Modern Pathology</i> , <b>2014</b> , 27, 972-82	9.8	44
119	Left atrial volume is an independent predictor of mortality in CAPD patients. <i>Nephrology Dialysis Transplantation</i> , <b>2011</b> , 26, 3732-9	4.3	41
118	Sclerosing encapsulating peritonitis as a complication of long-term continuous ambulatory peritoneal dialysis in Korea. <i>Nephrology</i> , <b>2003</b> , 8 Suppl, S33-9	2.2	40

117	Clinical outcomes, when matched at presentation, do not vary between adult-onset Henñh-Schñlein purpura nephritis and IgA nephropathy. <i>Kidney International</i> , <b>2012</b> , 82, 1304-12	9.9	36
116	High glucose activates the p38 MAPK pathway in cultured human peritoneal mesothelial cells. <i>Kidney International</i> , <b>2003</b> , 63, 958-68	9.9	31
115	Association of inflammation and protein-energy wasting with endothelial dysfunction in peritoneal dialysis patients. <i>Nephrology Dialysis Transplantation</i> , <b>2010</b> , 25, 1266-71	4.3	29
114	Influence of ketoanalog supplementation on the progression in chronic kidney disease patients who had training on low-protein diet. <i>Nephrology</i> , <b>2009</b> , 14, 750-7	2.2	29
113	Clinical characteristics of dialysis related sclerosing encapsulating peritonitis: multi-center experience in Korea. <i>Yonsei Medical Journal</i> , <b>2005</b> , 46, 104-11	3	29
112	ACE gene polymorphism and progression of diabetic nephropathy in Korean type 2 diabetic patients: effect of ACE gene DD on the progression of diabetic nephropathy. <i>American Journal of Kidney Diseases</i> , <b>2003</b> , 41, 943-9	7.4	28
111	Early initiation of continuous renal replacement therapy improves patient survival in severe progressive septic acute kidney injury. <i>Journal of Critical Care</i> , <b>2012</b> , 27, 743.e9-18	4	27
110	Cancer in patients on chronic dialysis in Korea. <i>Journal of Korean Medical Science</i> , <b>2009</b> , 24 Suppl, S95-S101	1.7	27
109	Urinary Potassium Excretion and Progression of CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , <b>2019</b> , 14, 330-340	6.9	26
108	Comparison of the Haas and the Oxford classifications for prediction of renal outcome in patients with IgA nephropathy. <i>Human Pathology</i> , <b>2014</b> , 45, 236-43	3.7	26
107	Progression of aortic arch calcification over 1 year is an independent predictor of mortality in incident peritoneal dialysis patients. <i>PLoS ONE</i> , <b>2012</b> , 7, e48793	3.7	26
106	Ability of nafamostat mesilate to prolong filter patency during continuous renal replacement therapy in patients at high risk of bleeding: a randomized controlled study. <i>PLoS ONE</i> , <b>2014</b> , 9, e108737	3.7	25
105	Hyponatremia as a predictor of mortality in peritoneal dialysis patients. <i>PLoS ONE</i> , <b>2014</b> , 9, e111373	3.7	25
104	Good glycemic control is associated with better survival in diabetic patients on peritoneal dialysis: a prospective observational study. <i>PLoS ONE</i> , <b>2012</b> , 7, e30072	3.7	24
103	Clinical implication of crescentic lesions in immunoglobulin A nephropathy. <i>Nephrology Dialysis Transplantation</i> , <b>2014</b> , 29, 356-64	4.3	23
102	Circulating Fibroblast Growth Factor-23 Levels are Associated with an Increased Risk of Anemia Development in Patients with Nondialysis Chronic Kidney Disease. <i>Scientific Reports</i> , <b>2018</b> , 8, 7294	4.9	22
101	Visceral fat thickness is associated with carotid atherosclerosis in peritoneal dialysis patients. <i>Obesity</i> , <b>2012</b> , 20, 1301-7	8	22
100	Combined vascular effects of HMG-CoA reductase inhibitor and angiotensin receptor blocker in non-diabetic patients undergoing peritoneal dialysis. <i>Nephrology Dialysis Transplantation</i> , <b>2011</b> , 26, 3722-8	4.3	22

99	Usefulness of 23S rRNA amplification by PCR in the detection of bacteria in CAPD peritonitis. <i>American Journal of Nephrology</i> , <b>2006</b> , 26, 115-20	4.6	22
98	Diastolic dysfunction is associated with an increased risk of contrast-induced nephropathy: a retrospective cohort study. <i>BMC Nephrology</i> , <b>2013</b> , 14, 146	2.7	20
97	The prevalence and associated risk factors of renal artery stenosis in patients undergoing cardiac catheterization. <i>Yonsei Medical Journal</i> , <b>2000</b> , 41, 219-25	3	20
96	Optimal proteinuria target for renoprotection in patients with IgA nephropathy. <i>PLoS ONE</i> , <b>2014</b> , 9, e101935	3.7	19
95	Statin therapy is associated with improved survival in incident peritoneal dialysis patients: propensity-matched comparison. <i>Nephrology Dialysis Transplantation</i> , <b>2011</b> , 26, 4090-4	4.3	19
94	Risk Factors for Adverse Outcomes after Peritonitis-Related Technique Failure. <i>Peritoneal Dialysis International</i> , <b>2008</b> , 28, 352-360	2.8	19
93	Factors associated with Hypokalemia in Continuous Ambulatory Peritoneal Dialysis Patients. <i>Electrolyte and Blood Pressure</i> , <b>2007</b> , 5, 102-10	1.1	19
92	Association of serum adiponectin level with albuminuria in chronic kidney disease patients. <i>Clinical and Experimental Nephrology</i> , <b>2016</b> , 20, 443-9	2.5	17
91	The impact of pretransplant 25-hydroxy vitamin D deficiency on subsequent graft function: an observational study. <i>BMC Nephrology</i> , <b>2012</b> , 13, 22	2.7	17
90	Elevated osteoprotegerin is associated with inflammation, malnutrition and new onset cardiovascular events in peritoneal dialysis patients. <i>Atherosclerosis</i> , <b>2011</b> , 219, 925-30	3.1	17
89	Gentamicin induced apoptosis of renal tubular epithelial (LLC-PK1) cells. <i>Korean Journal of Internal Medicine</i> , <b>2000</b> , 15, 218-23	2.5	17
88	Prognostic value of elevated cardiac troponin I in ESRD patients with sepsis. <i>Nephrology Dialysis Transplantation</i> , <b>2009</b> , 24, 1568-73	4.3	16
87	Glomerular crescents are responsible for chronic graft dysfunction in post-transplant IgA nephropathy. <i>Pathology International</i> , <b>2004</b> , 54, 837-42	1.8	16
86	Electrocardiographic left ventricular hypertrophy and outcome in hemodialysis patients. <i>PLoS ONE</i> , <b>2012</b> , 7, e35534	3.7	16
85	Smoking, Smoking Cessation, and Progression of Chronic Kidney Disease: Results From KNOW-CKD Study. <i>Nicotine and Tobacco Research</i> , <b>2021</b> , 23, 92-98	4.9	16
84	Dietary Protein Intake, Protein Energy Wasting, and the Progression of Chronic Kidney Disease: Analysis from the KNOW-CKD Study. <i>Nutrients</i> , <b>2019</b> , 11,	6.7	15
83	Long-term outcome of kidney transplantation in adult recipients with focal segmental glomerulosclerosis. <i>Yonsei Medical Journal</i> , <b>2001</b> , 42, 209-14	3	15
82	Association Between Serum High-Density Lipoprotein Cholesterol Levels and Progression of Chronic Kidney Disease: Results From the KNOW-CKD. <i>Journal of the American Heart Association</i> , <b>2019</b> , 8, e011162	6	14

81	The effect of specialized continuous renal replacement therapy team in acute kidney injury patients treatment. <i>Yonsei Medical Journal</i> , <b>2015</b> , 56, 658-65	3	14
80	Elevated cardiac troponin T predicts cardiovascular events in asymptomatic continuous ambulatory peritoneal dialysis patients without a history of cardiovascular disease. <i>American Journal of Nephrology</i> , <b>2009</b> , 29, 129-35	4.6	14
79	Baseline Cardiovascular Characteristics of Adult Patients with Chronic Kidney Disease from the KoreaN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>Journal of Korean Medical Science</i> , <b>2017</b> , 32, 231-239	4.7	13
78	A low serum bicarbonate concentration as a risk factor for mortality in peritoneal dialysis patients. <i>PLoS ONE</i> , <b>2013</b> , 8, e82912	3.7	13
77	Association of Serum Osteoprotegerin Levels with Bone Loss in Chronic Kidney Disease: Insights from the KNOW-CKD Study. <i>PLoS ONE</i> , <b>2016</b> , 11, e0166792	3.7	13
76	Association between health related quality of life and progression of chronic kidney disease. <i>Scientific Reports</i> , <b>2019</b> , 9, 19595	4.9	13
75	Normal body mass index with central obesity has increased risk of coronary artery calcification in Korean patients with chronic kidney disease. <i>Kidney International</i> , <b>2016</b> , 90, 1368-1376	9.9	12
74	Low circulating adiponectin levels are associated with insulin resistance in non-obese peritoneal dialysis patients. <i>Endocrine Journal</i> , <b>2012</b> , 59, 685-95	2.9	12
73	Spontaneous remission of IgA nephropathy associated with resolution of hepatitis A. <i>American Journal of Kidney Diseases</i> , <b>2010</b> , 56, 1163-7	7.4	12
72	Chlamydia pneumoniae accompanied by inflammation is associated with the progression of atherosclerosis in CAPD patients: a prospective study for 3 years. <i>Nephrology Dialysis Transplantation</i> , <b>2008</b> , 23, 1011-8	4.3	12
71	Clinical features and outcomes of focal segmental glomerulosclerosis pathologic variants in Korean adult patients. <i>BMC Nephrology</i> , <b>2014</b> , 15, 52	2.7	11
70	Predictors of mortality in patients returning to dialysis after allograft loss. <i>Blood Purification</i> , <b>2010</b> , 30, 56-63	3.1	11
69	Hyponatremia Predicts New-Onset Cardiovascular Events in Peritoneal Dialysis Patients. <i>PLoS ONE</i> , <b>2015</b> , 10, e0129480	3.7	11
68	Removal of kidney stones by extracorporeal shock wave lithotripsy is associated with delayed progression of chronic kidney disease. <i>Yonsei Medical Journal</i> , <b>2012</b> , 53, 708-14	3	10
67	Attenuation of nephritis in lupus-prone mice by thalidomide. <i>Rheumatology</i> , <b>2012</b> , 51, 2131-40	3.9	10
66	Segmental glomerulosclerosis in IgA nephropathy after renal transplantation: relationship with proteinuria and therapeutic response to enalapril. <i>Clinical Transplantation</i> , <b>2003</b> , 17, 108-13	3.8	10
65	Studies on bone markers and bone mineral density in patients with chronic renal failure. <i>Yonsei Medical Journal</i> , <b>1996</b> , 37, 350-6	3	10
64	Glomerular epithelial CD44 expression and segmental sclerosis in IgA nephropathy. <i>Clinical and Experimental Nephrology</i> , <b>2016</b> , 20, 871-877	2.5	9

63	The effect of renin-angiotensin system blockade on renal protection in chronic kidney disease patients with hyperkalemia. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , <b>2014</b> , 15, 491-7	9	9
62	Left atrial enlargement is associated with a rapid decline in residual renal function in ESRD patients on peritoneal dialysis. <i>Journal of the American Society of Echocardiography</i> , <b>2012</b> , 25, 421-7	5.8	9
61	Immunoglobulin A nephropathy associated with Plasmodium falciparum malaria. <i>Journal of Korean Medical Science</i> , <b>2012</b> , 27, 446-9	4.7	9
60	Estimating average glucose levels from glycated albumin in patients with end-stage renal disease. <i>Yonsei Medical Journal</i> , <b>2012</b> , 53, 578-86	3	9
59	Impact of cyclosporin on podocyte ZO-1 expression in puromycin aminonucleoside nephrosis rats. <i>Yonsei Medical Journal</i> , <b>2005</b> , 46, 141-8	3	9
58	Alcohol Consumption and Progression of Chronic Kidney Disease: Results From the Korean Cohort Study for Outcome in Patients with Chronic Kidney Disease. <i>Mayo Clinic Proceedings</i> , <b>2020</b> , 95, 293-305	6.4	9
57	Hypoadiponectinemia and the presence of metabolic syndrome in patients with chronic kidney disease: results from the KNOW-CKD study. <i>Diabetology and Metabolic Syndrome</i> , <b>2016</b> , 8, 75	5.6	9
56	Comparison of two creatinine-based equations for predicting decline in renal function in type 2 diabetic patients with nephropathy in a korean population. <i>International Journal of Endocrinology</i> , <b>2013</b> , 2013, 848963	2.7	8
55	Two distinct functional patterns of hepatitis C Virus (HCV)-specific T cell responses in seronegative, aviremic patients. <i>PLoS ONE</i> , <b>2013</b> , 8, e62319	3.7	8
54	High peritoneal transport status is not an independent risk factor for high mortality in patients treated with automated peritoneal dialysis. <i>Journal of Korean Medical Science</i> , <b>2010</b> , 25, 1313-7	4.7	8
53	Antiproteinuric effect of losartan in non-diabetic renal disease is not dependent on ACE insertion/deletion polymorphism. <i>Kidney and Blood Pressure Research</i> , <b>2006</b> , 29, 216-24	3.1	8
52	The effects of high glucose concentration on angiotensin II- or transforming growth factor-beta-induced DNA synthesis, hypertrophy and collagen synthesis in cultured rat mesangial cells. <i>Yonsei Medical Journal</i> , <b>1996</b> , 37, 302-11	3	8
51	Low Serum Bicarbonate Predicts Residual Renal Function Loss in Peritoneal Dialysis Patients. <i>Medicine (United States)</i> , <b>2015</b> , 94, e1276	1.8	7
50	Stepwise treatment using corticosteroids alone and in combination with cyclosporine in korean patients with idiopathic membranous nephropathy. <i>Yonsei Medical Journal</i> , <b>2013</b> , 54, 973-82	3	7
49	Relation of homocysteinemia to contrast-induced nephropathy in patients undergoing percutaneous coronary intervention. <i>American Journal of Cardiology</i> , <b>2011</b> , 108, 1086-91	3	7
48	Sagittal abdominal diameter is an independent predictor of all-cause and cardiovascular mortality in incident peritoneal dialysis patients. <i>PLoS ONE</i> , <b>2013</b> , 8, e77082	3.7	7
47	Albuminuria as a Risk Factor for Anemia in Chronic Kidney Disease: Result from the KoreaN Cohort Study for Outcomes in Patients With Chronic Kidney Disease (KNOW-CKD). <i>PLoS ONE</i> , <b>2015</b> , 10, e0139747	2.7	7
46	High fibroblast growth factor 23 is associated with coronary calcification in patients with high adiponectin: analysis from the KoreaN cohort study for Outcome in patients With Chronic Kidney Disease (KNOW-CKD) study. <i>Nephrology Dialysis Transplantation</i> , <b>2019</b> , 34, 123-129	4.3	7



45	Association between the ratio of insulin-like growth factor-I to insulin-like growth factor binding protein-3 and inflammation in incident automated peritoneal dialysis patients. <i>Growth Hormone and IGF Research</i> , <b>2013</b> , 23, 170-4	2	6
44	The effect of lovastatin on proliferation of cultured rat mesangial and aortic smooth muscle cells. <i>Yonsei Medical Journal</i> , <b>1995</b> , 36, 251-61	3	6
43	Association of Blood Pressure With the Progression of CKD: Findings From KNOW-CKD Study. <i>American Journal of Kidney Diseases</i> , <b>2021</b> , 78, 236-245	7.4	6
42	Urinary angiotensinogen level is associated with potassium homeostasis and clinical outcome in patients with polycystic kidney disease: a prospective cohort study. <i>BMC Nephrology</i> , <b>2019</b> , 20, 104	2.7	5
41	High serum adiponectin is associated with anemia development in chronic kidney disease: The results from the KNOW-CKD study. <i>Cytokine</i> , <b>2018</b> , 103, 1-9	4	5
40	Association between Urine Creatinine Excretion and Arterial Stiffness in Chronic Kidney Disease: Data from the KNOW-CKD Study. <i>Kidney and Blood Pressure Research</i> , <b>2016</b> , 41, 527-534	3.1	5
39	Urine Osmolality and Renal Outcome in Patients with Chronic Kidney Disease: Results from the KNOW-CKD. <i>Kidney and Blood Pressure Research</i> , <b>2019</b> , 44, 1089-1100	3.1	5
38	Membranous glomerulonephritis in a patient with myelodysplastic syndrome-refractory cytopenia with multilineage dysplasia. <i>Kidney Research and Clinical Practice</i> , <b>2013</b> , 32, 134-7	3.6	5
37	Modified Peritoneal Dialysis Catheter Insertion: Comparison with a Conventional Method. <i>Yonsei Medical Journal</i> , <b>2015</b> , 56, 981-6	3	5
36	Idiopathic retroperitoneal fibrosis associated with Hashimoto's thyroiditis in an old-aged man. <i>Yonsei Medical Journal</i> , <b>2008</b> , 49, 1032-5	3	5
35	Hepatitis G virus infection in hemodialysis and continuous ambulatory peritoneal dialysis patients. <i>Yonsei Medical Journal</i> , <b>1998</b> , 39, 116-21	3	5
34	A novel multiplex real-time PCR assay for the concurrent detection of hepatitis A, B and C viruses in patients with acute hepatitis. <i>PLoS ONE</i> , <b>2012</b> , 7, e49106	3.7	5
33	eGFR and coronary artery calcification in chronic kidney disease. <i>European Journal of Clinical Investigation</i> , <b>2019</b> , 49, e13101	4.6	4
32	The relationship between hemodialysis modality and insulin resistance in non-diabetic hemodialysis patients. <i>Blood Purification</i> , <b>2015</b> , 39, 224-229	3.1	4
31	Effect of peritoneal dialysis modality on the 1-year rate of decline of residual renal function. <i>Yonsei Medical Journal</i> , <b>2014</b> , 55, 141-8	3	4
30	Rapid decrease of intact parathyroid hormone could be a predictor of better response to cinacalcet in hemodialysis patients. <i>Yonsei Medical Journal</i> , <b>2013</b> , 54, 453-63	3	4
29	Influence of nutritional status on CAPD peritonitis. <i>Yonsei Medical Journal</i> , <b>1990</b> , 31, 65-70	3	4
28	Relationship between Cardiac Geometry and Serum Hepcidin in Chronic Kidney Disease: Analysis from the KNOW-CKD Study. <i>Journal of Korean Medical Science</i> , <b>2020</b> , 35, e2	4.7	4

27	The Correlation of Serum Osteoprotegerin with Non-Traditional Cardiovascular Risk Factors and Arterial Stiffness in Patients with Pre-Dialysis Chronic Kidney Disease: Results from the KNOW-CKD Study. <i>Journal of Korean Medical Science</i> , <b>2018</b> , 33, e322	4.7	4
26	A case of membranous nephropathy as a manifestation of graft-versus-host disease. <i>Kidney Research and Clinical Practice</i> , <b>2013</b> , 32, 39-42	3.6	3
25	Intensity of statin therapy and renal outcome in chronic kidney disease: Results from the Korean Cohort Study for Outcome in Patients With Chronic Kidney Disease. <i>Kidney Research and Clinical Practice</i> , <b>2020</b> , 39, 93-102	3.6	3
24	Smoking-Related Renal Histologic Injury in IgA Nephropathy Patients. <i>Yonsei Medical Journal</i> , <b>2016</b> , 57, 209-16	3	3
23	The effect of interactions between proteinuria, activity of fibroblast growth factor 23 and serum phosphate on renal progression in patients with chronic kidney disease: a result from the Korean cohort study for Outcome in patients With Chronic Kidney Disease study. <i>Nephrology Dialysis Transplantation</i> , <b>2020</b> , 35, 438-446	4.3	3
22	Urinary chloride concentration and progression of chronic kidney disease: results from the Korean cohort study for Outcomes in patients With Chronic Kidney Disease. <i>Nephrology Dialysis Transplantation</i> , <b>2021</b> , 36, 673-680	4.3	3
21	Cystatin C is Better than Serum Creatinine for Estimating Glomerular Filtration Rate to Detect Osteopenia in Chronic Kidney Disease Patients. <i>Yonsei Medical Journal</i> , <b>2017</b> , 58, 380-387	3	2
20	Renal histologic parameters influencing postoperative renal function in renal cell carcinoma patients. <i>Korean Journal of Pathology</i> , <b>2013</b> , 47, 557-62		2
19	Correlates of the severity of coronary atherosclerosis in long-term kidney transplant patients. <i>Journal of Korean Medical Science</i> , <b>2010</b> , 25, 706-11	4.7	2
18	Low bone mineral density is associated with coronary arterial calcification progression and incident cardiovascular events in patients with chronic kidney disease.. <i>CKJ: Clinical Kidney Journal</i> , <b>2022</b> , 15, 1194-127	4.5	2
17	Association of Body Weight Variability with Adverse Cardiovascular Outcomes in Patients with Pre-Dialysis Chronic Kidney Disease. <i>Nutrients</i> , <b>2021</b> , 13,	6.7	2
16	Minimal Change Disease in Systemic Lupus: Another Renal Manifestation of Lupus?. <i>The Ewha Medical Journal</i> , <b>2013</b> , 36, 139	0.1	1
15	Effect of baseline glomerular filtration rate on renal function following stenting for atherosclerotic renal artery stenosis. <i>Scandinavian Journal of Urology and Nephrology</i> , <b>2010</b> , 44, 169-74		1
14	Protective effect of urinary alkalinization on gentamicin nephrotoxicity in rats. <i>Yonsei Medical Journal</i> , <b>1988</b> , 29, 225-32	3	1
13	Serum uric acid is associated with coronary artery calcification in early chronic kidney disease: a cross-sectional study. <i>BMC Nephrology</i> , <b>2021</b> , 22, 247	2.7	1
12	The difference between cystatin C- and creatinine-based eGFR is associated with adverse cardiovascular outcome in patients with chronic kidney disease. <i>Atherosclerosis</i> , <b>2021</b> , 335, 53-61	3.1	1
11	Smoking Cessation and Coronary Artery Calcification in CKD. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , <b>2021</b> , 16, 870-879	6.9	0
10	Incidence of cardiovascular events and mortality in Korean patients with chronic kidney disease. <i>Scientific Reports</i> , <b>2021</b> , 11, 1131	4.9	0



9	Association of Left Ventricular Diastolic Dysfunction With Cardiovascular Outcomes in Patients With Pre-dialysis Chronic Kidney Disease: Findings From KNOW-CKD Study.. <i>Frontiers in Cardiovascular Medicine</i> , <b>2022</b> , 9, 844312	5.4	o
8	Balloon catheter looping technique for entry site angioplasty during endovascular management of thrombosed arteriovenous grafts by single access. <i>CardioVascular and Interventional Radiology</i> , <b>2014</b> , 37, 502-7	2.7	
7	The effect of anti-hypertensive drugs on DNA synthesis and proliferation of cultured rat aortic smooth muscle cells. <i>Yonsei Medical Journal</i> , <b>1997</b> , 38, 160-6	3	
6	A Case of Membranous Glomerulonephritis Coexisting with Focal Segmental Glomerulosclerosis. <i>Ewha Medical Journal</i> , <b>2006</b> , 29, 133		
5	Interleukin-1 beta, -6 and interferon-gamma productions in patients undergoing continuous ambulatory peritoneal dialysis. <i>Yonsei Medical Journal</i> , <b>1998</b> , 39, 240-6	3	
4	Effect of captopril on heavy proteinuria in patients with various glomerular diseases. <i>Yonsei Medical Journal</i> , <b>1992</b> , 33, 232-9	3	
3	A Clinical Study on the Anti-hypertensive Effect of Barnidipine HCl in Hypertensive Patients with Renal Disease. <i>Journal of the Korean Society for Clinical Pharmacology and Therapeutics</i> , <b>1996</b> , 4, 185		
2	A Case of Continuous Ambulatory Peritoneal Dialysis-Associated Peritonitis Caused by Mycobacterium abscessus. <i>Korean Journal of Medicine</i> , <b>2014</b> , 87, 110	0.5	
1	Association between the transtubular potassium gradient and progression of chronic kidney disease: results from KNOW-CKD. <i>Journal of Nephrology</i> , <b>2021</b> , 34, 2063-2072	4.8	