Halit Filiz

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7227127/publications.pdf

Version: 2024-02-01

1307594 839539 30 409 7 18 citations g-index h-index papers 30 30 30 399 citing authors docs citations times ranked all docs

#	Article	IF	CITATIONS
1	The Vertebrate Biodiversity of Turkey. , 2021, , 175-274.		13
2	A global-scale screening of non-native aquatic organisms to identify potentially invasive species under current and future climate conditions. Science of the Total Environment, 2021, 788, 147868.	8.0	80
3	Risk of invasiveness of non-native aquatic species in the eastern Mediterranean region under current and projected climate conditions., 2021, 88, 1130-1143.		12
4	First occurrence of a lesser amberjack Seriola fasciata (Bloch, 1793) in the Aegean coasts of Turkey with morphological and molecular identification. Regional Studies in Marine Science, 2020, 40, 101494.	0.7	2
5	Year-Round Aggregation of Sandbar Shark, <i>Carcharhinus plumbeus</i> (Nardo, 1827), in Boncuk Cove in the southern Aegean Sea, Turkey (Carcharhiniformes: Carcharhinidae). Zoology in the Middle East, 2019, 65, 35-39.	0.6	5
6	Biological aspects of two coexisting native and non-native fish species in the Aegean Sea: Pagellus erythrinus vs. Nemipterus randalli. Mediterranean Marine Science, 2019, 20, 594.	1.6	9
7	A Risk Screening Study on the Potential Invasiveness of Lessepsian fishes in the South-Eastern Coasts of Anatolia. Acta Ichthyologica Et Piscatoria, 2019, 49, 23-31.	0.7	30
8	Occurrence of <i>Lobotes surinamensis</i> (Osteichthyes: Lobotidae) in the Mediterranean: Historical and recent data. Zoology in the Middle East, 2017, 63, 43-47.	0.6	6
9	The Factors Increasing of Invasiveness Potential of Five Pufferfishes in the Eastern Mediterranean, Turkey. Natural and Engineering Sciences, 2017, 2, 22-30.	0.3	5
10	Length-Weight Relationships of Four Lessepsian Puffer Fish Specıes From Muğla Coasts of Turkey. Natural and Engineering Sciences, 2017, 2, 36-40.	0.3	4
11	Champsodontids at the gates: first record of <i>Champsodon vorax</i> Günther, 1867 from the Aegean Sea (Teleostei: Champsodontidae). Journal of Applied Ichthyology, 2016, 32, 120-122.	0.7	4
12	New sighting of the endangered species blackchin guitarfish, Rhinobatos cemiculus, in the South Aegean Sea. Marine Biodiversity Records, 2016, 9, .	1.2	2
13	Filling the gap: first confirmed record for the Ophisurus serpens (Anguilliformes: Ophichthidae) from the Anatolian coast of the South Aegean Sea. Marine Biodiversity Records, 2015, 8, .	1.2	4
14	First length-weight relationships of 11 fish species in the Aegean Sea. Journal of Applied Ichthyology, 2015, 31, 398-402.	0.7	8
15	Occurrence, length-weight and length-length relationships of <i>Champsodon nudivittis</i> (Ogilby,) Tj ETQq1 1	0.78431	4 rgBT /Overlo
16	A new record of the Blue-Barred Parrotfish, <i>Scarus ghobban </i> (Actinopterygii: Scaridae), from the coastal waters of Cyprus. Zoology in the Middle East, 2014, 60, 281-282.	0.6	1
17	Weight–length relations for 103 fish species from the southern Aegean Sea, Turkey. Acta Ichthyologica Et Piscatoria, 2014, 44, 263-269.	0.7	46
18	Marine Litter: Composition in Eastern Aegean Coasts. Open Access Library Journal (oalib), 2014, 01, 1-7.	0.2	4

#	Article	IF	CITATIONS
19	Diet composition of the five deep sea fish from the Aegean Sea Su Ürünleri Dergisi, 2013, 30, 61-67.	0.3	2
20	Are introduced gibel carp <i>Carassius gibelio</i> in Turkey more invasive in artificial than in natural waters?. Fisheries Management and Ecology, 2012, 19, 178-187.	2.0	50
21	Title is missing!. Turkish Journal of Fisheries and Aquatic Sciences, 2011, 11, .	0.9	4
22	The Occurrence of the Cadenat's Rockfish, <l>Scorpaena Loppei</l> (Actinopterygii:) Tj ETQq0 0 0 rgBT 75-78.	/Overlock 0.7	10 Tf 50 6
23	Age and growth, reproduction and diet of the black goby, (Gobius niger) from Aegean Sea, Turkey. Journal of Fisheriessciencescom, 2009, , .	0.2	8
24	First record of <l>Epigonus constanciae</l> (Giglioli, 1880), (deepwater cardinalfish) (Osteichthyes: Epigonidae), from Aegean Sea, Turkey. Acta Ichthyologica Et Piscatoria, 2007, 37, 91-94.	0.7	4
25	New records of <l>Bregmaceros atlanticus</l> (Bregmacerotidae), <l>Echiodon dentatus</l> (Carapidae), and <l>Nemichthys scolopaceus</l> (Nemichthyidae) from the Aegean Sea. Acta Ichthyologica Et Piscatoria, 2007, 37, 107-112.	0.7	12
26	Age and growth of the hollowsnout grenadier, Caelorinchus caelorhincus (Risso, 1810), in the Aegean Sea. Journal of Applied Ichthyology, 2006, 22, 285-287.	0.7	8
27	First record of the Small-toothed Argentine, <i>Glossanodon leioglossus </i> (Valenciennes, 1848) (Osteichthyes: Argentinidae), from Turkey. Zoology in the Middle East, 2005, 34, 110-112.	0.6	2
28	Length-weight relationships of 24 fish species from the North Aegean Sea, Turkey. Journal of Applied Ichthyology, 2004, 20, 431-432.	0.7	65
29	Development of a Filament on the Lower Lobe of the Caudal Fin of the Lessepsian Migrant Fish Nemipterus randalli (Actinopterygii: Clupeiformes: Nemipteridae). Proceedings of the Zoological Society, 0, , 1.	1.0	1
30	A NEW MAXIMUM LENGTH FOR THE PARROTFISH, Sparisoma cretense (Linnaeus, 1758) IN THE MEDITERRANEAN SEA. Journal of Aquaculture Engineering and Fisheries Research, 0, , 140-143.	0.6	5