

# Ann Depicker

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/7231462/ann-depicker-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

89  
papers

6,686  
citations

36  
h-index

81  
g-index

90  
ext. papers

7,561  
ext. citations

7.4  
avg, IF

5.65  
L-index

#	Paper	IF	Citations
89	Evaluating single-domain antibodies as carriers for targeted vaccine delivery to the small intestinal epithelium. <i>Journal of Controlled Release</i> , <b>2020</b> , 321, 416-429	11.7	6
88	Seed-produced anti-globulin VHH-Fc antibodies retrieve globulin precursors in the insoluble fraction and modulate the <i>Arabidopsis thaliana</i> seed subcellular morphology. <i>Plant Molecular Biology</i> , <b>2020</b> , 103, 597-608	4.6	2
87	Simplified monomeric VHH-Fc antibodies provide new opportunities for passive immunization. <i>Current Opinion in Biotechnology</i> , <b>2020</b> , 61, 96-101	11.4	12
86	Transformation strategies for stable expression of complex hetero-multimeric proteins like secretory immunoglobulin A in plants. <i>Plant Biotechnology Journal</i> , <b>2019</b> , 17, 1760-1769	11.6	4
85	Yeast-secreted, dried and food-admixed monomeric IgA prevents gastrointestinal infection in a piglet model. <i>Nature Biotechnology</i> , <b>2019</b> , 37, 527-530	44.5	29
84	A two-amino acid mutation in murine IgA enables downstream processing and purification on staphylococcal superantigen-like protein 7. <i>Journal of Biotechnology</i> , <b>2019</b> , 294, 26-29	3.7	4
83	Russell-Like Bodies in Plant Seeds Share Common Features With Prolamin Bodies and Occur Upon Recombinant Protein Production. <i>Frontiers in Plant Science</i> , <b>2019</b> , 10, 777	6.2	6
82	Transgene Silencing <b>2018</b> , 1-32		
81	In planta expression of nanobody-based designer chicken antibodies targeting <i>Campylobacter</i> . <i>PLoS ONE</i> , <b>2018</b> , 13, e0204222	3.7	10
80	High accumulation in tobacco seeds of hemagglutinin antigen from avian (H5N1) influenza. <i>Transgenic Research</i> , <b>2017</b> , 26, 775-789	3.3	8
79	Recombinant IgA production for mucosal passive immunization, advancing beyond the hurdles. <i>Cellular and Molecular Life Sciences</i> , <b>2016</b> , 73, 535-45	10.3	21
78	The case for plant-made veterinary immunotherapeutics. <i>Biotechnology Advances</i> , <b>2016</b> , 34, 597-604	17.8	32
77	Biomanufacturing of protective antibodies and other therapeutics in edible plant tissues for oral applications. <i>Plant Biotechnology Journal</i> , <b>2016</b> , 14, 1791-9	11.6	24
76	Tobacco seeds as efficient production platform for a biologically active anti-HBsAg monoclonal antibody. <i>Transgenic Research</i> , <b>2015</b> , 24, 897-909	3.3	12
75	Using GlycoDelete to produce proteins lacking plant-specific N-glycan modification in seeds. <i>Nature Biotechnology</i> , <b>2015</b> , 33, 1135-7	44.5	36
74	Plant expression systems for early stage discovery and development of lead therapeutic antibodies. <i>Human Antibodies</i> , <b>2015</b> , 23, 37-43	1.3	5
73	Comparison of VHH-Fc antibody production in <i>Arabidopsis thaliana</i> , <i>Nicotiana benthamiana</i> and <i>Pichia pastoris</i> . <i>Plant Biotechnology Journal</i> , <b>2015</b> , 13, 938-47	11.6	19

72	Single-domain antibodies targeting neuraminidase protect against an H5N1 influenza virus challenge. <i>Journal of Virology</i> , <b>2014</b> , 88, 8278-96	6.6	44
71	Detection and investigation of transitive gene silencing in plants. <i>Methods in Molecular Biology</i> , <b>2014</b> , 1112, 219-41	1.4	
70	Trafficking of endoplasmic reticulum-retained recombinant proteins is unpredictable in <i>Arabidopsis thaliana</i> . <i>Frontiers in Plant Science</i> , <b>2014</b> , 5, 473	6.2	20
69	Generation of VHH antibodies against the <i>Arabidopsis thaliana</i> seed storage proteins. <i>Plant Molecular Biology</i> , <b>2014</b> , 84, 83-93	4.6	11
68	Nanobody-based products as research and diagnostic tools. <i>Trends in Biotechnology</i> , <b>2014</b> , 32, 263-70	15.1	262
67	Boosting in planta production of antigens derived from the porcine reproductive and respiratory syndrome virus (PRRSV) and subsequent evaluation of their immunogenicity. <i>PLoS ONE</i> , <b>2014</b> , 9, e91386	3.7	12
66	T-DNA transfer and T-DNA integration efficiencies upon <i>Arabidopsis thaliana</i> root explant cocultivation and floral dip transformation. <i>Planta</i> , <b>2013</b> , 238, 1025-37	4.7	6
65	Fusion of an Fc chain to a VHH boosts the accumulation levels in <i>Arabidopsis</i> seeds. <i>Plant Biotechnology Journal</i> , <b>2013</b> , 11, 1006-16	11.6	30
64	Recombinant antibody production in <i>Arabidopsis</i> seeds triggers an unfolded protein response. <i>Plant Physiology</i> , <b>2013</b> , 161, 1021-33	6.6	25
63	Site-specific T-DNA integration in <i>Arabidopsis thaliana</i> mediated by the combined action of CRE recombinase and $\gamma$ C31 integrase. <i>Plant Journal</i> , <b>2013</b> , 75, 172-184	6.9	13
62	The efficiency of <i>Arabidopsis thaliana</i> floral dip transformation is determined not only by the <i>Agrobacterium</i> strain used but also by the physiology and the ecotype of the dipped plant. <i>Molecular Plant-Microbe Interactions</i> , <b>2013</b> , 26, 823-32	3.6	12
61	Transitive RNA silencing signals induce cytosine methylation of a transgenic but not an endogenous target. <i>Plant Journal</i> , <b>2013</b> , 74, 867-79	6.9	17
60	Epigenetic switches of tobacco transgenes associate with transient redistribution of histone marks in callus culture. <i>Epigenetics</i> , <b>2013</b> , 8, 666-76	5.7	5
59	Orally fed seeds producing designer IgAs protect weaned piglets against enterotoxigenic <i>Escherichia coli</i> infection. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2013</b> , 110, 11809-14	11.5	82
58	Role of plant expression systems in antibody production for passive immunization. <i>International Journal of Developmental Biology</i> , <b>2013</b> , 57, 587-93	1.9	20
57	Production of camel-like antibodies in plants. <i>Methods in Molecular Biology</i> , <b>2012</b> , 911, 305-24	1.4	18
56	High frequency of single-copy T-DNA transformants produced after floral dip in CRE-expressing <i>Arabidopsis</i> plants. <i>Methods in Molecular Biology</i> , <b>2012</b> , 847, 317-33	1.4	2
55	Production of monoclonal antibodies with a controlled N-glycosylation pattern in seeds of <i>Arabidopsis thaliana</i> . <i>Plant Biotechnology Journal</i> , <b>2011</b> , 9, 179-92	11.6	43

54	Non-food/feed seeds as biofactories for the high-yield production of recombinant pharmaceuticals. <i>Plant Biotechnology Journal</i> , <b>2011</b> , 9, 911-21	11.6	38
53	Characterization of the single-chain Fv-Fc antibody MBP10 produced in Arabidopsis alg3 mutant seeds. <i>Transgenic Research</i> , <b>2011</b> , 20, 1033-42	3.3	9
52	Expression of antibody fragments with a controlled N-glycosylation pattern and induction of endoplasmic reticulum-derived vesicles in seeds of Arabidopsis. <i>Plant Physiology</i> , <b>2011</b> , 155, 2036-48	6.6	44
51	Paramutation of tobacco transgenes by small RNA-mediated transcriptional gene silencing. <i>Epigenetics</i> , <b>2011</b> , 6, 650-60	5.7	13
50	Introns reduce transitivity proportionally to their length, suggesting that silencing spreads along the pre-mRNA. <i>Plant Journal</i> , <b>2010</b> , 64, 392-401	6.9	23
49	Cell culture-induced gradual and frequent epigenetic reprogramming of invertedly repeated tobacco transgene epialleles. <i>Plant Physiology</i> , <b>2009</b> , 149, 1493-504	6.6	45
48	Production of antibody fragments in Arabidopsis seeds. <i>Methods in Molecular Biology</i> , <b>2009</b> , 483, 89-101	1.4	3
47	Evaluation of seven promoters to achieve germline directed Cre-lox recombination in Arabidopsis thaliana. <i>Plant Cell Reports</i> , <b>2009</b> , 28, 1509-20	5.1	17
46	High frequency of single-copy T-DNA transformants produced by floral dip in CRE-expressing Arabidopsis plants. <i>Plant Journal</i> , <b>2009</b> , 59, 517-27	6.9	25
45	The T-DNA integration pattern in Arabidopsis transformants is highly determined by the transformed target cell. <i>Plant Journal</i> , <b>2009</b> , 60, 134-45	6.9	57
44	Trans-generation inheritance of methylation patterns in a tobacco transgene following a post-transcriptional silencing event. <i>Plant Journal</i> , <b>2008</b> , 54, 1049-62	6.9	25
43	Agrobacterium Tumefaciens-Mediated Transformation: Patterns of T-Dna Integration Into the Host Genome <b>2008</b> , 441-481		11
42	Stability of the T-DNA flanking regions in transgenic Arabidopsis thaliana plants under influence of abiotic stress and cultivation practices. <i>Plant Cell Reports</i> , <b>2008</b> , 27, 749-57	5.1	8
41	The influence of matrix attachment regions on transgene expression in Arabidopsis thaliana wild type and gene silencing mutants. <i>Plant Molecular Biology</i> , <b>2007</b> , 63, 533-43	4.6	19
40	Aberrant localization and underglycosylation of highly accumulating single-chain Fv-Fc antibodies in transgenic Arabidopsis seeds. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2007</b> , 104, 1430-5	11.5	109
39	Recombinational cloning with plant gateway vectors. <i>Plant Physiology</i> , <b>2007</b> , 145, 1144-54	6.6	289
38	Generation of single-copy T-DNA transformants in Arabidopsis by the CRE/loxP recombination-mediated resolution system. <i>Plant Physiology</i> , <b>2007</b> , 145, 1171-82	6.6	36
37	The trans-silencing capacity of invertedly repeated transgenes depends on their epigenetic state in tobacco. <i>Nucleic Acids Research</i> , <b>2006</b> , 34, 2280-93	20.1	31

36	Introduction of silencing-inducing transgenes does not affect expression of known transcripts. <i>FEBS Letters</i> , <b>2006</b> , 580, 4154-9	3.8	6
35	Sequence stability of the T-DNA - plant junctions in tissue culture in Arabidopsis transgenic lines. <i>Plant Cell Reports</i> , <b>2006</b> , 25, 1362-8	5.1	6
34	Stable high-level transgene expression in Arabidopsis thaliana using gene silencing mutants and matrix attachment regions. <i>Plant Journal</i> , <b>2004</b> , 39, 440-9	6.9	117
33	A technology platform for the fast production of monoclonal recombinant antibodies against plant proteins and peptides. <i>Journal of Immunological Methods</i> , <b>2004</b> , 294, 181-7	2.5	13
32	Epigenetic switch from posttranscriptional to transcriptional silencing is correlated with promoter hypermethylation. <i>Plant Physiology</i> , <b>2003</b> , 133, 1240-50	6.6	70
31	Qualitative and event-specific PCR real-time detection methods for StarLink maize. <i>European Food Research and Technology</i> , <b>2003</b> , 216, 259-263	3.4	41
30	T-DNA integration in Arabidopsis chromosomes. Presence and origin of filler DNA sequences. <i>Plant Physiology</i> , <b>2003</b> , 133, 2061-8	6.6	85
29	Genetic and epigenetic aspects of somaclonal variation: flower colour bud sports in azalea, a case study. <i>South African Journal of Botany</i> , <b>2003</b> , 69, 117-128	2.9	8
28	Boosting heterologous protein production in transgenic dicotyledonous seeds using Phaseolus vulgaris regulatory sequences. <i>Nature Biotechnology</i> , <b>2002</b> , 20, 1265-8	44.5	148
27	Expression of antibodies and Fab fragments in transgenic potato plants: a case study for bulk production in crop plants. <i>Molecular Breeding</i> , <b>2002</b> , 9, 271-282	3.4	43
26	GATEWAY vectors for Agrobacterium-mediated plant transformation. <i>Trends in Plant Science</i> , <b>2002</b> , 7, 193-5	13.1	2693
25	Characterisation of the Roundup Ready soybean insert. <i>European Food Research and Technology</i> , <b>2001</b> , 213, 107-112	3.4	104
24	Highly efficient targeting and accumulation of a F(ab) fragment within the secretory pathway and apoplast of Arabidopsis thaliana. <i>FEBS Journal</i> , <b>2001</b> , 268, 4251-60		37
23	Transgene silencing of invertedly repeated transgenes is released upon deletion of one of the transgenes involved. <i>Plant Molecular Biology</i> , <b>2001</b> , 46, 433-45	4.6	48
22	Production of antibodies and antibody fragments in plants. <i>Vaccine</i> , <b>2001</b> , 19, 2756-61	4.1	35
21	Determination of the T-DNA transfer and the T-DNA integration frequencies upon cocultivation of Arabidopsis thaliana root explants. <i>Molecular Plant-Microbe Interactions</i> , <b>2000</b> , 13, 658-65	3.6	50
20	Isolation and characterization of recombinant antibody fragments against CDC2a from Arabidopsis thaliana. <i>FEBS Journal</i> , <b>2000</b> , 267, 6775-83		13
19	Plants as bioreactors for protein production: avoiding the problem of transgene silencing. <i>Plant Molecular Biology</i> , <b>2000</b> , 43, 347-59	4.6	108

18	The plantibody approach: expression of antibody genes in plants to modulate plant metabolism or to obtain pathogen resistance. <i>Plant Molecular Biology</i> , <b>2000</b> , 43, 419-28	4.6	56
17	T-DNA vector backbone sequences are frequently integrated into the genome of transgenic plants obtained by <i>Agrobacterium</i> -mediated transformation. <i>Molecular Breeding</i> , <b>2000</b> , 6, 459-468	3.4	104
16	Analysis of the interaction between single-chain variable fragments and their antigen in a reducing intracellular environment using the two-hybrid system. <i>FEBS Letters</i> , <b>2000</b> , 467, 316-20	3.8	14
15	Plants as bioreactors for protein production: avoiding the problem of transgene silencing <b>2000</b> , 227-239		4
14	High level accumulation of single-chain variable fragments in the cytosol of transgenic <i>Petunia hybrida</i> . <i>FEBS Journal</i> , <b>1999</b> , 259, 426-34		65
13	The DNA sequences of T-DNA junctions suggest that complex T-DNA loci are formed by a recombination process resembling T-DNA integration. <i>Plant Journal</i> , <b>1999</b> , 20, 295-304	6.9	119
12	<i>Agrobacterium tumefaciens</i> transformation and cotransformation frequencies of <i>Arabidopsis thaliana</i> root explants and tobacco protoplasts. <i>Molecular Plant-Microbe Interactions</i> , <b>1998</b> , 11, 449-57	3.6	59
11	Post-transcriptional gene silencing in plants. <i>Current Opinion in Cell Biology</i> , <b>1997</b> , 9, 373-82	9	212
10	Use of phage display for isolation and characterization of single-chain variable fragments against dihydroflavonol 4-reductase from <i>Petunia hybrida</i> . <i>FEBS Letters</i> , <b>1997</b> , 403, 116-22	3.8	12
9	T-DNA integration patterns in co-transformed plant cells suggest that T-DNA repeats originate from co-integration of separate T-DNAs. <i>Plant Journal</i> , <b>1997</b> , 11, 15-29	6.9	222
8	Post-transcriptional silencing of a neomycin phosphotransferase II transgene correlates with the accumulation of unproductive RNAs and with increased cytosine methylation of 3' flanking regions. <i>Plant Journal</i> , <b>1997</b> , 12, 379-392	6.9	77
7	Intact antigen-binding MAK33 antibody and Fab fragment accumulate in intercellular spaces of <i>Arabidopsis thaliana</i> . <i>Plant Science</i> , <b>1996</b> , 114, 233-241	5.3	52
6	Bacterial and plant-produced scFv proteins have similar antigen-binding properties. <i>FEBS Letters</i> , <b>1996</b> , 386, 5-10	3.8	50
5	Different 5' leader sequences modulate $\beta$ -glucuronidase accumulation levels in transgenic <i>Nicotiana tabacum</i> plants. <i>Euphytica</i> , <b>1995</b> , 85, 209-216	2.1	29
4	Quantitative kinetic analysis of $\beta$ -glucuronidase activities using a computer-directed microtiter plate reader. <i>Plant Molecular Biology Reporter</i> , <b>1993</b> , 11, 21-31	1.7	41
3	Assembly of an antibody and its derived antibody fragment in <i>Nicotiana</i> and <i>Arabidopsis</i> . <i>Transgenic Research</i> , <b>1993</b> , 2, 227-37	3.3	155
2	Frequencies of simultaneous transformation with different T-DNAs and their relevance to the <i>Agrobacterium</i> /plant cell interaction. <i>Molecular Genetics and Genomics</i> , <b>1985</b> , 201, 477-484		126
1	Transgene Silencing1-32		10

