
Agnese Latorre

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7232532/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Paclitaxel, cisplatin and lonidamine in advanced ovarian cancer. A phase II study. European Journal of Cancer, 2001, 37, 364-368.	1.3	151
2	Cytoskeleton and paclitaxel sensitivity in breast cancer: The role of β-tubulins. International Journal of Cancer, 2007, 120, 2078-2085.	2.3	132
3	Biomarkers predictive for clinical efficacy of taxol-based chemotherapy in advanced breast cancer. Annals of Oncology, 2005, 16, iv14-iv19.	0.6	127
4	Adjuvant anastrozole versus exemestane versus letrozole, upfront or after 2 years of tamoxifen, in endocrine-sensitive breast cancer (FATA-GIM3): a randomised, phase 3 trial. Lancet Oncology, The, 2018, 19, 474-485.	5.1	59
5	Predicting of Sentinel Lymph Node Status in Breast Cancer Patients with Clinically Negative Nodes: A Validation Study. Cancers, 2021, 13, 352.	1.7	33
6	Oral vinorelbine plus capecitabine (oral vincap) combination in patients with advanced breast cancer (ABC). A phase II study of the GOIM (Gruppo Oncologico dell'Italia Meridionale). Annals of Oncology, 2006, 17, vii15-vii17.	0.6	31
7	Revertant and potentiating activity of lonidamine in patients with ovarian cancer previously treated with platinum Journal of Clinical Oncology, 1997, 15, 3208-3213.	0.8	30
8	A multicenter REtrospective observational study of first-line treatment with PERtuzumab, trastuzumab and taxanes for advanced HER2 positive breast cancer patients. RePer Study. Cancer Biology and Therapy, 2019, 20, 192-200.	1.5	30
9	Subcutaneous recombinant interleukin-2 plus chemotherapy with cisplatin and dacarbazine in metastatic melanoma. European Journal of Cancer, 1996, 32, 730-733.	1.3	25
10	Liposomal-encapsulated doxorubicin plus cyclophosphamide as first-line therapy in metastatic breast cancer: a phase II multicentric study. Annals of Oncology, 2007, 18, vi66-vi69.	0.6	24
11	A Roadmap towards Breast Cancer Therapies Supported by Explainable Artificial Intelligence. Applied Sciences (Switzerland), 2021, 11, 4881.	1.3	24
12	A Clinical Decision Support System for Predicting Invasive Breast Cancer Recurrence: Preliminary Results. Frontiers in Oncology, 2021, 11, 576007.	1.3	21
13	A multicenter prospective phase II randomized trial of epirubicin/vinorelbine versus pegylated liposomal doxorubicin/vinorelbine as first-line treatment in advanced breast cancer. A GOIM study. Journal of Experimental and Clinical Cancer Research, 2011, 30, 39.	3.5	20
14	A Phase I Study of Capecitabine in Combination with Vinorelbine in Advanced Breast Cancer. Clinical Breast Cancer, 2003, 4, 138-141.	1.1	19
15	Long-term Subcutaneous Recombinant Interleukin-2 as Maintenance Therapy: Biological Effects and Clinical Implications. Cancer Biotherapy, 1995, 10, 195-203.	0.6	18
16	Eribulin Mesylate as Third or Subsequent Line Chemotherapy for Elderly Patients with Locally Recurrent or Metastatic Breast Cancer: A Multicentric Observational Study of GIOGer (Italian Group) Tj ETQq0 0	0 ng₿T /Ov	erlosck 10 Tf
17	Robustness Evaluation of a Deep Learning Model on Sagittal and Axial Breast DCE-MRIs to Predict Pathological Complete Response to Neoadjuvant Chemotherapy. Journal of Personalized Medicine, 2022, 12, 953.	1.1	15

Demographic, tumor and clinical features of clinical trials versus clinical practice patients with HER2-positive early breast cancer: results of a prospective study. Journal of Cancer Research and 1.2 14 Clinical Oncology, 2016, 142, 669-678.

Agnese Latorre

#	ARTICLE	IF	CITATIONS
19	FISH testing of HER2 immunohistochemistry 1+ invasive breast cancer with unfavorable characteristics. Oncology Letters, 2016, 12, 3115-3122.	0.8	13
20	Efficacy and safety of eribulin in taxane-refractory patients in the â€real world'. Future Oncology, 2017, 13, 971-978.	1.1	13
21	Chemotherapy Options beyond the First Line in HER-Negative Metastatic Breast Cancer. Journal of Oncology, 2020, 2020, 1-17.	0.6	11
22	Dose intensity and efficacy of the combination of everolimus and exemestane (EVE/EXE) in a real-world population of hormone receptor-positive (ER+/PgR+), HER2-negative advanced breast cancer (ABC) patients: a multicenter Italian experience. Breast Cancer Research and Treatment, 2017, 163, 587-594.	1.1	9
23	Phase I/II study of gemcitabine plus mitoxantrone as salvage chemotherapy in metastatic breast cancer. British Journal of Cancer, 2003, 88, 491-495.	2.9	8
24	A Multicenter Phase II Randomized Trial of Docetaxel/Gemcitabine versus Docetaxel/Capecitabine as First-Line Treatment for Advanced Breast Cancer: A Gruppo Oncologico Italia Meridionale Study. Oncology, 2011, 81, 230-236.	0.9	8
25	An Invasive Disease Event-Free Survival Analysis to Investigate Ki67 Role with Respect to Breast Cancer Patients' Age: A Retrospective Cohort Study. Cancers, 2022, 14, 2215.	1.7	4
26	A broad Italian experience with eribulin mesylate in metastatic breast cancer patients: The ESEMPiO study Journal of Clinical Oncology, 2015, 33, e11539-e11539.	0.8	3
27	Downstream Signaling of Inflammasome Pathway Affects Patients' Outcome in the Context of Distinct Molecular Breast Cancer Subtypes. Pharmaceuticals, 2022, 15, 651.	1.7	3
28	Amifostine as chemoprotectant in metastatic breast cancer patients treated with doxorubicin. Oncology Reports, 2003, 10, 163.	1.2	2
29	Efficacy and safety of lenvatinib in an elderly patient with metastatic papillary thyroid carcinoma and cardiological comorbidity: a case report. Future Oncology, 2019, 15, 27-33.	1.1	1
30	Mitoxantrone, L-Leucovorin and 5-Fluorouracil: An Effective and Well Tolerated First-Line Treatment for Advanced Breast Cancer. Tumori, 1999, 85, 60-64.	0.6	0
31	Rapid tumor shrinkage with lapatinib plus capecitabine in a patient with massive liver involvement. Tumori, 2013, 99, 278e-81e.	0.6	0