

RocÃ-o Mateo-Gallego

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7238564/publications.pdf>

Version: 2024-02-01

22
papers

282
citations

840776

11
h-index

940533

16
g-index

23
all docs

23
docs citations

23
times ranked

475
citing authors

#	ARTICLE	IF	CITATIONS
1	Effect of the Consumption of Alcohol-Free Beers with Different Carbohydrate Composition on Postprandial Metabolic Response. <i>Nutrients</i> , 2022, 14, 1046.	4.1	3
2	Triglyceride Metabolism Modifies Lipoprotein(a) Plasma Concentration. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, 107, e3594-e3602.	3.6	5
3	Diagnostic yield of sequencing familial hypercholesterolemia genes in individuals with primary hypercholesterolemia. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2021, 74, 664-673.	0.6	5
4	An alcohol-free beer enriched with isomaltulose and a resistant dextrin modulates gut microbiome in subjects with type 2 diabetes mellitus and overweight or obesity: a pilot study. <i>Food and Function</i> , 2021, 12, 3635-3646.	4.6	19
5	Effect of Novel Alcohol-Free Beer Recipes Enriched With Isomaltulose and a Resistant Dextrin on the Metabolic Postprandial Response in Healthy Subjects. <i>Current Developments in Nutrition</i> , 2021, 5, 340.	0.3	0
6	High-protein energy-restricted diets induce greater improvement in glucose homeostasis but not in adipokines comparing to standard-protein diets in early-onset diabetic adults with overweight or obesity. <i>Clinical Nutrition</i> , 2020, 39, 1354-1363.	5.0	10
7	Effect of an alcohol-free beer enriched with isomaltulose and a resistant dextrin on insulin resistance in diabetic patients with overweight or obesity. <i>Clinical Nutrition</i> , 2020, 39, 475-483.	5.0	30
8	Predicted pathogenic mutations in STAP1 are not associated with clinically defined familial hypercholesterolemia. <i>Atherosclerosis</i> , 2020, 292, 143-151.	0.8	21
9	Glycerol kinase deficiency in adults: Description of 4 novel cases, systematic review and development of a clinical diagnostic score. <i>Atherosclerosis</i> , 2020, 315, 24-32.	0.8	3
10	Effect of Lifestyle Intervention in the Concentration of Adipoquines and Branched Chain Amino Acids in Subjects with High Risk of Developing Type 2 Diabetes: Feel4Diabetes Study. <i>Cells</i> , 2020, 9, 693.	4.1	7
11	Lipid-lowering response in subjects with the p.(Leu167del) mutation in the APOE gene. <i>Atherosclerosis</i> , 2019, 282, 143-147.	0.8	12
12	Dietary polyunsaturated fatty acids mediate the inverse association of stearoyl-CoA desaturase activity with the risk of fatty liver in dyslipidaemic individuals. <i>European Journal of Nutrition</i> , 2019, 58, 1561-1568.	3.9	6
13	Single Nucleotide Variants Associated With Polygenic Hypercholesterolemia in Families Diagnosed Clinically With Familial Hypercholesterolemia. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2018, 71, 351-356.	0.6	3
14	Cholesterol oversynthesis markers define familial combined hyperlipidemia versus other genetic hypercholesterolemias independently of body weight. <i>Journal of Nutritional Biochemistry</i> , 2018, 53, 48-57.	4.2	14
15	Efficacy of repeated phlebotomies in hypertriglyceridemia and iron overload: A prospective, randomized, controlled trial. <i>Journal of Clinical Lipidology</i> , 2018, 12, 1190-1198.	1.5	6
16	Association between non-cholesterol sterol concentrations and Achilles tendon thickness in patients with genetic familial hypercholesterolemia. <i>Journal of Translational Medicine</i> , 2018, 16, 6.	4.4	10
17	Effect of intensive LDL cholesterol lowering with PCSK9 monoclonal antibodies on tendon xanthoma regression in familial hypercholesterolemia. <i>Atherosclerosis</i> , 2017, 263, 92-96.	0.8	14
18	ABCG5/G8 gene is associated with hypercholesterolemias without mutation in candidate genes and noncholesterol sterols. <i>Journal of Clinical Lipidology</i> , 2017, 11, 1432-1440.e4.	1.5	33

#	ARTICLE	IF	CITATIONS
19	Adherence to a Mediterranean diet is associated with the presence and extension of atherosclerotic plaques in middle-aged asymptomatic adults: The Aragon Workers' Health Study. <i>Journal of Clinical Lipidology</i> , 2017, 11, 1372-1382.e4.	1.5	12
20	Lipid phenotype and heritage pattern in families with genetic hypercholesterolemia not related to LDLR, APOB, PCSK9, or APOE. <i>Journal of Clinical Lipidology</i> , 2016, 10, 1397-1405.e2.	1.5	12
21	Effect of Nicotinic acid/Laropirant in the lipoprotein(a) concentration with regard to baseline lipoprotein(a) concentration and LPA genotype. <i>Metabolism: Clinical and Experimental</i> , 2014, 63, 365-371.	3.4	15
22	Common Genetic Variants Contribute to Primary Hypertriglyceridemia Without Differences Between Familial Combined Hyperlipidemia and Isolated Hypertriglyceridemia. <i>Circulation: Cardiovascular Genetics</i> , 2014, 7, 814-821.	5.1	36