

Florian Duclot

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7239022/publications.pdf>

Version: 2024-02-01

23
papers

1,335
citations

471509

17
h-index

677142

22
g-index

23
all docs

23
docs citations

23
times ranked

2337
citing authors

#	ARTICLE	IF	CITATIONS
1	Epigenetics of Aggression. <i>Current Topics in Behavioral Neurosciences</i> , 2021, , 1.	1.7	0
2	Transcriptomic Regulations Underlying Pair-bond Formation and Maintenance in the Socially Monogamous Male and Female Prairie Vole. <i>Biological Psychiatry</i> , 2020, 91, 141-151.	1.3	14
3	Consequences of prenatal exposure to valproic acid in the socially monogamous prairie voles. <i>Scientific Reports</i> , 2019, 9, 2453.	3.3	18
4	Epigenetic regulation of motivated behaviors by histone deacetylase inhibitors. <i>Neuroscience and Biobehavioral Reviews</i> , 2019, 105, 305-317.	6.1	18
5	Hippocampal protein kinase D1 is necessary for DHPG-induced learning and memory impairments in rats. <i>PLoS ONE</i> , 2018, 13, e0195095.	2.5	3
6	Comparative Transcriptomic Analysis of the Effects of Antidepressant Drugs in Stress-Susceptible Mice. <i>Biological Psychiatry</i> , 2017, 81, 278-279.	1.3	1
7	Integrative analysis of sex differences in the rapid antidepressant effects of ketamine in preclinical models for individualized clinical outcomes. <i>Current Opinion in Behavioral Sciences</i> , 2017, 14, 19-26.	3.9	28
8	The Role of Early Growth Response 1 (EGR1) in Brain Plasticity and Neuropsychiatric Disorders. <i>Frontiers in Behavioral Neuroscience</i> , 2017, 11, 35.	2.0	268
9	Trichostatin A (TSA) facilitates formation of partner preference in male prairie voles (<i>Microtus</i>) Tj ETQq1 1 0.784314 1.1gBT /Overlock 10	2.1	27
10	Prediction of individual differences in fear response by novelty seeking, and disruption of contextual fear memory reconsolidation by ketamine. <i>Neuropharmacology</i> , 2016, 109, 293-305.	4.1	39
11	The estrous cycle surpasses sex differences in regulating the transcriptome in the rat medial prefrontal cortex and reveals an underlying role of early growth response 1. <i>Genome Biology</i> , 2015, 16, 256.	8.8	54
12	Regulated internalization of NMDA receptors drives PKD1-mediated suppression of the activity of residual cell-surface NMDA receptors. <i>Molecular Brain</i> , 2015, 8, 75.	2.6	6
13	Methyl Supplementation Attenuates Cocaine-Seeking Behaviors and Cocaine-Induced c-Fos Activation in a DNA Methylation-Dependent Manner. <i>Journal of Neuroscience</i> , 2015, 35, 8948-8958.	3.6	101
14	Epigenetic mechanisms underlying the role of brain-derived neurotrophic factor in depression and response to antidepressants. <i>Journal of Experimental Biology</i> , 2015, 218, 21-31.	1.7	82
15	The Anxiolytic and Antidepressant-like Effects of Testosterone and Estrogen in Gonadectomized Male Rats. <i>Biological Psychiatry</i> , 2015, 78, 259-269.	1.3	88
16	Histone deacetylase inhibitors facilitate partner preference formation in female prairie voles. <i>Nature Neuroscience</i> , 2013, 16, 919-924.	14.8	117
17	Individual Differences in Novelty Seeking Predict Subsequent Vulnerability to Social Defeat through a Differential Epigenetic Regulation of Brain-Derived Neurotrophic Factor Expression. <i>Journal of Neuroscience</i> , 2013, 33, 11048-11060.	3.6	90
18	Cognitive impairments in adult mice with constitutive inactivation of <i>RIP140</i> gene expression. <i>Genes, Brain and Behavior</i> , 2012, 11, 69-78.	2.2	36

#	ARTICLE	IF	CITATIONS
19	Individual differences in the effect of social defeat on anhedonia and histone acetylation in the rat hippocampus. <i>Hormones and Behavior</i> , 2011, 59, 331-337.	2.1	91
20	Individual differences in novelty-seeking behavior in rats as a model for psychosocial stress-related mood disorders. <i>Physiology and Behavior</i> , 2011, 104, 296-305.	2.1	41
21	Alteration of working memory but not in anxiety or stress response in p300/CBP associated factor (PCAF) histone acetylase knockout mice bred on a C57BL/6 background. <i>Neuroscience Letters</i> , 2010, 475, 179-183.	2.1	33
22	Mice knock out for the histone acetyltransferase p300/CREB binding protein-associated factor develop a resistance to amyloid toxicity. <i>Neuroscience</i> , 2010, 167, 850-863.	2.3	47
23	Altered Memory Capacities and Response to Stress in p300/CBP-Associated Factor (PCAF) Histone Acetylase Knockout Mice. <i>Neuropsychopharmacology</i> , 2008, 33, 1584-1602.	5.4	133