

Vasileios I Sakalis

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7244130/publications.pdf>

Version: 2024-02-01

19
papers

242
citations

1478505

6
h-index

996975

15
g-index

20
all docs

20
docs citations

20
times ranked

208
citing authors

#	ARTICLE	IF	CITATIONS
1	European Association of Urology Guidelines on the Diagnosis and Management of Female Non-neurogenic Lower Urinary Tract Symptoms. Part 1: Diagnostics, Overactive Bladder, Stress Urinary Incontinence, and Mixed Urinary Incontinence. <i>European Urology</i> , 2022, 82, 49-59.	1.9	87
2	Medical Treatment of Nocturia in Men with Lower Urinary Tract Symptoms: Systematic Review by the European Association of Urology Guidelines Panel for Male Lower Urinary Tract Symptoms. <i>European Urology</i> , 2017, 72, 757-769.	1.9	59
3	Transurethral holmium laser intravesical tape excision following TVT procedure: results from seven patients in a 12-month follow-up. <i>International Urogynecology Journal</i> , 2012, 23, 769-777.	1.4	14
4	Management of stress urinary incontinence in spinal cord injured female patients with a mid-urethral tape – a single center experience. <i>Journal of Spinal Cord Medicine</i> , 2018, 41, 703-709.	1.4	13
5	Pain point system scale (PPSS): a method for postoperative pain estimation in retrospective studies. <i>Journal of Pain Research</i> , 2012, 5, 503.	2.0	12
6	European Association of Urology Guidelines on the Management of Female Non-neurogenic Lower Urinary Tract Symptoms. Part 2: Underactive Bladder, Bladder Outlet Obstruction, and Nocturia. <i>European Urology</i> , 2022, 82, 60-70.	1.9	12
7	Combination of solifenacin with tamsulosin reduces prostate volume and vascularity as opposed to tamsulosin monotherapy in patients with benign prostate enlargement and overactive bladder symptoms: Results from a randomized pilot study. <i>International Journal of Urology</i> , 2018, 25, 737-745.	1.0	8
8	Diagnostic Tests for Female Bladder Outlet Obstruction: A Systematic Review from the European Association of Urology Non-neurogenic Female LUTS Guidelines Panel. <i>European Urology Focus</i> , 2022, 8, 1015-1030.	3.1	8
9	Mitrofanoff cystolitholopaxy and bladder biopsy using a mini-PCNL kit. <i>Journal of Clinical Urology</i> , 2017, 10, 36-38.	0.1	6
10	What Is the Most Effective Treatment for Nocturia or Nocturnal Incontinence in Adult Women?. <i>European Urology Focus</i> , 2021, 7, 453-463.	3.1	6
11	What Is the Most Effective Management of the Primary Tumor in Men with Invasive Penile Cancer: A Systematic Review of the Available Treatment Options and Their Outcomes. <i>European Urology Open Science</i> , 2022, 40, 58-94.	0.4	6
12	Improving Guideline Adherence in Urology. <i>European Urology Focus</i> , 2022, 8, 1545-1552.	3.1	5
13	The effect of pharmacotherapy on prostate volume, prostate perfusion and prostate-specific antigens (prostate morphometric parameters) in patients with lower urinary tract symptoms and benign prostatic obstruction. A systematic review and meta-analysis. <i>Central European Journal of Urology</i> , 2021, 74, 388-421.	0.3	2
14	Detrusor Overactivity May Be a Prognostic Factor for Better Response to Combination Therapy Over Monotherapy in Male Patients With Benign Prostatic Enlargement and Storage Lower Urinary Tract Symptoms. <i>International Neurourology Journal</i> , 2021, 25, 69-76.	1.2	2
15	Do Drugs Work for ŸŸŸ Following Prostate Cancer Surgery. <i>Current Drug Targets</i> , 2020, 21, 1515-1526.	2.1	2
16	A pure microcytic bladder carcinoma synchronous to prostatic adenocarcinoma. <i>Rare Tumors</i> , 2011, 3, 95-97.	0.6	0
17	Reply to Jae Hung Jung, Julia L. Han, and Philipp Dahm's Letter to the Editor re: Vasileios I. Sakalis, Markos Karavitakis, Dina Bedretdinova, et al. Medical Treatment of Nocturia in Men with Lower Urinary Tract Symptoms: Systematic Review by the European Association of Urology Guidelines Panel for Male Lower Urinary Tract Symptoms. <i>Eur Urol</i> 2017;72:757-769. <i>European Urology</i> , 2018, 73, e60-e62.	1.9	0
18	Macroplastique and Botox are superior to Macroplastique alone in the management of neurogenic vesicoureteric reflux in spinal cord injury population with presumed healthy bladders. <i>Journal of Spinal Cord Medicine</i> , 2019, 42, 478-484.	1.4	0

#	ARTICLE	IF	CITATIONS
19	Subtle errors of bladder wall thickness measurement have a significant impact on the calculation of ultrasound-estimated bladder weight. A pilot study. Medical Ultrasonography, 2018, 20, 292.	0.8	0