

Isabelle Herter-Aeberli

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7245239/isabelle-herter-aeberli-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

77
papers

6,592
citations

30
h-index

81
g-index

84
ext. papers

8,358
ext. citations

6.1
avg, IF

6.07
L-index

#	Paper	IF	Citations
77	The Potential of Prebiotic and Probiotic Supplementation During Obese Pregnancy to Improve Maternal and Offspring Metabolic Health and Reduce Obesity Risk-A Narrative Review.. <i>Frontiers in Nutrition</i> , 2022 , 9, 819882	6.2	3
76	Kinetics of iron absorption from ferrous fumarate with and without galacto-oligosaccharides determined from stable-isotope appearance curves in women. <i>American Journal of Clinical Nutrition</i> , 2021 ,	7	1
75	The effects of reducing chronic inflammation in overweight women on serum hepcidin and iron absorption with and without supplemental ascorbic acid. <i>British Journal of Nutrition</i> , 2021 , 126, 877-884 ^{3.6}		
74	Strengthening the immunity of the Swiss population with micronutrients: A narrative review and call for action. <i>Clinical Nutrition ESPEN</i> , 2021 , 43, 39-48	1.3	9
73	Kinetics of Iron Absorption From Ferrous Fumarate With Galacto-oligosaccharides Measured by Stable-isotope Appearance Curves in Iron Depleted Women in Switzerland. <i>Current Developments in Nutrition</i> , 2021 , 5, 1317-1317	0.4	78
72	Vegan Diet and Bone Health-Results from the Cross-Sectional RBVD Study. <i>Nutrients</i> , 2021 , 13,	6.7	12
71	A test to measure oral iron absorption and glucose tolerance simultaneously in 18 to 55 year old premenopausal women. <i>Clinical Nutrition ESPEN</i> , 2021 , 46, 325-329	1.3	
70	Inadequate Status and Low Awareness of Folate in Switzerland-A Call to Strengthen Public Health Measures to Ensure Sufficient Intakes. <i>Nutrients</i> , 2020 , 12,	6.7	3
69	In women, central obesity predicts higher inflammation, higher serum hepcidin, lower absorption and hypoferrremia. <i>Proceedings of the Nutrition Society</i> , 2020 , 79,	2.9	1
68	Height and body-mass index trajectories of school-aged children and adolescents from 1985 to 2019 in 200 countries and territories: a pooled analysis of 2181 population-based studies with 65 million participants. <i>Lancet, The</i> , 2020 , 396, 1511-1524	40	73
67	The effect of central obesity on inflammation, hepcidin, and iron metabolism in young women. <i>International Journal of Obesity</i> , 2020 , 44, 1291-1300	5.5	15
66	Compared to Normal-Weight Pregnant Women, Overweight Pregnant Women Fail to Upregulate Iron Absorption in the Third Trimester Despite Iron Deficiency and Their Infants Are Born with Lower Body Iron Stores. <i>Blood</i> , 2020 , 136, 51-52	2.2	
65	Vitamin and Mineral Status in a Vegan Diet. <i>Deutsches A&#x0308;rztblatt International</i> , 2020 , 117, 575-582	2.5	23
64	International Waist Circumference Percentile Cutoffs for Central Obesity in Children and Adolescents Aged 6 to 18 Years. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	22
63	Acute Consumption of Prebiotic Galacto-Oligosaccharides Increases Iron Absorption from Ferrous Fumarate, but not from Ferrous Sulfate and Ferric Pyrophosphate: Stable Iron Isotope Studies in Iron-Depleted Young Women. <i>Journal of Nutrition</i> , 2020 , 150, 2391-2397	4.1	5
62	Addition of Whole Wheat Flour During Injera Fermentation Degrades Phytic Acid and Triples Iron Absorption from Fortified Tef in Young Women. <i>Journal of Nutrition</i> , 2020 , 150, 2666-2672	4.1	
61	Effectiveness of increased salt iodine concentration on iodine status: trend analysis of cross-sectional national studies in Switzerland. <i>European Journal of Nutrition</i> , 2020 , 59, 581-593	5.2	12

60	Significant Decrease in Childhood Obesity and Waist Circumference over 15 Years in Switzerland: A Repeated Cross-Sectional Study. <i>Nutrients</i> , 2019 , 11,	6.7	7
59	Asymptomatic Infection in Preschool Children and Young Women Does Not Predict Iron Bioavailability from Iron-Fortified Foods. <i>Nutrients</i> , 2019 , 11,	6.7	3
58	Effects of feed iodine concentrations and milk processing on iodine concentrations of cowsQmilk and dairy products, and potential impact on iodine intake in Swiss adults. <i>British Journal of Nutrition</i> , 2019 , 122, 172-185	3.6	6
57	Rising rural body-mass index is the main driver of the global obesity epidemic in adults. <i>Nature</i> , 2019 , 569, 260-264	50.4	278
56	Effects of exergame training combined with omega-3 fatty acids on the elderly brain: a randomized double-blind placebo-controlled trial. <i>BMC Geriatrics</i> , 2019 , 19, 81	4.1	7
55	Greater blood volume and Hb mass in obese women quantified by the carbon monoxide-rebreathing method affects interpretation of iron biomarkers and iron requirements. <i>International Journal of Obesity</i> , 2019 , 43, 999-1008	5.5	14
54	Validation of a Food Frequency Questionnaire to Assess Intake of Polyunsaturated Fatty Acids in Switzerland. <i>Nutrients</i> , 2019 , 11,	6.7	6
53	Iodine bioavailability from cow milk: a randomized, crossover balance study in healthy iodine-replete adults. <i>American Journal of Clinical Nutrition</i> , 2019 , 110, 102-110	7	8
52	Consumption of Galacto-Oligosaccharides Increases Iron Absorption from Ferrous Fumarate: A Stable Iron Isotope Study in Iron-Depleted Young Women. <i>Journal of Nutrition</i> , 2019 , 149, 738-746	4.1	16
51	The main determinants of iodine in cowsQmilk in Switzerland are farm type, season and teat dipping. <i>British Journal of Nutrition</i> , 2018 , 119, 559-569	3.6	16
50	Micronutrient status and intake in omnivores, vegetarians and vegans in Switzerland. <i>European Journal of Nutrition</i> , 2017 , 56, 283-293		120
49	Moderate-to-Severe Iodine Deficiency in the "First 1000 Days" Causes More Thyroid Hypofunction in Infants Than in Pregnant or Lactating Women. <i>Journal of Nutrition</i> , 2017 , 147, 589-595	4.1	19
48	The effect of body composition and serum inflammatory markers on the functional muscle-bone unit in premenopausal women. <i>International Journal of Obesity</i> , 2017 , 41, 1203-1206	5.5	4
47	Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 1289 million children, adolescents, and adults. <i>Lancet, The</i> , 2017 , 390, 2627-2642	40	2980
46	In Haitian women and preschool children, iron absorption from wheat flour-based meals fortified with sodium iron EDTA is higher than that from meals fortified with ferrous fumarate, and is not affected by <i>Helicobacter pylori</i> infection in children. <i>British Journal of Nutrition</i> , 2017 , 118, 273-279	3.6	9
45	Dried Blood Spot Thyroglobulin as a Biomarker of Iodine Status in Pregnant Women. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017 , 102, 23-32	5.6	44
44	The Potential of Fermentation and Contamination of Teff by Soil to Influence Iron Intake and Bioavailability from Injera Flatbread. <i>International Journal for Vitamin and Nutrition Research</i> , 2017 , 87, 75-84	1.7	4
43	Increased risk of iron deficiency and reduced iron absorption but no difference in zinc, vitamin A or B-vitamin status in obese women in India. <i>European Journal of Nutrition</i> , 2016 , 55, 2411-2421		12

42	Risk factors for overweight and obesity in Swiss primary school children: results from a representative national survey. <i>European Journal of Nutrition</i> , 2016 , 55, 621-629	5.2	9
41	The effects of fat loss after bariatric surgery on inflammation, serum hepcidin, and iron absorption: a prospective 6-mo iron stable isotope study. <i>American Journal of Clinical Nutrition</i> , 2016 , 104, 1030-1038	7	29
40	Oral Vitamin D Supplements Increase Serum 25-Hydroxyvitamin D in Postmenopausal Women and Reduce Bone Calcium Flux Measured by ⁴¹ Ca Skeletal Labeling. <i>Journal of Nutrition</i> , 2015 , 145, 2333-40	4.1	4
39	Iodine Supplementation Decreases Hypercholesterolemia in Iodine-Deficient, Overweight Women: A Randomized Controlled Trial. <i>Journal of Nutrition</i> , 2015 , 145, 2067-75	4.1	21
38	In overweight and obese women, dietary iron absorption is reduced and the enhancement of iron absorption by ascorbic acid is one-half that in normal-weight women. <i>American Journal of Clinical Nutrition</i> , 2015 , 102, 1389-97	7	48
37	Efficient ⁴¹ Ca measurements for biomedical applications. <i>Nuclear Instruments & Methods in Physics Research B</i> , 2015 , 361, 273-276	1.2	8
36	Phytic acid degrading lactic acid bacteria in tef-injera fermentation. <i>International Journal of Food Microbiology</i> , 2014 , 190, 54-60	5.8	53
35	Pediatric adiposity stabilized in Switzerland between 1999 and 2012. <i>European Journal of Nutrition</i> , 2014 , 53, 865-75	5.2	15
34	Sugar-sweetened beverages with moderate amounts of fructose, but not sucrose, induce Fatty Acid synthesis in healthy young men: a randomized crossover study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, 2164-72	5.6	11
33	Antioxidant supplements reduced oxidative stress and stabilized liver function tests but did not reduce inflammation in a randomized controlled trial in obese children and adolescents. <i>Journal of Nutrition</i> , 2014 , 144, 193-201	4.1	44
32	Disturbed eating at high altitude: influence of food preferences, acute mountain sickness and satiation hormones. <i>European Journal of Nutrition</i> , 2013 , 52, 625-35	5.2	33
31	Thyroglobulin is a sensitive measure of both deficient and excess iodine intakes in children and indicates no adverse effects on thyroid function in the UIC range of 100-299 µg/L: a UNICEF/ICCIDD study group report. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 1271-80	5.6	146
30	Reduced insulin sensitivity as a marker for acute mountain sickness?. <i>High Altitude Medicine and Biology</i> , 2013 , 14, 240-50	1.9	9
29	A composite score combining waist circumference and body mass index more accurately predicts body fat percentage in 6- to 13-year-old children. <i>European Journal of Nutrition</i> , 2013 , 52, 247-53	5.2	12
28	Response to Comment on: Aeberli et al. Moderate amounts of fructose consumption impair insulin sensitivity in healthy young men: a randomized controlled trial. <i>Diabetes Care</i> 2013;36:150-156. <i>Diabetes Care</i> , 2013 , 36, e105	14.6	1
27	Moderate amounts of fructose consumption impair insulin sensitivity in healthy young men: a randomized controlled trial. <i>Diabetes Care</i> , 2013 , 36, 150-6	14.6	138
26	Micronutrient supply and health outcomes in children. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2013 , 16, 328-38	3.8	7
25	Overweight impairs efficacy of iron supplementation in iron-deficient South African children: a randomized controlled intervention. <i>International Journal of Obesity</i> , 2013 , 37, 24-30	5.5	36

24	Reply to JS White. <i>American Journal of Clinical Nutrition</i> , 2012 , 95, 773-774	7	
23	Waist circumference and waist-to-height ratio percentiles in a nationally representative sample of 6-13 year old children in Switzerland. <i>Swiss Medical Weekly</i> , 2011 , 141, w13227	3.1	31
22	Evidence that the prevalence of childhood overweight is plateauing: data from nine countries. <i>Pediatric Obesity</i> , 2011 , 6, 342-60		429
21	Ten repeat collections for urinary iodine from spot samples or 24-hour samples are needed to reliably estimate individual iodine status in women. <i>Journal of Nutrition</i> , 2011 , 141, 2049-54	4.1	220
20	Low to moderate sugar-sweetened beverage consumption impairs glucose and lipid metabolism and promotes inflammation in healthy young men: a randomized controlled trial. <i>American Journal of Clinical Nutrition</i> , 2011 , 94, 479-85	7	229
19	The thin-fat phenotype and global metabolic disease risk. <i>Current Opinion in Clinical Nutrition and Metabolic Care</i> , 2011 , 14, 542-7	3.8	30
18	Sharply higher rates of iron deficiency in obese Mexican women and children are predicted by obesity-related inflammation rather than by differences in dietary iron intake. <i>American Journal of Clinical Nutrition</i> , 2011 , 93, 975-83	7	129
17	Baseline leptin and leptin reduction predict improvements in metabolic variables and long-term fat loss in obese children and adolescents: a prospective study of an inpatient weight-loss program. <i>American Journal of Clinical Nutrition</i> , 2011 , 93, 695-702	7	29
16	Decrease in the prevalence of paediatric adiposity in Switzerland from 2002 to 2007. <i>Public Health Nutrition</i> , 2010 , 13, 806-11	3.3	38
15	During rapid weight loss in obese children, reductions in TSH predict improvements in insulin sensitivity independent of changes in body weight or fat. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, 5412-8	5.6	52
14	Does obesity increase risk for iron deficiency? A review of the literature and the potential mechanisms. <i>International Journal for Vitamin and Nutrition Research</i> , 2010 , 80, 263-70	1.7	112
13	The Swiss iodized salt program provides adequate iodine for school children and pregnant women, but weaning infants not receiving iodine-containing complementary foods as well as their mothers are iodine deficient. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010 , 95, 5217-24	5.6	102
12	Stabilization of the prevalence of childhood obesity in Switzerland. <i>Swiss Medical Weekly</i> , 2010 , 140, w13046	3.1	26
11	Iodine treatment in children with subclinical hypothyroidism due to chronic iodine deficiency decreases thyrotropin and C-peptide concentrations and improves the lipid profile. <i>Thyroid</i> , 2009 , 19, 1099-104	6.2	30
10	Overweight children have higher circulating hepcidin concentrations and lower iron status but have dietary iron intakes and bioavailability comparable with normal weight children. <i>International Journal of Obesity</i> , 2009 , 33, 1111-7	5.5	124
9	Diet determines features of the metabolic syndrome in 6- to 14-year-old children. <i>International Journal for Vitamin and Nutrition Research</i> , 2009 , 79, 14-23	1.7	21
8	The increase of fatty acid-binding protein aP2 in overweight and obese children: interactions with dietary fat and impact on measures of subclinical inflammation. <i>International Journal of Obesity</i> , 2008 , 32, 1513-20	5.5	21
7	Dietary determinants of subclinical inflammation, dyslipidemia and components of the metabolic syndrome in overweight children: a review. <i>International Journal of Obesity</i> , 2008 , 32 Suppl 6, S11-8	5.5	42

6	Adiposity in women and children from transition countries predicts decreased iron absorption, iron deficiency and a reduced response to iron fortification. <i>International Journal of Obesity</i> , 2008 , 32, 1098-104	5.5	142
5	Serum retinol-binding protein 4 concentration and its ratio to serum retinol are associated with obesity and metabolic syndrome components in children. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007 , 92, 4359-65	5.6	115
4	Interaction of positional isomers of quercetin glucuronides with the transporter ABCC2 (cMOAT, MRP2). <i>Drug Metabolism and Disposition</i> , 2007 , 35, 1262-8	4	68
3	Fructose intake is a predictor of LDL particle size in overweight schoolchildren. <i>American Journal of Clinical Nutrition</i> , 2007 , 86, 1174-8	7	74
2	Glycaemic control, insulin resistance and obesity 2007 , 43-57		1
1	Dietary intakes of fat and antioxidant vitamins are predictors of subclinical inflammation in overweight Swiss children. <i>American Journal of Clinical Nutrition</i> , 2006 , 84, 748-55	7	91