

Isabel Fabregat

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7246301/publications.pdf>

Version: 2024-02-01

148
papers

8,832
citations

36203

51
h-index

46693

89
g-index

149
all docs

149
docs citations

149
times ranked

11971
citing authors

#	ARTICLE	IF	CITATIONS
1	Mitochondrial bioenergetics boost macrophage activation, promoting liver regeneration in metabolically compromised animals. <i>Hepatology</i> , 2022, 75, 550-566.	3.6	25
2	Proteoglycans in Cancer: Friends or Enemies? A Special Focus on Hepatocellular Carcinoma. <i>Cancers</i> , 2022, 14, 1902.	1.7	11
3	The Tumor Microenvironment Drives Intrahepatic Cholangiocarcinoma Progression. <i>International Journal of Molecular Sciences</i> , 2022, 23, 4187.	1.8	4
4	NADPH oxidase 4 (Nox4) deletion accelerates liver regeneration in mice. <i>Redox Biology</i> , 2021, 40, 101841.	3.9	13
5	Epithelial-Mesenchymal Transition (EMT) Induced by TGF- β 2 in Hepatocellular Carcinoma Cells Reprograms Lipid Metabolism. <i>International Journal of Molecular Sciences</i> , 2021, 22, 5543.	1.8	35
6	Anti-miR-518d-5p overcomes liver tumor cell death resistance through mitochondrial activity. <i>Cell Death and Disease</i> , 2021, 12, 555.	2.7	10
7	2-[18F]FDG PET/CT as a Predictor of Microvascular Invasion and High Histological Grade in Patients with Hepatocellular Carcinoma. <i>Cancers</i> , 2021, 13, 2554.	1.7	10
8	The TGF- β 2 Pathway: A Pharmacological Target in Hepatocellular Carcinoma?. <i>Cancers</i> , 2021, 13, 3248.	1.7	37
9	Anti-TGF- β 2 (Transforming Growth Factor β 2) Therapy With Betaglycan-Derived P144 Peptide Gene Delivery Prevents the Formation of Aortic Aneurysm in a Mouse Model of Marfan Syndrome. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2021, 41, e440-e452.	1.1	12
10	The TGF- β 2/NADPH Oxidases Axis in the Regulation of Liver Cell Biology in Health and Disease. <i>Cells</i> , 2021, 10, 2312.	1.8	14
11	Direct and Indirect Effect of TGF- β 2 on Treg Transendothelial Recruitment in HCC Tissue Microenvironment. <i>International Journal of Molecular Sciences</i> , 2021, 22, 11765.	1.8	7
12	Clathrin switches transforming growth factor- β 2 role to pro-tumorigenic in liver cancer. <i>Journal of Hepatology</i> , 2020, 72, 125-134.	1.8	30
13	Relevance of epidermal growth factor receptor kinase activity in a model of cholestatic liver injury. <i>Journal of Hepatology</i> , 2020, 73, S202.	1.8	0
14	Editorial Special Issue TGF-Beta/BMP Signaling Pathway. <i>Cells</i> , 2020, 9, 2363.	1.8	2
15	Encapsulating TGF- β 1 Inhibitory Peptides P17 and P144 as a Promising Strategy to Facilitate Their Dissolution and to Improve Their Functionalization. <i>Pharmaceutics</i> , 2020, 12, 421.	2.0	13
16	A Signaling Crosstalk between BMP9 and HGF/c-Met Regulates Mouse Adult Liver Progenitor Cell Survival. <i>Cells</i> , 2020, 9, 752.	1.8	10
17	Case Report: An EGFR-Targeted 4-1BB-agonistic Trimerbody Does Not Induce Hepatotoxicity in Transgenic Mice With Liver Expression of Human EGFR. <i>Frontiers in Immunology</i> , 2020, 11, 614363.	2.2	5
18	Calcium Regulates HCC Proliferation as well as EGFR Recycling/Degradation and Could Be a New Therapeutic Target in HCC. <i>Cancers</i> , 2019, 11, 1588.	1.7	6

#	ARTICLE	IF	CITATIONS
19	Downregulation of Epidermal Growth Factor Receptor in hepatocellular carcinoma facilitates Transforming Growth Factor- β -induced epithelial to amoeboid transition. <i>Cancer Letters</i> , 2019, 464, 15-24.	3.2	25
20	Paradoxical role of the NADPH oxidase NOX4 in early preneoplastic stages of hepatocytes induced by amino acid deprivation. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2019, 1863, 714-722.	1.1	4
21	c-Met Signaling Is Essential for Mouse Adult Liver Progenitor Cells Expansion After Transforming Growth Factor- β -Induced Epithelial to Mesenchymal Transition and Regulates Cell Phenotypic Switch. <i>Stem Cells</i> , 2019, 37, 1108-1118.	1.4	19
22	Snail mediates crosstalk between TGF β 2 and LXR \pm in hepatocellular carcinoma. <i>Cell Death and Differentiation</i> , 2018, 25, 885-903.	5.0	34
23	Redox stress in Marfan syndrome: Dissecting the role of the NADPH oxidase NOX4 in aortic aneurysm. <i>Free Radical Biology and Medicine</i> , 2018, 118, 44-58.	1.3	57
24	Galunisertib suppresses the staminal phenotype in hepatocellular carcinoma by modulating CD44 expression. <i>Cell Death and Disease</i> , 2018, 9, 373.	2.7	31
25	Altered TGF- β 2 endocytic trafficking contributes to the increased signaling in Marfan syndrome. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2018, 1864, 554-562.	1.8	16
26	Transforming Growth Factor- β -Induced Cell Plasticity in Liver Fibrosis and Hepatocarcinogenesis. <i>Frontiers in Oncology</i> , 2018, 8, 357.	1.3	243
27	Revisiting the liver: from development to regeneration - what we ought to know!. <i>International Journal of Developmental Biology</i> , 2018, 62, 441-451.	0.3	14
28	TGF- β 2 and the Tissue Microenvironment: Relevance in Fibrosis and Cancer. <i>International Journal of Molecular Sciences</i> , 2018, 19, 1294.	1.8	231
29	Bone morphogenetic protein 9 as a key regulator of liver progenitor cells in DDC-induced cholestatic liver injury. <i>Liver International</i> , 2018, 38, 1664-1675.	1.9	26
30	Transforming growth factor- β -induced plasticity causes a migratory stemness phenotype in hepatocellular carcinoma. <i>Cancer Letters</i> , 2017, 392, 39-50.	3.2	69
31	The NADPH oxidase NOX4 represses epithelial to amoeboid transition and efficient tumour dissemination. <i>Oncogene</i> , 2017, 36, 3002-3014.	2.6	57
32	European contribution to the study of ROS: A summary of the findings and prospects for the future from the COST action BM1203 (EU-ROS). <i>Redox Biology</i> , 2017, 13, 94-162.	3.9	242
33	Role of the Transforming Growth Factor- β 2 in regulating hepatocellular carcinoma oxidative metabolism. <i>Scientific Reports</i> , 2017, 7, 12486.	1.6	54
34	Exploring liver physiology, pathology, TGF- β 2, EMT, stemness and new developments in liver cancer. <i>Hepatic Oncology</i> , 2017, 4, 9-13.	4.2	5
35	Hybrid polymeric-protein nano-carriers (HPPNC) for targeted delivery of TGF β 2 inhibitors to hepatocellular carcinoma cells. <i>Journal of Materials Science: Materials in Medicine</i> , 2017, 28, 120.	1.7	26
36	The level of caveolin-1 expression determines response to TGF- β 2 as a tumour suppressor in hepatocellular carcinoma cells. <i>Cell Death and Disease</i> , 2017, 8, e3098-e3098.	2.7	25

#	ARTICLE	IF	CITATIONS
37	The TGF- β pathway: a pharmacological target in hepatocellular carcinoma?. <i>Hepatic Oncology</i> , 2017, 4, 35-38.	4.2	2
38	Resminostat induces changes in epithelial plasticity of hepatocellular carcinoma cells and sensitizes them to sorafenib-induced apoptosis. <i>Oncotarget</i> , 2017, 8, 110367-110379.	0.8	26
39	TGF- β 1 and TGF- β 2 abundance in liver diseases of mice and men. <i>Oncotarget</i> , 2016, 7, 19499-19518.	0.8	52
40	New Insights into the Crossroads between EMT and Stemness in the Context of Cancer. <i>Journal of Clinical Medicine</i> , 2016, 5, 37.	1.0	110
41	Caveolin-1 dependent activation of the metalloprotease TACE/ADAM17 by TGF- β in hepatocytes requires activation of Src and the NADPH oxidase NOX1. <i>FEBS Journal</i> , 2016, 283, 1300-1310.	2.2	21
42	Dissecting the role of epidermal growth factor receptor catalytic activity during liver regeneration and hepatocarcinogenesis. <i>Hepatology</i> , 2016, 63, 604-619.	3.6	47
43	TGF- β signalling and liver disease. <i>FEBS Journal</i> , 2016, 283, 2219-2232.	2.2	457
44	The rationale for targeting TGF- β in chronic liver diseases. <i>European Journal of Clinical Investigation</i> , 2016, 46, 349-361.	1.7	60
45	Apoptosis in liver carcinogenesis and chemotherapy. <i>Hepatic Oncology</i> , 2015, 2, 381-397.	4.2	13
46	BMP9-Induced Survival Effect in Liver Tumor Cells Requires p38MAPK Activation. <i>International Journal of Molecular Sciences</i> , 2015, 16, 20431-20448.	1.8	22
47	Vascular Smooth Muscle Cell Phenotypic Changes in Patients With Marfan Syndrome. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2015, 35, 960-972.	1.1	116
48	Mechanisms regulating cell membrane localization of the chemokine receptor CXCR4 in human hepatocarcinoma cells. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2015, 1853, 1205-1218.	1.9	18
49	Role of NADPH oxidases in the redox biology of liver fibrosis. <i>Redox Biology</i> , 2015, 6, 106-111.	3.9	127
50	TGF- β in Hepatic Stellate Cell Activation and Liver Fibrogenesis: Updated. <i>Current Pathobiology Reports</i> , 2015, 3, 291-305.	1.6	36
51	A mesenchymal-like phenotype and expression of CD44 predict lack of apoptotic response to sorafenib in liver tumor cells. <i>International Journal of Cancer</i> , 2015, 136, E161-72.	2.3	108
52	Cross-Talk Between TGF- β and NADPH Oxidases During Liver Fibrosis and Hepatocarcinogenesis. <i>Current Pharmaceutical Design</i> , 2015, 21, 5964-5976.	0.9	31
53	A Trifluorinated Thiazoline Scaffold Leading to Pro-apoptotic Agents Targeting Prohibitins. <i>Angewandte Chemie - International Edition</i> , 2014, 53, 10150-10154.	7.2	35
54	The NADPH oxidase NOX4 inhibits hepatocyte proliferation and liver cancer progression. <i>Free Radical Biology and Medicine</i> , 2014, 69, 338-347.	1.3	78

#	ARTICLE	IF	CITATIONS
55	Po1 ^{1/4} Deficiency Increases Resistance to Oxidative Damage and Delays Liver Aging. PLoS ONE, 2014, 9, e93074.	1.1	6
56	TGF-beta Signaling in Cancer Treatment. Current Pharmaceutical Design, 2014, 20, 2934-2947.	0.9	155
57	Role of the tissue microenvironment as a therapeutic target in hepatocellular carcinoma. World Journal of Gastroenterology, 2014, 20, 4128.	1.4	34
58	Overactivation of the TGF- β 2 pathway confers a mesenchymal-like phenotype and CXCR4-dependent migratory properties to liver tumor cells. Hepatology, 2013, 58, 2032-2044.	3.6	113
59	Mouse Hepatic Oval Cells Require Met-Dependent PI3K to Impair TGF- β 2-Induced Oxidative Stress and Apoptosis. PLoS ONE, 2013, 8, e53108.	1.1	26
60	Differential Inhibition of the TGF- β 2 Signaling Pathway in HCC Cells Using the Small Molecule Inhibitor LY2157299 and the D10 Monoclonal Antibody against TGF- β 2 Receptor Type II. PLoS ONE, 2013, 8, e67109.	1.1	86
61	The Transforming Growth Factor-Beta (TGF- β 2) in Liver Fibrosis. , 2013, , 255-277.		1
62	BMP9 Is a Proliferative and Survival Factor for Human Hepatocellular Carcinoma Cells. PLoS ONE, 2013, 8, e69535.	1.1	67
63	Protein-tyrosine Phosphatase 1B (PTP1B) Deficiency Confers Resistance to Transforming Growth Factor- β 2 (TGF- β 2)-induced Suppressor Effects in Hepatocytes. Journal of Biological Chemistry, 2012, 287, 15263-15274.	1.6	25
64	BMPS and Liver: More Questions than Answers. Current Pharmaceutical Design, 2012, 18, 4114-4125.	0.9	17
65	Cell Fusion Reprogramming Leads to a Specific Hepatic Expression Pattern during Mouse Bone Marrow Derived Hepatocyte Formation In Vivo. PLoS ONE, 2012, 7, e33945.	1.1	13
66	Lack of amino acids in mouse hepatocytes in culture induces the selection of preneoplastic cells. Cellular Signalling, 2012, 24, 325-332.	1.7	5
67	EGFR is dispensable for c-Met-mediated proliferation and survival activities in mouse adult liver oval cells. Cellular Signalling, 2012, 24, 505-513.	1.7	15
68	Lack of aminoacids in mouse hepatocytes in culture induces the selection of preneoplastic cells. BMC Proceedings, 2012, 6, .	1.8	0
69	Sorafenib sensitizes hepatocellular carcinoma cells to physiological apoptotic stimuli. Journal of Cellular Physiology, 2012, 227, 1319-1325.	2.0	66
70	NADPH Oxidase NOX4 Mediates Stellate Cell Activation and Hepatocyte Cell Death during Liver Fibrosis Development. PLoS ONE, 2012, 7, e45285.	1.1	134
71	ROS Production Is Essential for the Apoptotic Function of E2F1 in Pheochromocytoma and Neuroblastoma Cell Lines. PLoS ONE, 2012, 7, e51544.	1.1	10
72	Dissecting the effect of targeting the epidermal growth factor receptor on TGF- β 2-induced-apoptosis in human hepatocellular carcinoma cells. Journal of Hepatology, 2011, 55, 351-358.	1.8	48

#	ARTICLE	IF	CITATIONS
73	Reciprocal regulation of NADPH oxidases and the cyclooxygenase-2 pathway. <i>Free Radical Biology and Medicine</i> , 2011, 51, 1789-1798.	1.3	44
74	The NADPH oxidase inhibitor VAS2870 impairs cell growth and enhances TGF- β -induced apoptosis of liver tumor cells. <i>Biochemical Pharmacology</i> , 2011, 81, 917-924.	2.0	44
75	The tyrphostin AG1478 inhibits proliferation and induces death of liver tumor cells through EGF receptor-dependent and independent mechanisms. <i>Biochemical Pharmacology</i> , 2011, 82, 1583-1592.	2.0	13
76	The transforming growth factor- β (TGF- β) mediates acquisition of a mesenchymal stem cell-like phenotype in human liver cells. <i>Journal of Cellular Physiology</i> , 2011, 226, 1214-1223.	2.0	92
77	Snail1 suppresses TGF- β -induced apoptosis and is sufficient to trigger EMT in hepatocytes. <i>Journal of Cell Science</i> , 2010, 123, 3467-3477.	1.2	134
78	NADPH Oxidase NOX1 Controls Autocrine Growth of Liver Tumor Cells through Up-regulation of the Epidermal Growth Factor Receptor Pathway. <i>Journal of Biological Chemistry</i> , 2010, 285, 24815-24824.	1.6	65
79	Growth factor- and cytokine-driven pathways governing liver stemness and differentiation. <i>World Journal of Gastroenterology</i> , 2010, 16, 5148.	1.4	37
80	Dysregulation of apoptosis in hepatocellular carcinoma cells. <i>World Journal of Gastroenterology</i> , 2009, 15, 513.	1.4	241
81	Overactivation of the MEK/ERK Pathway in Liver Tumor Cells Confers Resistance to TGF- β -Induced Cell Death through Impairing Up-regulation of the NADPH Oxidase NOX4. <i>Cancer Research</i> , 2009, 69, 7595-7602.	0.4	106
82	The inhibition of the epidermal growth factor (EGF) pathway enhances TGF- β -induced apoptosis in rat hepatoma cells through inducing oxidative stress coincident with a change in the expression pattern of the NADPH oxidases (NOX) isoforms. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2009, 1793, 253-263.	1.9	76
83	Role of CXCR4/SDF-1 α in the migratory phenotype of hepatoma cells that have undergone epithelial-mesenchymal transition in response to the transforming growth factor- β . <i>Cellular Signalling</i> , 2009, 21, 1595-1606.	1.7	68
84	Genetically modified animal models recapitulating molecular events altered in human hepatocarcinogenesis. <i>Clinical and Translational Oncology</i> , 2009, 11, 208-214.	1.2	7
85	Isolation and characterization of a putative liver progenitor population after treatment of fetal rat hepatocytes with TGF- β . <i>Journal of Cellular Physiology</i> , 2008, 215, 846-855.	2.0	21
86	Inhibition of the EGF receptor blocks autocrine growth and increases the cytotoxic effects of doxorubicin in rat hepatoma cells. <i>Biochemical Pharmacology</i> , 2008, 75, 1935-1945.	2.0	38
87	Upregulation of the NADPH oxidase NOX4 by TGF- β in hepatocytes is required for its pro-apoptotic activity. <i>Journal of Hepatology</i> , 2008, 49, 965-976.	1.8	197
88	Deletion of the Met Tyrosine Kinase in Liver Progenitor Oval Cells Increases Sensitivity to Apoptosis in Vitro. <i>American Journal of Pathology</i> , 2008, 172, 1238-1247.	1.9	30
89	Activation of NADPH oxidase by transforming growth factor- β in hepatocytes mediates up-regulation of epidermal growth factor receptor ligands through a nuclear factor- κ B-dependent mechanism. <i>Biochemical Journal</i> , 2007, 405, 251-259.	1.7	97
90	TGF- β dependent regulation of oxygen radicals during transdifferentiation of activated hepatic stellate cells to myofibroblastoid cells. <i>Comparative Hepatology</i> , 2007, 6, 1.	0.9	57

#	ARTICLE	IF	CITATIONS
91	Survival and apoptosis: a dysregulated balance in liver cancer. <i>Liver International</i> , 2007, 27, 155-162.	1.9	197
92	Apoptotic action of E2F1 requires glycogen synthase kinase 3- β activity in PC12 cells. <i>Journal of Neurochemistry</i> , 2007, 102, 2020-2028.	2.1	10
93	Differential intracellular signalling induced by TGF- β 2 in rat adult hepatocytes and hepatoma cells: Implications in liver carcinogenesis. <i>Cellular Signalling</i> , 2007, 19, 683-694.	1.7	84
94	Autocrine production of TGF- β 1 confers resistance to apoptosis after an epithelial-mesenchymal transition process in hepatocytes: Role of EGF receptor ligands. <i>Experimental Cell Research</i> , 2006, 312, 2860-2871.	1.2	65
95	Syndecan-2 expression increases serum-withdrawal-induced apoptosis, mediated by re-distribution of Fas into lipid rafts, in stably transfected Swiss 3T3 cells. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2006, 11, 2065-2075.	2.2	5
96	EGF blocks NADPH oxidase activation by TGF- β 2 in fetal rat hepatocytes, impairing oxidative stress, and cell death. <i>Journal of Cellular Physiology</i> , 2006, 207, 322-330.	2.0	70
97	Hematopoietic mobilization in mice increases the presence of bone marrow-derived hepatocytes via in vivo cell fusion. <i>Hepatology</i> , 2006, 43, 108-116.	3.6	120
98	Efficient execution of cell death in non-glycolytic cells requires the generation of ROS controlled by the activity of mitochondrial H ⁺ -ATP synthase. <i>Carcinogenesis</i> , 2006, 27, 925-935.	1.3	91
99	Involvement of EGF receptor and c-Src in the survival signals induced by TGF- β 1 in hepatocytes. <i>Oncogene</i> , 2005, 24, 4580-4587.	2.6	135
100	c-Myc regulates cell size and ploidy but is not essential for postnatal proliferation in liver. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 7286-7291.	3.3	85
101	Snail blocks the cell cycle and confers resistance to cell death. <i>Genes and Development</i> , 2004, 18, 1131-1143.	2.7	738
102	Source of early reactive oxygen species in the apoptosis induced by transforming growth factor- β 2 in fetal rat hepatocytes. <i>Free Radical Biology and Medicine</i> , 2004, 36, 16-26.	1.3	127
103	IRS-2 mediates the antiapoptotic effect of insulin in neonatal hepatocytes. <i>Hepatology</i> , 2004, 40, 1285-1294.	3.6	55
104	Resistance to TGF- β 2-induced apoptosis in regenerating hepatocytes. <i>Journal of Cellular Physiology</i> , 2004, 201, 385-392.	2.0	23
105	Transforming growth factor-beta activates both pro-apoptotic and survival signals in fetal rat hepatocytes. <i>Experimental Cell Research</i> , 2004, 292, 209-218.	1.2	61
106	Increased Generation of Hepatocytes Expressing Bone Marrow-Derived Markers after Hematopoietic Progenitors TM Mobilization in a Murine Model of Hepatic Damage. <i>Blood</i> , 2004, 104, 3600-3600.	0.6	0
107	Vitamin E blocks early events induced by 1-methyl-4-phenylpyridinium (MPP ⁺) in cerebellar granule cells. <i>Journal of Neurochemistry</i> , 2003, 84, 305-315.	2.1	44
108	Molecular Mechanisms of Insulin Resistance in IRS-2-Deficient Hepatocytes. <i>Diabetes</i> , 2003, 52, 2239-2248.	0.3	136

#	ARTICLE	IF	CITATIONS
109	Long-Term Treatment with Insulin Induces Apoptosis in Brown Adipocytes: Role of Oxidative Stress. <i>Endocrinology</i> , 2003, 144, 5390-5401.	1.4	19
110	clAP-1, but not XIAP, is cleaved by caspases during the apoptosis induced by TGF- β 2 in fetal rat hepatocytes. <i>FEBS Letters</i> , 2002, 520, 93-96.	1.3	29
111	Liver cell proliferation requires methionine adenosyltransferase 2A mRNA up-regulation. <i>Hepatology</i> , 2002, 35, 1381-1391.	3.6	38
112	The epithelial mesenchymal transition confers resistance to the apoptotic effects of transforming growth factor Beta in fetal rat hepatocytes. <i>Molecular Cancer Research</i> , 2002, 1, 68-78.	1.5	172
113	Short-chain ceramide regulates hepatic methionine adenosyltransferase expression. <i>Journal of Hepatology</i> , 2001, 34, 192-201.	1.8	13
114	Activation of p38MAPK by TGF- β 2 in fetal rat hepatocytes requires radical oxygen production, but is dispensable for cell death. <i>FEBS Letters</i> , 2001, 499, 225-229.	1.3	38
115	Functional pleiotropy of an intramolecular triplex-forming fragment from the 3'-UTR of the rat Pigrgene. <i>Physiological Genomics</i> , 2001, 5, 53-65.	1.0	13
116	Activation of caspases occurs downstream from radical oxygen species production, Bcl-xL down-regulation, and early cytochrome C release in apoptosis induced by transforming growth factor β 2 in rat fetal hepatocytes. <i>Hepatology</i> , 2001, 34, 548-556.	3.6	110
117	Reactive oxygen species (ROS) mediates the mitochondrial-dependent apoptosis induced by transforming growth factor β 2 in fetal hepatocytes. <i>FASEB Journal</i> , 2001, 15, 741-751.	0.2	288
118	Epidermal Growth Factor Impairs the Cytochrome C/Caspase-3 Apoptotic Pathway Induced by Transforming Growth Factor β 2 in Rat Fetal Hepatocytes Via a Phosphoinositide 3-Kinase-Dependent Pathway. <i>Hepatology</i> , 2000, 32, 528-535.	3.6	76
119	Fibronectin regulates morphology, cell organization and gene expression of rat fetal hepatocytes in primary culture. <i>Journal of Hepatology</i> , 2000, 32, 242-250.	1.8	52
120	Glucocorticoid receptor down-Regulates c-jun amino terminal kinases induced by tumor necrosis factor α in fetal rat hepatocyte primary cultures. <i>Hepatology</i> , 1999, 29, 849-857.	3.6	24
121	Effects of growth and differentiation factors on the epithelial-mesenchymal transition in cultured neonatal rat hepatocytes. <i>Journal of Hepatology</i> , 1999, 31, 895-904.	1.8	45
122	Phorbol esters down-regulate alpha-fetoprotein gene expression without affecting growth in fetal hepatocytes in primary culture. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1998, 1402, 151-164.	1.9	10
123	Transforming Growth Factor- β 2 (TGF- β 2) and EGF Promote Cord-like Structures That Indicate Terminal Differentiation of Fetal Hepatocytes in Primary Culture. <i>Experimental Cell Research</i> , 1998, 242, 27-37.	1.2	22
124	Epidermal growth factor, but not hepatocyte growth factor, suppresses the apoptosis induced by transforming growth factor-beta in fetal hepatocytes in primary culture. <i>FEBS Letters</i> , 1996, 384, 14-18.	1.3	68
125	Apoptosis Induced by Transforming Growth Factor- β 2 in Fetal Hepatocyte Primary Cultures. <i>Journal of Biological Chemistry</i> , 1996, 271, 7416-7422.	1.6	248
126	Regulation of gene expression by interleukin-6 in fetal rat hepatocyte primary cultures: Role of epidermal growth factor and dexamethasone. <i>Hepatology</i> , 1995, 22, 1769-1775.	3.6	11

#	ARTICLE	IF	CITATIONS
127	Transforming growth factor β modulates growth and differentiation of fetal hepatocytes in primary culture. <i>Journal of Cellular Physiology</i> , 1995, 165, 398-405.	2.0	62
128	Noradrenergic modulation of albumin expression in growth-stimulated adult rat hepatocytes in primary culture. <i>Journal of Cellular Physiology</i> , 1994, 158, 513-517.	2.0	5
129	Growth stimulation of rat fetal hepatocytes in response to hepatocyte growth factor: modulation of c-myc and c-fos expression. <i>Biochemical and Biophysical Research Communications</i> , 1992, 189, 684-690.	1.0	47
130	Differential proliferative response of cultured fetal and regenerating hepatocytes to growth factors and hormones. <i>Experimental Cell Research</i> , 1992, 202, 495-500.	1.2	64
131	Regulation of albumin expression in fetal rat hepatocytes cultured under proliferative conditions: Role of epidermal growth factor and hormones. <i>Journal of Cellular Physiology</i> , 1992, 152, 95-101.	2.0	38
132	The role of NADPH in the regulation of glucose-6-phosphate and 6-phosphogluconate dehydrogenases in rat adipose tissue. <i>Molecular and Cellular Biochemistry</i> , 1991, 105, 1-5.	1.4	17
133	Possible involvement of NADPH requirement in regulation of Glucose-6-Phosphate and 6-Phosphogluconate dehydrogenase levels in rat liver. <i>Molecular and Cellular Biochemistry</i> , 1990, 95, 107-115.	1.4	8
134	[D-Arg1, D-Phe5, D-Trp7,9, Leu11] substance P, a neuropeptide antagonist, blocks binding, Ca ²⁺ -mobilizing, and mitogenic effects of endothelin and vasoactive intestinal contractor in mouse 3T3 cells. <i>Journal of Cellular Physiology</i> , 1990, 145, 88-94.	2.0	31
135	Solubilization of the bombesin receptor from Swiss 3T3 cell membranes. <i>FEBS Letters</i> , 1990, 263, 80-84.	1.3	30
136	Vasoactive intestinal contractor, a novel peptide, shares a common receptor with endothelin-1 and stimulates Ca ²⁺ mobilization and DNA synthesis in Swiss 3T3 cells. <i>Biochemical and Biophysical Research Communications</i> , 1990, 167, 161-167.	1.0	28
137	Rates of lipogenesis in fetal hepatocytes in suspension and in primary culture: hormonal effects. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1989, 1012, 320-324.	1.9	30
138	Precocious induction of malic enzyme by nutritional and hormonal factors in rat foetal hepatocyte primary cultures. <i>Biochemical and Biophysical Research Communications</i> , 1989, 161, 1028-1034.	1.0	13
139	Hormonal regulation of malic enzyme expression in primary cultures of foetal brown adipocytes. <i>Biochemical and Biophysical Research Communications</i> , 1989, 163, 341-347.	1.0	15
140	Inhibition of fatty acid biosynthesis by bezafibrate in different rat cells. <i>Biochemical Pharmacology</i> , 1989, 38, 2505-2510.	2.0	4
141	Induction of malic enzyme genetic expression in rat foetal hepatocyte primary cultures. <i>Biochemical Society Transactions</i> , 1989, 17, 172-173.	1.6	1
142	Fetal rat brown adipocyte primary cultures: characterization of a system for lipid synthesis studies. <i>Biochemical Society Transactions</i> , 1988, 16, 274-275.	1.6	1
143	Short-term control of the pentose phosphate cycle by insulin could be modulated by the NADPH/NADP ratio in rat adipocytes and hepatocytes. <i>Biochemical and Biophysical Research Communications</i> , 1987, 146, 920-925.	1.0	47
144	NADPH/NADP ratio could regulate the glyoxylate cycle in <i>Tetrahymena pyriformis</i> . <i>Comparative Biochemistry and Physiology Part B: Comparative Biochemistry</i> , 1987, 88, 851-854.	0.2	1

#	ARTICLE	IF	CITATIONS
145	The NADPH consumption regulates the NADPH-producing pathways (pentose phosphate cycle and malic) Tj ETQq1_1_0.784314 rgBT (0	1.4	15
146	The pentose phosphate cycle is regulated by NADPH/NADP ratio in rat liver. Archives of Biochemistry and Biophysics, 1985, 236, 110-118.	1.4	55
147	Interaction with protein SH groups could be involved in adriamycin cardiotoxicity. Biochemical Medicine, 1984, 32, 289-295.	0.5	9
148	Citrate synthase of Tetrahymena pyriformis: Evolutionary and regulatory aspects. Archives of Biochemistry and Biophysics, 1983, 220, 354-360.	1.4	5