

Mark Ashworth

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/724902/mark-ashworth-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

182
papers

3,574
citations

33
h-index

52
g-index

209
ext. papers

4,378
ext. citations

3.2
avg, IF

5.5
L-index

#	Paper	IF	Citations
182	Inequalities in developing multimorbidity over time: A population-based cohort study from an urban, multi-ethnic borough in the United Kingdom.. <i>Lancet Regional Health - Europe, The</i> , 2022 , 12, 100247		1
181	Osteoporosis referral and treatment among people with severe mental illness: A ten-year data linkage study.. <i>Journal of Psychiatric Research</i> , 2022 , 147, 94-102	5.2	
180	Multimorbidity clusters among people with serious mental illness: a representative primary and secondary data linkage cohort study.. <i>Psychological Medicine</i> , 2022 , 1-12	6.9	1
179	What factors influence differential uptake of NHS Health Checks, diabetes and hypertension reviews among women in ethnically diverse South London? Cross-sectional analysis of 63,000 primary care records. <i>EclinicalMedicine</i> , 2022 , 49, 101471	11.3	
178	Impact of primary care funding on patient satisfaction: a retrospective longitudinal study of English general practice, 2013-2016. <i>British Journal of General Practice</i> , 2021 , 71, e47-e54	1.6	1
177	Hypertension and cardiovascular risk factor management in a multi-ethnic cohort of adults with CKD: a cross sectional study in general practice. <i>Journal of Nephrology</i> , 2021 , 1	4.8	0
176	Access to mental healthcare in the year after first fit note: a longitudinal study of linked clinical records. <i>BMJ Open</i> , 2021 , 11, e044725	3	
175	Impact of multimorbidity on healthcare costs and utilisation: a systematic review of the UK literature. <i>British Journal of General Practice</i> , 2021 , 71, e39-e46	1.6	20
174	Identifying multimorbidity clusters with the highest primary care use: 15 years of evidence from a multi-ethnic metropolitan population. <i>British Journal of General Practice</i> , 2021 ,	1.6	2
173	Epilepsy and mortality: a retrospective cohort analysis with a nested case-control study identifying causes and risk factors from primary care and linkage-derived data. <i>BMJ Open</i> , 2021 , 11, e052841	3	0
172	Health condition at first fit note and number of fit notes: a longitudinal study of primary care records in south London. <i>BMJ Open</i> , 2021 , 11, e043889	3	
171	Identifying longitudinal clusters of multimorbidity in an urban setting: A population-based cross-sectional study. <i>Lancet Regional Health - Europe, The</i> , 2021 , 3, 100047		7
170	Spatio-temporal associations of air pollutant concentrations, GP respiratory consultations and respiratory inhaler prescriptions: a 5-year study of primary care in the borough of Lambeth, South London. <i>Environmental Health</i> , 2021 , 20, 54	6	2
169	Safety of reducing antibiotic prescribing in primary care: a mixed-methods study. <i>Health Services and Delivery Research</i> , 2021 , 9, 1-126	1.5	2
168	Adapting the definition of multimorbidity - development of a locality-based consensus for selecting included Long Term Conditions. <i>BMC Family Practice</i> , 2021 , 22, 124	2.6	4
167	Cancer detection via primary care urgent referral and association with practice characteristics: a retrospective cross-sectional study in England from 2009/2010 to 2018/2019. <i>British Journal of General Practice</i> , 2021 , 71, e826-e835	1.6	3
166	Evaluation of NHS Practitioner Health: capturing mental health outcomes using five instruments. <i>BJPsych Open</i> , 2021 , 7, e106	5	0

165	Comment on: Influence of prior antibiotic use on risk of rheumatoid arthritis: case-control study in general practice. Reply. <i>Rheumatology</i> , 2021 , 60, e150	3.9	
164	Prevalence and predictors of potentially inappropriate prescribing in middle-aged adults: a repeated cross-sectional study. <i>British Journal of General Practice</i> , 2021 , 71, e491-e497	1.6	3
163	Development and validation of the Maudsley Modified Patient Health Questionnaire (MM-PHQ-9). <i>BJPsych Open</i> , 2021 , 7, e123	5	0
162	Inequalities in glycemic management in people living with type 2 diabetes mellitus and severe mental illnesses: cohort study from the UK over 10 years. <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9,	4.5	1
161	Applying resolved and remission codes reduced prevalence of multimorbidity in an urban multi-ethnic population. <i>Journal of Clinical Epidemiology</i> , 2021 , 140, 135-148	5.7	1
160	PRINCE Secondary: transdiagnostic cognitive behaviour therapy for persistent physical symptoms. <i>Psychological Medicine</i> , 2021 , 1-2	6.9	
159	A case-control and cohort study to determine the relationship between ethnic background and severe COVID-19. <i>EClinicalMedicine</i> , 2020 , 28, 100574	11.3	26
158	Multimorbidity and fit note receipt in working-age adults with long-term health conditions. <i>Psychological Medicine</i> , 2020 , 1-10	6.9	3
157	Association between use of urgent suspected cancer referral and mortality and stage at diagnosis: a 5-year national cohort study. <i>British Journal of General Practice</i> , 2020 , 70, e389-e398	1.6	20
156	Is health research undertaken where the burden of disease is greatest? Observational study of geographical inequalities in recruitment to research in England 2013-2018. <i>BMC Medicine</i> , 2020 , 18, 133	11.4	7
155	Study protocol for the antidepressant advisor (ADeSS): a decision support system for antidepressant treatment for depression in UK primary care: a feasibility study. <i>BMJ Open</i> , 2020 , 10, e035905	3	5
154	Demographic variation in fit note receipt and long-term conditions in south London. <i>Occupational and Environmental Medicine</i> , 2020 , 77, 418-426	2.1	9
153	Risks of use and non-use of antibiotics in primary care: qualitative study of prescribers' views. <i>BMJ Open</i> , 2020 , 10, e038851	3	5
152	Reducing weight and increasing physical activity in people at high risk of cardiovascular disease: a randomised controlled trial comparing the effectiveness of enhanced motivational interviewing intervention with usual care. <i>Heart</i> , 2020 , 106, 447-454	5.1	11
151	Ethics education and moral decision-making in clinical commissioning: an interview study. <i>British Journal of General Practice</i> , 2020 , 70, e45-e54	1.6	1
150	The association between colorectal cancer and prior antibiotic prescriptions: case control study. <i>British Journal of Cancer</i> , 2020 , 122, 912-917	8.7	17
149	Proportion of young people in the general population consulting general practitioners: Potential for mental health screening and prevention. <i>Microbial Biotechnology</i> , 2020 , 14, 631-635	3.3	4
148	Lean implementation within healthcare: imaging as fertile ground. <i>Journal of Health Organization and Management</i> , 2020 , ahead-of-print,	1.9	1

147	Uncoded chronic kidney disease in primary care: a cross-sectional study of inequalities and cardiovascular disease risk management. <i>British Journal of General Practice</i> , 2020 , 70, e785-e792	1.6	3
146	BMC family practice integrated GP care for patients with persistent physical symptoms: feasibility cluster randomised trial. <i>BMC Family Practice</i> , 2020 , 21, 207	2.6	1
145	Probability of sepsis after infection consultations in primary care in the United Kingdom in 2002-2017: Population-based cohort study and decision analytic model. <i>PLoS Medicine</i> , 2020 , 17, e1003202	11.6	12
144	Personalising Outcome Measurement in Substance Misuse Treatment: the Feasibility and Psychometrics of Two Individualised Outcome Measures. <i>International Journal of Mental Health and Addiction</i> , 2020 , 1	8.8	1
143	Peritonsillar Abscess and Antibiotic Prescribing for Respiratory Infection in Primary Care: A Population-Based Cohort Study and Decision-Analytic Model. <i>Annals of Family Medicine</i> , 2020 , 18, 390-396	2.9	1
142	Prevalence of suspected COVID-19 infection in patients from ethnic minority populations: a cross-sectional study in primary care. <i>British Journal of General Practice</i> , 2020 , 70, e696-e704	1.6	7
141	Influence of prior antibiotic use on risk of rheumatoid arthritis: case control study in general practice. <i>Rheumatology</i> , 2020 , 59, 1281-1287	3.9	7
140	“Where Are Things I Want to Say But You Do Not Ask” a Comparison Between Standardised and Individualised Evaluations in Substance Use Treatment. <i>International Journal of Mental Health and Addiction</i> , 2020 , 18, 861-874	8.8	4
139	Serious bacterial infections and antibiotic prescribing in primary care: cohort study using electronic health records in the UK. <i>BMJ Open</i> , 2020 , 10, e036975	3	10
138	Persistent physical symptoms reduction intervention: a system change and evaluation in secondary care (PRINCE secondary) - a CBT-based transdiagnostic approach: study protocol for a randomised controlled trial. <i>BMC Psychiatry</i> , 2019 , 19, 307	4.2	8
137	Individualised or Standardised Outcome Measures: A Co-habitation?. <i>Administration and Policy in Mental Health and Mental Health Services Research</i> , 2019 , 46, 425-428	3.1	13
136	Content and source analysis of popular tweets following a recent case of diphtheria in Spain. <i>European Journal of Public Health</i> , 2019 , 29, 117-122	2.1	12
135	PSYCHLOPS in Polish primary care: how do clients conceptualise their problems on a patient-generated outcome measure?. <i>Heliyon</i> , 2019 , 5, e02209	3.6	0
134	User Involvement in the Design of a Data-Driven Self-Management Decision Support Tool for Stroke Survivors 2019 ,		2
133	Electronically delivered interventions to reduce antibiotic prescribing for respiratory infections in primary care: cluster RCT using electronic health records and cohort study. <i>Health Technology Assessment</i> , 2019 , 23, 1-70	4.4	11
132	Enhanced motivational interviewing for reducing weight and increasing physical activity in adults with high cardiovascular risk: the MOVE IT three-arm RCT. <i>Health Technology Assessment</i> , 2019 , 23, 1-144	4.4	7
131	Effectiveness and safety of electronically delivered prescribing feedback and decision support on antibiotic use for respiratory illness in primary care: REDUCE cluster randomised trial. <i>BMJ, The</i> , 2019 , 364, l236	5.9	54
130	Relationship between general practice capitation funding and the quality of primary care in England: a cross-sectional, 3-year study. <i>BMJ Open</i> , 2019 , 9, e030624	3	5

129	Persistent physical symptoms reduction intervention: a system change and evaluation (PRINCE)-integrated GP care for persistent physical symptoms: protocol for a feasibility and cluster randomised waiting list, controlled trial. <i>BMJ Open</i> , 2019 , 9, e025513	3	4
128	Journey to multimorbidity: longitudinal analysis exploring cardiovascular risk factors and sociodemographic determinants in an urban setting. <i>BMJ Open</i> , 2019 , 9, e031649	3	24
127	Is the rule of halves still relevant today? A cross-sectional analysis of hypertension detection, treatment and control in an urban community. <i>Journal of Hypertension</i> , 2019 , 37, 2470-2480	1.9	9
126	The development and testing of PSYCHLOPS Kids: a new child-centred outcome measure. <i>Child and Adolescent Mental Health</i> , 2019 , 24, 54-65	2.2	6
125	A Collaborative Decision Support Tool for Managing Chronic Conditions. <i>Studies in Health Technology and Informatics</i> , 2019 , 264, 644-648	0.5	2
124	Antimicrobial prescribing in post-graduate training practices: a cross-sectional study of prescribing data in general practices in England. <i>Education for Primary Care</i> , 2018 , 29, 139-143	0.9	4
123	Population and patient factors affecting emergency department attendance in London: retrospective cohort analysis of linked primary and secondary care records. <i>British Journal of General Practice</i> , 2018 , 68, e157-e167	1.6	11
122	Association between prior antibiotic therapy and subsequent risk of community-acquired infections: a systematic review. <i>Journal of Antimicrobial Chemotherapy</i> , 2018 , 73, 287-296	5.1	20
121	Enhanced Invitations Using the Question-Behavior Effect and Financial Incentives to Promote Health Check Uptake in Primary Care. <i>Annals of Behavioral Medicine</i> , 2018 , 52, 594-605	4.5	8
120	Cardiovascular risk at health checks performed opportunistically or following an invitation letter. Cohort study. <i>Journal of Public Health</i> , 2018 , 40, e151-e156	3.5	4
119	Automated recruitment and randomisation for an efficient randomised controlled trial in primary care. <i>Trials</i> , 2018 , 19, 341	2.8	4
118	The role of spoken language in cardiovascular health inequalities: a cross-sectional study of people with non-English language preference. <i>BJGP Open</i> , 2018 , 1, bjgpopen17X101241	3.1	1
117	Capturing and missing the patient's story through outcome measures: A thematic comparison of patient-generated items in PSYCHLOPS with CORE-OM and PHQ-9. <i>Health Expectations</i> , 2018 , 21, 615-619	3.7	15
116	Evaluation of the Hospital Anxiety and Depression Scale (HADS) in screening stroke patients for symptoms: Item Response Theory (IRT) analysis. <i>Journal of Affective Disorders</i> , 2018 , 228, 33-40	6.6	22
115	The CONSULT System 2018 ,		1
114	An integrated care programme in London: qualitative evaluation. <i>Journal of Integrated Care</i> , 2018 , 26, 296-308	0.7	8
113	Improving medication adherence in stroke survivors: the intervention development process. <i>BMC Health Services Research</i> , 2018 , 18, 772	2.9	2
112	Response bias to a randomised controlled trial of a lifestyle intervention in people at high risk of cardiovascular disease: a cross-sectional analysis. <i>BMC Public Health</i> , 2018 , 18, 1092	4.1	7

111	Type 2 diabetes mellitus in people with severe mental illness: inequalities by ethnicity and age. Cross-sectional analysis of 588 408 records from the UK. <i>Diabetic Medicine</i> , 2017 , 34, 916-924	3.5	27
110	Psychological Determinants of Medication Adherence in Stroke Survivors: a Systematic Review of Observational Studies. <i>Annals of Behavioral Medicine</i> , 2017 , 51, 833-845	4.5	23
109	Costs and Outcomes of Increasing Access to Bariatric Surgery: Cohort Study and Cost-Effectiveness Analysis Using Electronic Health Records. <i>Value in Health</i> , 2017 , 20, 85-92	3.3	58
108	National disparities in the relationship between antimicrobial resistance and antimicrobial consumption in Europe: an observational study in 29 countries-authors' response. <i>Journal of Antimicrobial Chemotherapy</i> , 2017 , 72, 3500	5.1	1
107	Impact of primary care funding on secondary care utilisation and patient outcomes: a retrospective cross-sectional study of English general practice. <i>British Journal of General Practice</i> , 2017 , 67, e792-e799 ^{1.6}		7
106	Does outcome measurement of treatment for substance use disorder reflect the personal concerns of patients? A scoping review of measures recommended in Europe. <i>Drug and Alcohol Dependence</i> , 2017 , 179, 299-308	4.9	16
105	Ethnic differences in smoking intensity and COPD risk: an observational study in primary care. <i>Npj Primary Care Respiratory Medicine</i> , 2017 , 27, 50	3.2	14
104	GP Training practices in England: a description of their unique features based on national data. <i>Education for Primary Care</i> , 2017 , 28, 313-318	0.9	6
103	National disparities in the relationship between antimicrobial resistance and antimicrobial consumption in Europe: an observational study in 29 countries. <i>Journal of Antimicrobial Chemotherapy</i> , 2017 , 72, 3199-3204	5.1	19
102	Financiamento para medicina de família na primeira década: vida após o QOF. <i>Revista Brasileira De Medicina De Família E Comunidade</i> , 2017 , 12, 1-6	0.3	1
101	Cardiovascular disease treatment among patients with severe mental illness: a data linkage study between primary and secondary care. <i>British Journal of General Practice</i> , 2016 , 66, e374-81	1.6	32
100	Safety of reduced antibiotic prescribing for self limiting respiratory tract infections in primary care: cohort study using electronic health records. <i>BMJ, The</i> , 2016 , 354, i3410	5.9	74
99	Electronically delivered, multicomponent intervention to reduce unnecessary antibiotic prescribing for respiratory infections in primary care: a cluster randomised trial using electronic health records-REDUCE Trial study original protocol. <i>BMJ Open</i> , 2016 , 6, e010892	3	9
98	Prostate-specific antigen testing in inner London general practices: are those at higher risk most likely to get tested?. <i>BMJ Open</i> , 2016 , 6, e011356	3	10
97	Long term condition morbidity in English general practice: a cross-sectional study using three composite morbidity measures. <i>BMC Family Practice</i> , 2016 , 17, 166	2.6	6
96	"It is not just about the alcohol": service users' views about individualised and standardised clinical assessment in a therapeutic community for alcohol dependence. <i>Substance Abuse Treatment, Prevention, and Policy</i> , 2016 , 11, 25	3.4	12
95	Ethnicity and cardiovascular health inequalities in people with severe mental illnesses: protocol for the E-CHASM study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016 , 51, 627-38	4.5	12
94	Antibiotic prescribing and patient satisfaction in primary care in England: cross-sectional analysis of national patient survey data and prescribing data. <i>British Journal of General Practice</i> , 2016 , 66, e40-6	1.6	72

93	Costs and outcomes of increasing access to bariatric surgery for obesity: cohort study and cost-effectiveness analysis using electronic health records. <i>Health Services and Delivery Research</i> , 2016 , 4, 1-120	1.5	10
92	Location of care for people with serious mental illness (LOCAPE): implications for service use and costs using a mixed-methods approach. <i>Health Services and Delivery Research</i> , 2016 , 4, 1-108	1.5	3
91	Enhanced invitation methods and uptake of health checks in primary care: randomised controlled trial and cohort study using electronic health records. <i>Health Technology Assessment</i> , 2016 , 20, 1-92	4.4	21
90	Does COPD risk vary by ethnicity? A retrospective cross-sectional study. <i>International Journal of COPD</i> , 2016 , 11, 739-46	3	26
89	Cervical and breast cancer screening uptake among women with serious mental illness: a data linkage study. <i>BMC Cancer</i> , 2016 , 16, 819	4.8	35
88	Associations between multimorbidity, healthcare utilisation and health status: evidence from 16 European countries. <i>Age and Ageing</i> , 2016 , 45, 431-5	3	153
87	The relationship between prior antimicrobial prescription and meningitis: a case-control study. <i>British Journal of General Practice</i> , 2016 , 66, e228-33	1.6	4
86	Authors' reply to Honeyford and colleagues and Roland. <i>BMJ, The</i> , 2015 , 350, h1681	5.9	
85	Comparing the effectiveness of an enhanced MOTivational intErviewing InTervention (MOVE IT) with usual care for reducing cardiovascular risk in high risk subjects: study protocol for a randomised controlled trial. <i>Trials</i> , 2015 , 16, 112	2.8	13
84	Alcohol use, socioeconomic deprivation and ethnicity in older people. <i>BMJ Open</i> , 2015 , 5, e007525	3	29
83	Nursing consultations and control of diabetes in general practice: a retrospective observational study. <i>British Journal of General Practice</i> , 2015 , 65, e642-8	1.6	18
82	Socioeconomic deprivation and accident and emergency attendances: cross-sectional analysis of general practices in England. <i>British Journal of General Practice</i> , 2015 , 65, e649-54	1.6	25
81	Challenges of the Pandemic Response in Primary Care during Pre-Vaccination Period: A Qualitative Study. <i>Israel Journal of Health Policy Research</i> , 2015 , 4, 32	1.7	15
80	Investigating the relationship between quality of primary care and premature mortality in England: a spatial whole-population study. <i>BMJ, The</i> , 2015 , 350, h904	5.9	63
79	Hypertension: a cross-sectional study of the role of multimorbidity in blood pressure control. <i>BMC Family Practice</i> , 2015 , 16, 98	2.6	9
78	Continued high rates of antibiotic prescribing to adults with respiratory tract infection: survey of 568 UK general practices. <i>BMJ Open</i> , 2014 , 4, e006245	3	152
77	Patterns of physical co-/multi-morbidity among patients with serious mental illness: a London borough-based cross-sectional study. <i>BMC Family Practice</i> , 2014 , 15, 117	2.6	33
76	Patient experience and the role of postgraduate GP training: a cross-sectional analysis of national Patient Survey data in England. <i>British Journal of General Practice</i> , 2014 , 64, e168-77	1.6	16

75	Dangerous Ideas: resistant superbugs and the potential of the microbiome. <i>British Journal of General Practice</i> , 2014 , 64, 637	1.6	
74	Using a patient-generated mental-health measure 'PSYCHLOPS' to explore problems in patients with coronary heart disease. <i>British Journal of General Practice</i> , 2014 , 64, e354-63	1.6	5
73	Process evaluation of a point-of-care cluster randomised trial using a computer-delivered intervention to reduce antibiotic prescribing in primary care. <i>BMC Health Services Research</i> , 2014 , 14, 594	2.9	19
72	Variations in statin prescribing for primary cardiovascular disease prevention: cross-sectional analysis. <i>BMC Health Services Research</i> , 2014 , 14, 414	2.9	21
71	User-generated quality standards for youth mental health in primary care: a participatory research design using mixed methods. <i>BMJ Quality and Safety</i> , 2014 , 23, 857-66	5.4	15
70	Incidence of type 2 diabetes after bariatric surgery: population-based matched cohort study. <i>Lancet Diabetes and Endocrinology</i> , 2014 , 2, 963-8	18.1	53
69	Point-of-care cluster randomized trial in stroke secondary prevention using electronic health records. <i>Stroke</i> , 2014 , 45, 2066-71	6.7	20
68	Electronic health records for intervention research: a cluster randomized trial to reduce antibiotic prescribing in primary care (eCRT study). <i>Annals of Family Medicine</i> , 2014 , 12, 344-51	2.9	56
67	Enhancing the patient involvement in outcomes: a study protocol of personalised outcome measurement in the treatment of substance misuse. <i>BMC Psychiatry</i> , 2013 , 13, 337	4.2	10
66	Ethnicity and stroke risk in patients with atrial fibrillation. <i>Heart</i> , 2013 , 99, 1087-92	5.1	23
65	The Public Health Impact score: a new measure of public health effectiveness for general practices in England. <i>British Journal of General Practice</i> , 2013 , 63, e291-9	1.6	8
64	Influenza pandemic 2009/A/H1N1 management policies in primary care: a comparative analysis of three countries. <i>Australian Health Review</i> , 2013 , 37, 291-9	1.8	6
63	Trends in antiepileptic drug utilisation in UK primary care 1993-2008: cohort study using the General Practice Research Database. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2012 , 21, 466-70	3.2	74
62	Quality standards for child and adolescent mental health in primary care. <i>BMC Family Practice</i> , 2012 , 13, 51	2.6	11
61	A pilot randomised controlled trial of personalised care for depressed patients with symptomatic coronary heart disease in South London general practices: the UPBEAT-UK RCT protocol and recruitment. <i>BMC Psychiatry</i> , 2012 , 12, 58	4.2	11
60	Cluster randomized trial in the general practice research database: 2. Secondary prevention after first stroke (eCRT study): study protocol for a randomized controlled trial. <i>Trials</i> , 2012 , 13, 181	2.8	11
59	The quality and outcomes framework in the United kingdom: indicators in transition. <i>Journal of Ambulatory Care Management</i> , 2012 , 35, 200-5	0.8	2
58	Achievement of cholesterol targets and prescribing of higher-cost statins: a cross-sectional study in general practice. <i>British Journal of General Practice</i> , 2012 , 62, e815-20	1.6	1

57	Managing hypertension in general practice: a cross-sectional study of treatment and ethnicity. <i>British Journal of General Practice</i> , 2012 , 62, e703-9	1.6	7
56	The relationship between asthma and depression in primary care patients: a historical cohort and nested case control study. <i>PLoS ONE</i> , 2011 , 6, e20750	3.7	47
55	Measuring psychological change during cognitive behaviour therapy in primary care: a Polish study using 'PSYCHLOPS' (Psychological Outcome Profiles). <i>PLoS ONE</i> , 2011 , 6, e27378	3.7	16
54	Epilepsy mortality and risk factors for death in epilepsy: a population-based study. <i>British Journal of General Practice</i> , 2011 , 61, e271-8	1.6	71
53	Ethnic differences in blood pressure monitoring and control in south east London. <i>British Journal of General Practice</i> , 2011 , 61, 190-6	1.6	24
52	Up-beat UK: a programme of research into the relationship between coronary heart disease and depression in primary care patients. <i>BMC Family Practice</i> , 2011 , 12, 38	2.6	23
51	Utility of electronic patient records in primary care for stroke secondary prevention trials. <i>BMC Public Health</i> , 2011 , 11, 86	4.1	21
50	Cluster randomised trial in the General Practice Research Database: 1. Electronic decision support to reduce antibiotic prescribing in primary care (eCRT study). <i>Trials</i> , 2011 , 12, 115	2.8	23
49	Ethnic isolation and psychosis: re-examining the ethnic density effect. <i>Psychological Medicine</i> , 2011 , 41, 1263-9	6.9	52
48	Identifying poorly performing general practices in England: a longitudinal study using data from the quality and outcomes framework. <i>Journal of Health Services Research and Policy</i> , 2011 , 16, 21-7	2.4	22
47	Family practices' achievement of diabetes quality of care targets and risk of screen-detected diabetic retinopathy. <i>PLoS ONE</i> , 2010 , 5, e10424	3.7	4
46	Impact of pay for performance on inequalities in health care: systematic review. <i>Journal of Health Services Research and Policy</i> , 2010 , 15, 178-84	2.4	66
45	What underlies the perception that a medical intervention is effective? An exploratory study among smokers. <i>Nicotine and Tobacco Research</i> , 2010 , 12, 508-15	4.9	5
44	Are problems prevalent and stable in non-clinical populations? Problems and test-retest stability of a patient-generated measure, PSYCHLOPS (Psychological Outcome Profiles), in a non-clinical student sample. <i>British Journal of Guidance and Counselling</i> , 2010 , 38, 431-439	0.8	5
43	Differences in the primary care management of patients with psychosis from two ethnic groups: a population-based cross-sectional study. <i>Family Practice</i> , 2010 , 27, 439-46	1.9	13
42	Parental help-seeking in primary care for child and adolescent mental health concerns: qualitative study. <i>British Journal of Psychiatry</i> , 2010 , 197, 476-81	5.4	78
41	Nurse staffing and quality of care in UK general practice: cross-sectional study using routinely collected data. <i>British Journal of General Practice</i> , 2010 , 60, 36-48	1.6	47
40	Can a self-referral system help improve access to psychological treatments?. <i>British Journal of General Practice</i> , 2010 , 60, 365-71	1.6	22

39	Developing a computer delivered, theory based intervention for guideline implementation in general practice. <i>BMC Family Practice</i> , 2010 , 11, 90	2.6	47
38	Quality and Outcomes Framework: smoke and mirrors?. <i>Quality in Primary Care</i> , 2010 , 18, 127-31		6
37	Variations in anxiolytic and hypnotic prescribing by GPs: a cross-sectional analysis using data from the UK Quality and Outcomes Framework. <i>British Journal of General Practice</i> , 2009 , 59, e191-8	1.6	27
36	Which practices are high antibiotic prescribers? A cross-sectional analysis. <i>British Journal of General Practice</i> , 2009 , 59, e315-20	1.6	44
35	Selection of medical diagnostic codes for analysis of electronic patient records. Application to stroke in a primary care database. <i>PLoS ONE</i> , 2009 , 4, e7168	3.7	57
34	Measuring psychological outcomes after cognitive behaviour therapy in primary care: a comparison between a new patient-generated measure PSYCHLOPS (Psychological Outcome Profiles) and HADS (Hospital Anxiety and Depression Scale). <i>Journal of Mental Health</i> , 2009 , 18, 169-177	2.7	23
33	Selective decrease in consultations and antibiotic prescribing for acute respiratory tract infections in UK primary care up to 2006. <i>Journal of Public Health</i> , 2009 , 31, 512-20	3.5	59
32	Population intermediate outcomes of diabetes under pay-for-performance incentives in England from 2004 to 2008. <i>Diabetes Care</i> , 2009 , 32, 427-9	14.6	45
31	A tale of two systems: perceptions of primary care for depression in London and Melbourne. <i>Family Practice</i> , 2009 , 26, 210-20	1.9	6
30	Schizophrenia in black Caribbeans living in the UK: an exploration of underlying causes of the high incidence rate. <i>British Journal of General Practice</i> , 2008 , 58, 429-34	1.6	37
29	Effect of social deprivation on blood pressure monitoring and control in England: a survey of data from the quality and outcomes framework. <i>BMJ, The</i> , 2008 , 337, a2030	5.9	73
28	Quality improvement in UK primary care: the role of financial incentives. <i>Journal of Ambulatory Care Management</i> , 2008 , 31, 220-5	0.8	7
27	Ethnic density, physical illness, social deprivation and antidepressant prescribing in primary care: ecological study. <i>British Journal of Psychiatry</i> , 2008 , 193, 235-9	5.4	33
26	PSYCHLOPS is a psychometric outcome measure that is finding a niche. <i>Counselling and Psychotherapy Research</i> , 2007 , 7, 201-202	1.3	2
25	Social deprivation and statin prescribing: a cross-sectional analysis using data from the new UK general practitioner 'Quality and Outcomes Framework'. <i>Journal of Public Health</i> , 2007 , 29, 40-7	3.5	38
24	Developing primary care services for high-dose benzodiazepine-dependent patients: A consultation survey. <i>Drugs: Education, Prevention and Policy</i> , 2007 , 14, 429-442	1.2	3
23	Comparing population estimates for drug users, the prevalence of blood-borne viruses and the provision of services in one London borough: Evidence of a gap in provision?. <i>Drugs: Education, Prevention and Policy</i> , 2007 , 14, 205-215	1.2	
22	The relationship between social deprivation and the quality of primary care: a national survey using indicators from the UK Quality and Outcomes Framework. <i>British Journal of General Practice</i> , 2007 , 57, 441-8	1.6	55

21	The relationship between general practice characteristics and quality of care: a national survey of quality indicators used in the UK Quality and Outcomes Framework, 2004-5. <i>BMC Family Practice</i> , 2006 , 7, 68	2.6	72
20	The experiences of therapists using a new client-centred psychometric instrument, PSYCHLOPS (Psychological Outcome Profiles). <i>Counselling and Psychotherapy Research</i> , 2005 , 5, 37-42	1.3	10
19	Accuracy of the pressure scale of sphygmomanometers in clinical use within primary care. <i>Blood Pressure Monitoring</i> , 2005 , 10, 181-8	1.3	53
18	Which general practices have higher list inflation? An exploratory study. <i>Family Practice</i> , 2005 , 22, 529-31	1.9	21
17	What factors are associated with improvement after brief psychological interventions in primary care? Issues arising from using routine outcome measurement to inform clinical practice. <i>Counselling and Psychotherapy Research</i> , 2005 , 5, 273-280	1.3	10
16	Variations in antibiotic prescribing and consultation rates for acute respiratory infection in UK general practices 1995-2000. <i>British Journal of General Practice</i> , 2005 , 55, 603-8	1.6	100
15	PCT / primary care referrals: current and future trends. <i>British Journal of Health Care Management</i> , 2004 , 10, 110-115	0.4	
14	Linking research and development in primary care: primary care trusts, primary care research networks and primary care academics. <i>Primary Health Care Research and Development</i> , 2004 , 5, 255-263	1.6	2
13	Why has antibiotic prescribing for respiratory illness declined in primary care? A longitudinal study using the General Practice Research Database. <i>Journal of Public Health</i> , 2004 , 26, 268-74	3.5	103
12	How are primary care organizations using financial incentives to influence prescribing?. <i>Journal of Public Health</i> , 2004 , 26, 48-51	3.5	8
11	A client-generated psychometric instrument: The development of PSYCHLOPS. <i>Counselling and Psychotherapy Research</i> , 2004 , 4, 27-31	1.3	71
10	Perceptions of psychological content in the GP consultation--the role of practice, personal and prescribing attributes. <i>Family Practice</i> , 2003 , 20, 373-5	1.9	7
9	The development of prescribing incentive schemes in primary care: a longitudinal survey. <i>British Journal of General Practice</i> , 2003 , 53, 468-70	1.6	5
8	Partnership effects in general practice: identification of clustering using intra-class correlation coefficients. <i>British Journal of General Practice</i> , 2003 , 53, 863-5	1.6	
7	Satisfaction with respite care: a pilot study. <i>Health and Social Care in the Community</i> , 2002 , 10, 479-84	2.6	21
6	Benzodiazepines: addiction and abuse. <i>Drugs: Education, Prevention and Policy</i> , 2002 , 9, 389-397	1.2	
5	Demand, appropriateness and prescribing of 'lifestyle drugs': a consultation survey in general practice. <i>Family Practice</i> , 2002 , 19, 236-41	1.9	10
4	Prescribing incentive schemes in two NHS regions: cross sectional survey. <i>BMJ, The</i> , 2002 , 324, 1187-8	5.9	8

3	Psychiatric referral rates and the influence of on-site mental health workers in general practice. <i>British Journal of General Practice</i> , 2002 , 52, 39-41	1.6	7
2	'Time and space': carers' views about respite care. <i>Health and Social Care in the Community</i> , 2000 , 8, 50-56.6	6.6	62
1	Efficacy of therapist-delivered transdiagnostic CBT for patients with persistent physical symptoms in secondary care: a randomised controlled trial. <i>Psychological Medicine</i> , 1-11	6.9	2