

Hirofumi Taki

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7250306/publications.pdf>

Version: 2024-02-01

11
papers

159
citations

1478505

6
h-index

1281871

11
g-index

11
all docs

11
docs citations

11
times ranked

293
citing authors

#	ARTICLE	IF	CITATIONS
1	Physiologic Target, Molecular Evolution, and Pathogenic Functions of a Monoclonal Anti-“Citrullinated Protein Antibody Obtained From a Patient With Rheumatoid Arthritis. <i>Arthritis and Rheumatology</i> , 2020, 72, 2040-2049.	5.6	6
2	Autoantibodies reactive to PEP08 are clinically related with morbidity and severity of interstitial lung disease in connective tissue diseases. <i>European Journal of Immunology</i> , 2018, 48, 1717-1727.	2.9	6
3	Monoclonal Antibody Against Citrullinated Peptides Obtained From Rheumatoid Arthritis Patients Reacts With Numerous Citrullinated Microbial and Food Proteins. <i>Arthritis and Rheumatology</i> , 2015, 67, 2020-2031.	5.6	29
4	Exacerbation of symptoms in Blau syndrome/early-onset sarcoidosis following delivery. <i>European Journal of Dermatology</i> , 2015, 25, 620-622.	0.6	2
5	Recurrence of IgG4-related disease following treatment with rituximab. <i>Modern Rheumatology</i> , 2013, 23, 1226-1230.	1.8	18
6	Hereditary angioedema with a de novo mutation of exon 8 in the C1 inhibitor gene showing recurrent edema of the hands around the peripheral joints: Importance for the differential diagnosis of joint swelling. <i>Arthritis and Rheumatism</i> , 2001, 44, 974-977.	6.7	9
7	Interleukin-1 β and tumor necrosis factor α synergistically stimulate prostaglandin E2-dependent production of interleukin-11 in rheumatoid synovial fibroblasts. <i>Arthritis and Rheumatism</i> , 1998, 41, 2004-2013.	6.7	57
8	Interleukin-1 β and tumor necrosis factor α synergistically stimulate prostaglandin E2-dependent production of interleukin-11 in rheumatoid synovial fibroblasts. <i>Arthritis and Rheumatism</i> , 1998, 41, 2004-2013.	6.7	4
9	Intravenous injection of tridihomo- γ -linolenoyl-glycerol into mice and its effects on delayed-type hypersensitivity. <i>Lipids</i> , 1993, 28, 873-876.	1.7	10
10	Intravenous Infusion of Tridihomo- γ -Linolenoyl-Glycerol Reduces Leukotriene B4 Production in the Rat and Rabbit. <i>Clinical Science</i> , 1993, 84, 511-516.	4.3	13
11	Effect of eicosapentaenoic acid ethyl ester on proteinuria of streptozotocin-induced diabetes mellitus in rats. <i>Lipids</i> , 1989, 24, 765-768.	1.7	5