

Lidia Marszał,

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7262822/publications.pdf>

Version: 2024-02-01

15
papers

107
citations

1478505

6
h-index

1372567

10
g-index

16
all docs

16
docs citations

16
times ranked

197
citing authors

#	ARTICLE	IF	CITATIONS
1	Fish assemblage changes relative to environmental factors and time in the Warta River, Poland, and its oxbow lakes. <i>Journal of Fish Biology</i> , 2004, 64, 483-501.	1.6	37
2	The food preferences of three-spined stickleback, <i>Gasterosteus aculeatus</i> L., downstream from a dam reservoir. <i>Oceanological and Hydrobiological Studies</i> , 2009, 38, 39-50.	0.7	12
3	Disturbed fish fauna zonation as an indicator of large-scale human impact: A case study (2011-2012) of the large, lowland Warta River, Poland. <i>Journal of Applied Ichthyology</i> , 2017, 33, 174-188.	0.7	11
4	Spatially diverse recovery (1986–2012) of fish fauna in the Warta River, Poland: The role of recolonizers' availability after large-area degradation. <i>Ecological Engineering</i> , 2017, 102, 612-624.	3.6	9
5	Habitat preferences of Ukrainian brook lamprey <i>Eudontomyzon mariae</i> ammocoetes in the lowland rivers of Central Europe. <i>Journal of Fish Biology</i> , 2016, 88, 477-491.	1.6	8
6	Recovery of fish fauna in the upper Warta River, Poland: Long-term (1963–2012) monitoring study. <i>Ecological Informatics</i> , 2016, 33, 109-118.	5.2	6
7	Intra-Seasonal and Brood-Size Dependent Variation in the Diet of Black Stork (<i>Ciconia nigra</i>) Nestlings. <i>Waterbirds</i> , 2018, 41, 268-275.	0.3	5
8	Age estimation of black stork (<i>Ciconia nigra</i>) nestlings from wing, bill, head, and tarsus lengths at the time of ringing. <i>Turkish Journal of Zoology</i> , 2018, 42, 132-136.	0.9	5
9	Brood sex ratio and nestling physiological condition as indicators of the influence of weather conditions on breeding black storks <i>Ciconia nigra</i> . <i>Ecological Indicators</i> , 2019, 104, 313-320.	6.3	5
10	Reproductive traits of the spirlin <i>Alburnoides bipunctatus</i> in the Vistula River basin. <i>Reproductive Biology</i> , 2015, 15, 184-187.	1.9	2
11	Diet of the endangered spirlin (<i>Alburnoides bipunctatus</i>) at the centre of its distribution in Europe. <i>Marine and Freshwater Research</i> , 2018, 69, 1712.	1.3	2
12	Riverine Fish Fauna in Poland. <i>Handbook of Environmental Chemistry</i> , 2020, , 195-238.	0.4	2
13	Reproductive traits of the established population of invasive western tubenose goby, <i>Proterorhinus semilunaris</i> (Actinopterygii: Perciformes: Gobiidae), in the Vistula River, Poland. <i>Acta Ichthyologica Et Piscatoria</i> , 2019, 49, 355-364.	0.7	2
14	Feeding activity of spined loach <i>Cobitis</i> sp. in Lake Lucie, Poland. <i>Folia Biologica</i> , 2003, 51 Suppl, 159-65.	0.5	1
15	Water velocity preferences of two benthic fishes, spined loach <i>Cobitis taenia</i> and Northern golden loach <i>Sabanejewia baltica</i> : an experimental approach. , 2021, 88, 1096-1102.		0