

Marianne Hope Abel

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/726886/publications.pdf>

Version: 2024-02-01

10
papers

466
citations

1306789

7
h-index

1588620

8
g-index

10
all docs

10
docs citations

10
times ranked

573
citing authors

#	ARTICLE	IF	CITATIONS
1	Suboptimal Maternal Iodine Intake Is Associated with Impaired Child Neurodevelopment at 3 Years of Age in the Norwegian Mother and Child Cohort Study. <i>Journal of Nutrition</i> , 2017, 147, 1314-1324.	1.3	136
2	Risk of Suboptimal Iodine Intake in Pregnant Norwegian Women. <i>Nutrients</i> , 2013, 5, 424-440.	1.7	78
3	Maternal Iodine Intake and Offspring Attention-Deficit/Hyperactivity Disorder: Results from a Large Prospective Cohort Study. <i>Nutrients</i> , 2017, 9, 1239.	1.7	70
4	Maternal Iodine Status is Associated with Offspring Language Skills in Infancy and Toddlerhood. <i>Nutrients</i> , 2018, 10, 1270.	1.7	58
5	Iodine Intake is Associated with Thyroid Function in Mild to Moderately Iodine Deficient Pregnant Women. <i>Thyroid</i> , 2018, 28, 1359-1371.	2.4	54
6	Insufficient maternal iodine intake is associated with subfecundity, reduced foetal growth, and adverse pregnancy outcomes in the Norwegian Mother, Father and Child Cohort Study. <i>BMC Medicine</i> , 2020, 18, 211.	2.3	38
7	Language delay and poorer school performance in children of mothers with inadequate iodine intake in pregnancy: results from follow-up at 8Âyears in the Norwegian Mother and Child Cohort Study. <i>European Journal of Nutrition</i> , 2019, 58, 3047-3058.	1.8	30
8	Iron status in mid-pregnancy and associations with interpregnancy interval, hormonal contraceptives, dietary factors and supplement use. <i>British Journal of Nutrition</i> , 2021, 126, 1270-1280.	1.2	2
9	Mild-to-moderate iodine deficiency is associated with lower birthweight and increased risk of preterm delivery in a large Norwegian pregnancy cohort. <i>Proceedings of the Nutrition Society</i> , 2020, 79, .	0.4	0
10	Inadequate iodine intake is associated with subfecundity in mild-to-moderately iodine deficient Norwegian women. <i>Proceedings of the Nutrition Society</i> , 2020, 79, .	0.4	0