

# Kris Bogaerts

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7271014/publications.pdf>

Version: 2024-02-01

46  
papers

2,892  
citations

257101

24  
h-index

243296

44  
g-index

46  
all docs

46  
docs citations

46  
times ranked

4093  
citing authors

#	ARTICLE	IF	CITATIONS
1	Bleeding risk differences after TAVR according to the ARC-HBR criteria: insights from SCOPE 2. <i>EuroIntervention</i> , 2022, 18, 503-513.	1.4	5
2	Diagnostic performance of kSORT, a blood-based mRNA assay for noninvasive detection of rejection after kidney transplantation: A retrospective multicenter cohort study. <i>American Journal of Transplantation</i> , 2021, 21, 740-750.	2.6	22
3	Two-year outcomes after percutaneous coronary intervention with drug-eluting stents or bare-metal stents in elderly patients with coronary artery disease. <i>Catheterization and Cardiovascular Interventions</i> , 2021, 97, E607-E613.	0.7	0
4	Comparison of Self-Expanding Bioprostheses for Transcatheter Aortic Valve Replacement in Patients With Symptomatic Severe Aortic Stenosis. <i>Circulation</i> , 2020, 142, 2431-2442.	1.6	96
5	The Second Strategic Reperfusion Early After Myocardial Infarction (STREAM-2) study optimizing pharmacoinvasive reperfusion strategy in older ST-elevation myocardial infarction patients. <i>American Heart Journal</i> , 2020, 226, 140-146.	1.2	13
6	Prevention of Cardiogenic Shock After Acute Myocardial Infarction. <i>Circulation</i> , 2019, 139, 137-139.	1.6	7
7	Drug-eluting stents in elderly patients with coronary artery disease (SENIOR): a randomised single-blind trial. <i>Lancet</i> , 2018, 391, 41-50.	6.3	307
8	The Belgian trial with azithromycin for acute COPD exacerbations requiring hospitalization: an investigator-initiated study protocol for a multicenter, randomized, double-blind, placebo-controlled trial. <i>International Journal of COPD</i> , 2016, 11, 687.	0.9	13
9	Drug Treatment of STEMI in the Elderly: Focus on Fibrinolytic Therapy and Insights from the STREAM Trial. <i>Drugs and Aging</i> , 2016, 33, 109-118.	1.3	5
10	Final analysis of a prospective trial on functional imaging for nodal staging in patients with prostate cancer at high risk for lymph node involvement. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015, 33, 109.e23-109.e31.	0.8	63
11	Missing Oral Health-Related Data in the interRAI-HC - Associations with Selected Variables of General Health and the Effect of Multiple Imputation on the Relationship between Oral and General Health. <i>PLoS ONE</i> , 2015, 10, e0146065.	1.1	8
12	ST-Segment-Elevation Myocardial Infarction Patients Randomized to a Pharmacoinvasive Strategy or Primary Percutaneous Coronary Intervention. <i>Circulation</i> , 2014, 130, 1139-1145.	1.6	79
13	Individualized dosimetry-based activity reduction of <sup>90</sup> Y-DOTATOC prevents severe and rapid kidney function deterioration from peptide receptor radionuclide therapy. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2014, 41, 1141-57.	3.3	25
14	Dental attendance in preschool children – a prospective study. <i>International Journal of Paediatric Dentistry</i> , 2013, 23, 84-93.	1.0	31
15	Mapping of Pelvic Lymph Node Metastases in Prostate Cancer. <i>European Urology</i> , 2013, 63, 450-458.	0.9	216
16	Contribution of four common pulmonary function tests to diagnosis of patients with respiratory symptoms: a prospective cohort study. <i>Lancet Respiratory Medicine</i> , 2013, 1, 705-713.	5.2	25
17	Fibrinolysis or Primary PCI in ST-Segment Elevation Myocardial Infarction. <i>New England Journal of Medicine</i> , 2013, 368, 1379-1387.	13.9	595
18	Risk factors for caries incidence in a cohort of Flemish preschool children. <i>Clinical Oral Investigations</i> , 2012, 16, 805-812.	1.4	19

#	ARTICLE	IF	CITATIONS
19	Statin Use and the Presence of Microalbuminuria. Results from the ERICABEL Trial: A Non-Interventional Epidemiological Cohort Study. <i>PLoS ONE</i> , 2012, 7, e31639.	1.1	11
20	Intraluminal Thrombus in Facilitated Versus Primary Percutaneous Coronary Intervention. <i>Journal of the American College of Cardiology</i> , 2011, 57, 1867-1873.	1.2	31
21	Prospective Evaluation of 11C-Choline Positron Emission Tomography/Computed Tomography and Diffusion-Weighted Magnetic Resonance Imaging for the Nodal Staging of Prostate Cancer with a High Risk of Lymph Node Metastases. <i>European Urology</i> , 2011, 60, 125-130.	0.9	142
22	Prediction of cardiogenic shock using plasma B-type natriuretic peptide and the N-terminal fragment of its pro-hormone concentrations in ST elevation myocardial infarction: An analysis from the ASSENT-4 Percutaneous Coronary Intervention Trial. <i>Critical Care Medicine</i> , 2010, 38, 1793-1801.	0.4	17
23	Middle Ear Cholesteatoma: Non-echo-planar Diffusion-weighted MR Imaging versus Delayed Gadolinium-enhanced T1-weighted MR Imaging Value in Detection. <i>Radiology</i> , 2010, 255, 866-872.	3.6	123
24	Plasma N-terminal fragment of the prohormone B-type natriuretic peptide concentrations in relation to time to treatment and Thrombolysis in Myocardial Infarction (TIMI) flow: A substudy of the Assessment of the Safety and Efficacy of a New Treatment Strategy with Percutaneous Coronary Intervention (ASSENT IV-PCI) trial. <i>American Heart Journal</i> , 2010, 159, 131-140.	1.2	19
25	Reperfusion before percutaneous coronary intervention in ST-elevation myocardial infarction patients is associated with lower N-terminal pro-brain natriuretic peptide levels during follow-up, irrespective of pre-treatment with full-dose fibrinolysis. <i>European Heart Journal</i> , 2009, 30, 2213-2219.	1.0	4
26	The Impact of Place of Enrollment and Delay to Reperfusion on 90-Day Post-Infarction Mortality in the ASSENT-4 PCI Trial. <i>JACC: Cardiovascular Interventions</i> , 2009, 2, 925-930.	1.1	17
27	Estimating local and global measures of association for bivariate interval censored data with a smooth estimate of the density. <i>Statistics in Medicine</i> , 2008, 27, 5941-5955.	0.8	10
28	Modeling the association of bivariate interval-censored data using the copula approach. <i>Statistics in Medicine</i> , 2008, 27, 6379-6392.	0.8	11
29	A New Nonparametric Approach for Baseline Covariate Adjustment for Two-Group Comparative Studies. <i>Biometrics</i> , 2008, 64, 1110-1116.	0.8	46
30	Anxiety and depression in the first six months after stroke. A longitudinal multicentre study. <i>Disability and Rehabilitation</i> , 2008, 30, 1858-1866.	0.9	100
31	Motor and Functional Recovery After Stroke. <i>Stroke</i> , 2007, 38, 2101-2107.	1.0	115
32	Age, outcomes, and treatment effects of fibrinolytic and antithrombotic combinations: Findings from Assessment of the Safety and Efficacy of a New Thrombolytic (ASSENT)-3 and ASSENT-3 PLUS. <i>American Heart Journal</i> , 2006, 152, 684.e1-684.e9.	1.2	35
33	Multivariate survival analysis for the identification of factors associated with cavity formation in permanent first molars. <i>European Journal of Oral Sciences</i> , 2005, 113, 145-152.	0.7	37
34	Use of Time by Stroke Patients. <i>Stroke</i> , 2005, 36, 1977-1983.	1.0	155
35	A New, Fast Algorithm to Find the Regions of Possible Support for Bivariate Interval-Censored Data. <i>Journal of Computational and Graphical Statistics</i> , 2004, 13, 330-340.	0.9	18
36	Efficacy of tenecteplase in combination with enoxaparin, abciximab, or unfractionated heparin: one-year follow-up results of the assessment of the safety of a new thrombolytic-3 (ASSENT-3) randomized trial in acute myocardial infarction. <i>American Heart Journal</i> , 2004, 147, 993-998.	1.2	38

#	ARTICLE	IF	CITATIONS
37	The emergence of permanent teeth in Flemish children. <i>Community Dentistry and Oral Epidemiology</i> , 2003, 31, 30-39.	0.9	65
38	Impact of caries experience in the deciduous molars on the emergence of the successors. <i>European Journal of Oral Sciences</i> , 2003, 111, 106-110.	0.7	25
39	The effect of fluorides and caries in primary teeth on permanent tooth emergence. <i>Community Dentistry and Oral Epidemiology</i> , 2003, 31, 463-470.	0.9	20
40	One-year follow-up of the ASSENT-2 trial: A double-blind, randomized comparison of single-bolus tenecteplase and front-loaded alteplase in 16,949 patients with ST-elevation acute myocardial infarction. <i>American Heart Journal</i> , 2003, 146, 27-32.	1.2	45
41	Prediction of outcome of physiotherapy in advanced Parkinson's disease. <i>Clinical Rehabilitation</i> , 2002, 16, 886-893.	1.0	25
42	On the variability of covariate adjustment. <i>Contemporary Clinical Trials</i> , 2002, 23, 127-142.	2.0	8
43	Assessing risk indicators for dental caries in the primary dentition. <i>Community Dentistry and Oral Epidemiology</i> , 2001, 29, 424-434.	0.9	86
44	Long-term fluctuations of pressure pain thresholds in healthy men, normally menstruating women and oral contraceptive users. <i>European Journal of Pain</i> , 2001, 5, 27-37.	1.4	77
45	Development of an Activity Scale for Individuals With Advanced Parkinson Disease: Reliability and "On-Off" Variability. <i>Physical Therapy</i> , 2000, 80, 1087-1096.	1.1	43
46	Survival Analysis with Interval-Censored Data. , 0, , .		40