

# Martin Srayko

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7274427/publications.pdf>

Version: 2024-02-01

32  
papers

2,121  
citations

394421

19  
h-index

434195

31  
g-index

34  
all docs

34  
docs citations

34  
times ranked

2282  
citing authors

#	ARTICLE	IF	CITATIONS
1	Sperm-specific glycogen synthase kinase 3 is required for sperm motility and the post-fertilization signal for female meiosis II in <i>Caenorhabditis elegans</i> . <i>Development (Cambridge)</i> , 2022, 149, .	2.5	2
2	Microtubule reorganization during female meiosis in <i>C. elegans</i> . <i>ELife</i> , 2021, 10, .	6.0	11
3	Kinetochores Recruitment of the Spindle and Kinetochores-Associated (Ska) Complex Is Regulated by Centrosomal PP2A in <i>Caenorhabditis elegans</i> . <i>Genetics</i> , 2019, 212, 509-522.	2.9	9
4	A Switch in Microtubule Orientation during <i>C. elegans</i> Meiosis. <i>Current Biology</i> , 2018, 28, 2991-2997.e2.	3.9	39
5	Protein phosphatase 2A is crucial for sarcomere organization in <i>Caenorhabditis elegans</i> striated muscle. <i>Molecular Biology of the Cell</i> , 2018, 29, 2084-2097.	2.1	14
6	Maternal MEMI Promotes Female Meiosis II in Response to Fertilization in <i>Caenorhabditis elegans</i> . <i>Genetics</i> , 2016, 204, 1461-1477.	2.9	6
7	G2-phase arrest prevents bristle progenitor self-renewal and synchronizes cell divisions with cell fate differentiation. <i>Development (Cambridge)</i> , 2016, 143, 1160-9.	2.5	16
8	The KLP-7 Residue S546 Is a Putative Aurora Kinase Site Required for Microtubule Regulation at the Centrosome in <i>C. elegans</i> . <i>PLoS ONE</i> , 2015, 10, e0132593.	2.5	35
9	Dual Phosphorylation of Cdk1 Coordinates Cell Proliferation with Key Developmental Processes in <i>Drosophila</i> . <i>Genetics</i> , 2014, 196, 197-210.	2.9	31
10	Measuring Microtubule Growth and Gliding in <i>Caenorhabditis elegans</i> Embryos. <i>Methods in Molecular Biology</i> , 2014, 1136, 103-116.	0.9	3
11	Suppressor mutations identify amino acids in PAA-1/PR65 that facilitate regulatory RSA-1/B $\epsilon$ 3 subunit targeting of PP2A to centrosomes in <i>C. elegans</i> . <i>Biology Open</i> , 2013, 2, 88-94.	1.2	6
12	Laulimalide Induces Dose-Dependent Modulation of Microtubule Behaviour in the <i>C. elegans</i> Embryo. <i>PLoS ONE</i> , 2013, 8, e71889.	2.5	8
13	Correlative Light and Electron Microscopy of Intermediate Stages of Meiotic Spindle Assembly in the Early <i>Caenorhabditis elegans</i> Embryo. <i>Methods in Cell Biology</i> , 2012, 111, 223-234.	1.1	16
14	The Role of $\beta$ -Tubulin in Centrosomal Microtubule Organization. <i>PLoS ONE</i> , 2012, 7, e29795.	2.5	48
15	Visualization of dynein-dependent microtubule gliding at the cell cortex: implications for spindle positioning. <i>Journal of Cell Biology</i> , 2011, 194, 377-386.	5.2	63
16	The <i>elegans</i> of spindle assembly. <i>Cellular and Molecular Life Sciences</i> , 2010, 67, 2195-2213.	5.4	51
17	Meiotic kinetochores get pushed aside by a CLS act. <i>Nature Cell Biology</i> , 2010, 12, 849-851.	10.3	3
18	Efficient chaperone-mediated tubulin biogenesis is essential for cell division and cell migration in <i>C. elegans</i> . <i>Developmental Biology</i> , 2008, 313, 320-334.	2.0	66

#	ARTICLE	IF	CITATIONS
19	Functional Interaction between Phosducin-like Protein 2 and Cytosolic Chaperonin Is Essential for Cytoskeletal Protein Function and Cell Cycle Progression. <i>Molecular Biology of the Cell</i> , 2007, 18, 2336-2345.	2.1	50
20	The <i>C. elegans</i> RSA Complex Localizes Protein Phosphatase 2A to Centrosomes and Regulates Mitotic Spindle Assembly. <i>Cell</i> , 2007, 128, 115-127.	28.9	87
21	Cortical Microtubule Contacts Position the Spindle in <i>C. elegans</i> Embryos. <i>Cell</i> , 2007, 129, 499-510.	28.9	212
22	Correlative Light and Electron Microscopy of Early <i>Caenorhabditis elegans</i> Embryos in Mitosis. <i>Methods in Cell Biology</i> , 2007, 79, 101-119.	1.1	99
23	Katanin Disrupts the Microtubule Lattice and Increases Polymer Number in <i>C. elegans</i> Meiosis. <i>Current Biology</i> , 2006, 16, 1944-1949.	3.9	152
24	The <i>C. elegans</i> Centrosome during Early Embryonic Development. , 2005, , 225-250.		0
25	An Essential Function of the <i>C. elegans</i> Ortholog of TPX2 Is to Localize Activated Aurora A Kinase to Mitotic Spindles. <i>Developmental Cell</i> , 2005, 9, 237-248.	7.0	105
26	Identification and Characterization of Factors Required for Microtubule Growth and Nucleation in the Early <i>C. elegans</i> Embryo. <i>Developmental Cell</i> , 2005, 9, 223-236.	7.0	208
27	The <i>Caenorhabditis elegans</i> Microtubule-severing Complex MEI-1/MEI-2 Katanin Interacts Differently with Two Superficially Redundant $\beta$ -Tubulin Isoforms. <i>Molecular Biology of the Cell</i> , 2004, 15, 142-150.	2.1	60
28	<i>Caenorhabditis elegans</i> TAC-1 and ZYG-9 Form a Complex that Is Essential for Long Astral and Spindle Microtubules. <i>Current Biology</i> , 2003, 13, 1506-1511.	3.9	104
29	The BTB protein MEL-26 is a substrate-specific adaptor of the CUL-3 ubiquitin-ligase. <i>Nature</i> , 2003, 425, 311-316.	27.8	378
30	Mutational analysis of <i>bli-4/kpc-4</i> reveals critical residues required for proprotein convertase function in <i>C. elegans</i> . <i>Gene</i> , 2000, 252, 15-25.	2.2	19
31	MEI-1/MEI-2 katanin-like microtubule severing activity is required for <i>Caenorhabditis elegans</i> meiosis. <i>Genes and Development</i> , 2000, 14, 1072-1084.	5.9	172
32	RNA-Mediated Interference of <i>acdc25</i> Homolog in <i>Caenorhabditis elegans</i> Results in Defects in the Embryonic Cortical Membrane, Meiosis, and Mitosis. <i>Developmental Biology</i> , 1999, 206, 15-32.	2.0	48