

Ernest Arenas

List of Publications by Citations

Source: <https://exaly.com/author-pdf/7275213/ernest-arenas-publications-by-citations.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

155
papers

12,232
citations

61
h-index

108
g-index

175
ext. papers

14,299
ext. citations

10.6
avg, IF

6.1
L-index

#	Paper	IF	Citations
155	Molecular Architecture of the Mouse Nervous System. <i>Cell</i> , 2018 , 174, 999-1014.e22	56.2	1081
154	Functional receptor for GDNF encoded by the c-ret proto-oncogene. <i>Nature</i> , 1996 , 381, 785-9	50.4	724
153	Oligodendrocyte heterogeneity in the mouse juvenile and adult central nervous system. <i>Science</i> , 2016 , 352, 1326-1329	33.3	497
152	Emerging roles of Wnts in the adult nervous system. <i>Nature Reviews Neuroscience</i> , 2010 , 11, 77-86	13.5	487
151	Molecular Diversity of Midbrain Development in Mouse, Human, and Stem Cells. <i>Cell</i> , 2016 , 167, 566-580.e19	56.2	425
150	GDNF prevents degeneration and promotes the phenotype of brain noradrenergic neurons in vivo. <i>Neuron</i> , 1995 , 15, 1465-73	13.9	317
149	Induction of a midbrain dopaminergic phenotype in Nurr1-overexpressing neural stem cells by type 1 astrocytes. <i>Nature Biotechnology</i> , 1999 , 17, 653-9	44.5	311
148	Differential regulation of midbrain dopaminergic neuron development by Wnt-1, Wnt-3a, and Wnt-5a. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 12747-52	11.5	302
147	Neuroprotection through delivery of glial cell line-derived neurotrophic factor by neural stem cells in a mouse model of Parkinson's disease. <i>Journal of Neuroscience</i> , 2001 , 21, 8108-18	6.6	253
146	Normal feeding behavior, body weight and leptin response require the neuropeptide Y Y2 receptor. <i>Nature Medicine</i> , 1999 , 5, 1188-93	50.5	240
145	Histone H2AX-dependent GABA(A) receptor regulation of stem cell proliferation. <i>Nature</i> , 2008 , 451, 460-4	50.4	218
144	Neurotrophin-3 prevents the death of adult central noradrenergic neurons in vivo. <i>Nature</i> , 1994 , 367, 368-71	50.4	198
143	A Wnt1-regulated genetic network controls the identity and fate of midbrain-dopaminergic progenitors in vivo. <i>Development (Cambridge)</i> , 2006 , 133, 89-98	6.6	196
142	Neurogenin 2 is required for the development of ventral midbrain dopaminergic neurons. <i>Development (Cambridge)</i> , 2006 , 133, 495-505	6.6	183
141	How to make a midbrain dopaminergic neuron. <i>Development (Cambridge)</i> , 2015 , 142, 1918-36	6.6	181
140	Induction of functional dopamine neurons from human astrocytes in vitro and mouse astrocytes in a Parkinson's disease model. <i>Nature Biotechnology</i> , 2017 , 35, 444-452	44.5	178
139	Brain-derived neurotrophic factor, neurotrophin-3, and neurotrophin-4/5 prevent the death of striatal projection neurons in a rodent model of Huntington's disease. <i>Journal of Neurochemistry</i> , 2000 , 75, 2190-9	6	154

138	Wnt-5a induces Dishevelled phosphorylation and dopaminergic differentiation via a CK1-dependent mechanism. <i>Journal of Cell Science</i> , 2007 , 120, 586-95	5.3	141
137	BDNF regulates reelin expression and Cajal-Retzius cell development in the cerebral cortex. <i>Neuron</i> , 1998 , 21, 305-15	13.9	135
136	Differential effects of glial cell line-derived neurotrophic factor and neurturin on developing and adult substantia nigra dopaminergic neurons. <i>Journal of Neurochemistry</i> , 1999 , 73, 70-8	6	135
135	Beta-arrestin is a necessary component of Wnt/beta-catenin signaling in vitro and in vivo. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 6690-5	11.5	129
134	Wnt5a-treated midbrain neural stem cells improve dopamine cell replacement therapy in parkinsonian mice. <i>Journal of Clinical Investigation</i> , 2008 , 118, 149-60	15.9	128
133	Communication via gap junctions underlies early functional and beneficial interactions between grafted neural stem cells and the host. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 5184-9	11.5	119
132	GSK-3beta inhibition/beta-catenin stabilization in ventral midbrain precursors increases differentiation into dopamine neurons. <i>Journal of Cell Science</i> , 2004 , 117, 5731-7	5.3	118
131	Purified Wnt-5a increases differentiation of midbrain dopaminergic cells and dishevelled phosphorylation. <i>Journal of Neurochemistry</i> , 2005 , 92, 1550-3	6	103
130	Liver X receptors and oxysterols promote ventral midbrain neurogenesis in vivo and in human embryonic stem cells. <i>Cell Stem Cell</i> , 2009 , 5, 409-19	18	101
129	Identification of midbrain floor plate radial glia-like cells as dopaminergic progenitors. <i>Glia</i> , 2008 , 56, 809-20	9	101
128	Interactions of Wnt/beta-catenin signaling and sonic hedgehog regulate the neurogenesis of ventral midbrain dopamine neurons. <i>Journal of Neuroscience</i> , 2010 , 30, 9280-91	6.6	98
127	Neural progenitor cells engineered to secrete GDNF show enhanced survival, neuronal differentiation and improve cognitive function following traumatic brain injury. <i>European Journal of Neuroscience</i> , 2006 , 23, 2119-34	3.5	98
126	Nurr1-RXR heterodimers mediate RXR ligand-induced signaling in neuronal cells. <i>Genes and Development</i> , 2003 , 17, 3036-47	12.6	96
125	Cerebrospinal fluid steroidomics: are bioactive bile acids present in brain?. <i>Journal of Biological Chemistry</i> , 2010 , 285, 4666-79	5.4	94
124	Derivation of mouse embryonic stem cells. <i>Nature Protocols</i> , 2006 , 1, 2082-7	18.8	91
123	Expression of brain-derived neurotrophic factor in cortical neurons is regulated by striatal target area. <i>Journal of Neuroscience</i> , 2001 , 21, 117-24	6.6	90
122	Brain endogenous liver X receptor ligands selectively promote midbrain neurogenesis. <i>Nature Chemical Biology</i> , 2013 , 9, 126-33	11.7	88
121	Wnt5a cooperates with canonical Wnts to generate midbrain dopaminergic neurons in vivo and in stem cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, E602-10	11.5	87

120	The extracellular domain of Lrp5/6 inhibits noncanonical Wnt signaling in vivo. <i>Molecular Biology of the Cell</i> , 2009 , 20, 924-36	3.5	87
119	Midbrain dopaminergic neurogenesis and behavioural recovery in a salamander lesion-induced regeneration model. <i>Development (Cambridge)</i> , 2007 , 134, 2881-7	6.6	87
118	Ventral midbrain glia express region-specific transcription factors and regulate dopaminergic neurogenesis through Wnt-5a secretion. <i>Molecular and Cellular Neurosciences</i> , 2006 , 31, 251-62	4.8	85
117	Functional integration of grafted neural stem cell-derived dopaminergic neurons monitored by optogenetics in an in vitro Parkinson model. <i>PLoS ONE</i> , 2011 , 6, e17560	3.7	84
116	Adenosine A1 receptor-mediated modulation of dopamine D1 receptors in stably cotransfected fibroblast cells. <i>Journal of Biological Chemistry</i> , 1998 , 273, 4718-24	5.4	84
115	Genetic identification of cell types underlying brain complex traits yields insights into the etiology of Parkinson's disease. <i>Nature Genetics</i> , 2020 , 52, 482-493	36.3	79
114	Towards stem cell replacement therapies for Parkinson's disease. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 396, 152-6	3.4	79
113	Control of neural stem cell adhesion and density by an electronic polymer surface switch. <i>Langmuir</i> , 2008 , 24, 14133-8	4	79
112	The p75 neurotrophin receptor interacts with multiple MAGE proteins. <i>Journal of Biological Chemistry</i> , 2002 , 277, 49101-4	5.4	79
111	Adenosine A2A receptors modulate the binding characteristics of dopamine D2 receptors in stably cotransfected fibroblast cells. <i>European Journal of Pharmacology</i> , 1996 , 316, 325-31	5.3	79
110	Valproic acid induces differentiation and inhibition of proliferation in neural progenitor cells via the beta-catenin-Ras-ERK-p21Cip/WAF1 pathway. <i>BMC Cell Biology</i> , 2008 , 9, 66		78
109	Parkin protects dopaminergic neurons from excessive Wnt/beta-catenin signaling. <i>Biochemical and Biophysical Research Communications</i> , 2009 , 388, 473-8	3.4	74
108	Wnt-3a utilizes a novel low dose and rapid pathway that does not require casein kinase 1-mediated phosphorylation of Dvl to activate beta-catenin. <i>Cellular Signalling</i> , 2007 , 19, 610-6	4.9	74
107	Wnt5a regulates midbrain dopaminergic axon growth and guidance. <i>PLoS ONE</i> , 2011 , 6, e18373	3.7	73
106	Wnt5a regulates ventral midbrain morphogenesis and the development of A9-A10 dopaminergic cells in vivo. <i>PLoS ONE</i> , 2008 , 3, e3517	3.7	73
105	Neuroprotection of striatal neurons against kainate excitotoxicity by neurotrophins and GDNF family members. <i>Journal of Neurochemistry</i> , 2001 , 78, 1287-96	6	73
104	Heterotrimeric G protein-dependent WNT-5A signaling to ERK1/2 mediates distinct aspects of microglia proinflammatory transformation. <i>Journal of Neuroinflammation</i> , 2012 , 9, 111	10.1	72
103	BDNF up-regulates TrkB protein and prevents the death of CA1 neurons following transient forebrain ischemia. <i>Brain Pathology</i> , 1998 , 8, 253-61	6	71

102	Wnt signaling in midbrain dopaminergic neuron development and regenerative medicine for Parkinson's disease. <i>Journal of Molecular Cell Biology</i> , 2014 , 6, 42-53	6.3	69
101	Cholestenic acids regulate motor neuron survival via liver X receptors. <i>Journal of Clinical Investigation</i> , 2014 , 124, 4829-42	15.9	69
100	Beta-arrestin and casein kinase 1/2 define distinct branches of non-canonical WNT signalling pathways. <i>EMBO Reports</i> , 2008 , 9, 1244-50	6.5	67
99	An efficient method for the derivation of mouse embryonic stem cells. <i>Stem Cells</i> , 2006 , 24, 844-9	5.8	65
98	Neural progenitors organize in small-world networks to promote cell proliferation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, E1524-32	11.5	63
97	Effects of BDNF and NT-4/5 on striatonigral neuropeptides or nigral GABA neurons in vivo. <i>European Journal of Neuroscience</i> , 1996 , 8, 1707-17	3.5	62
96	Differential expression of Wnts after spinal cord contusion injury in adult rats. <i>PLoS ONE</i> , 2011 , 6, e27009	9.7	62
95	Inhibition of mitochondrial complex III blocks neuronal differentiation and maintains embryonic stem cell pluripotency. <i>PLoS ONE</i> , 2013 , 8, e82095	3.7	61
94	NTera2: a model system to study dopaminergic differentiation of human embryonic stem cells. <i>Stem Cells and Development</i> , 2005 , 14, 517-34	4.4	59
93	Involvement of nerve growth factor and its receptor in the regulation of the cholinergic function in aged rats. <i>Journal of Neurochemistry</i> , 1991 , 57, 1483-7	6	58
92	Wnt/beta-catenin signaling blockade promotes neuronal induction and dopaminergic differentiation in embryonic stem cells. <i>Stem Cells</i> , 2009 , 27, 2917-27	5.8	57
91	Function of Wnts in dopaminergic neuron development. <i>Neurodegenerative Diseases</i> , 2006 , 3, 5-11	2.3	57
90	LifeTime and improving European healthcare through cell-based interceptive medicine. <i>Nature</i> , 2020 , 587, 377-386	50.4	56
89	Cripto as a target for improving embryonic stem cell-based therapy in Parkinson's disease. <i>Stem Cells</i> , 2005 , 23, 471-6	5.8	55
88	Wnt2 regulates progenitor proliferation in the developing ventral midbrain. <i>Journal of Biological Chemistry</i> , 2010 , 285, 7246-53	5.4	54
87	Cxcl12/Cxcr4 signaling controls the migration and process orientation of A9-A10 dopaminergic neurons. <i>Development (Cambridge)</i> , 2013 , 140, 4554-64	6.6	53
86	Wnt5a is required for endothelial differentiation of embryonic stem cells and vascularization via pathways involving both Wnt/beta-catenin and protein kinase Calpha. <i>Circulation Research</i> , 2009 , 104, 372-9	15.7	53
85	A PBX1 transcriptional network controls dopaminergic neuron development and is impaired in Parkinson's disease. <i>EMBO Journal</i> , 2016 , 35, 1963-78	13	52

84	Wnts are expressed in the spinal cord of adult mice and are differentially induced after injury. <i>Journal of Neurotrauma</i> , 2014 , 31, 565-81	5.4	52
83	Increased survival of dopaminergic neurons in striatal grafts of fetal ventral mesencephalic cells exposed to neurotrophin-3 or glial cell line-derived neurotrophic factor. <i>Cell Transplantation</i> , 2000 , 9, 45-53	4	52
82	Neuropeptide Y alters sedation through a hypothalamic Y1-mediated mechanism. <i>European Journal of Neuroscience</i> , 2001 , 13, 2241-6	3.5	46
81	Effect of opioids on acetylcholine release evoked by K ⁺ or glutamic acid from rat neostriatal slices. <i>Brain Research</i> , 1990 , 523, 51-6	3.7	46
80	SFRP1 and SFRP2 dose-dependently regulate midbrain dopamine neuron development in vivo and in embryonic stem cells. <i>Stem Cells</i> , 2012 , 30, 865-75	5.8	45
79	Vang-like protein 2 and Rac1 interact to regulate adherens junctions. <i>Journal of Cell Science</i> , 2010 , 123, 472-83	5.3	45
78	Disruption of EphA/ephrin-a signaling in the nigrostriatal system reduces dopaminergic innervation and dissociates behavioral responses to amphetamine and cocaine. <i>Molecular and Cellular Neurosciences</i> , 2004 , 26, 418-28	4.8	44
77	Dynamic temporal and cell type-specific expression of Wnt signaling components in the developing midbrain. <i>Experimental Cell Research</i> , 2006 , 312, 1626-36	4.2	43
76	WNT5A is transported via lipoprotein particles in the cerebrospinal fluid to regulate hindbrain morphogenesis. <i>Nature Communications</i> , 2019 , 10, 1498	17.4	42
75	Analysis of neural crest-derived clones reveals novel aspects of facial development. <i>Science Advances</i> , 2016 , 2, e1600060	14.3	42
74	Analysis of bioactive oxysterols in newborn mouse brain by LC/MS. <i>Journal of Lipid Research</i> , 2012 , 53, 2469-83	6.3	41
73	Crucial role of TrkB ligands in the survival and phenotypic differentiation of developing locus coeruleus noradrenergic neurons. <i>Development (Cambridge)</i> , 2003 , 130, 3535-45	6.6	40
72	Differential regulation of the expression of nerve growth factor, brain-derived neurotrophic factor, and neurotrophin-3 after excitotoxicity in a rat model of Huntington's disease. <i>Neurobiology of Disease</i> , 1998 , 5, 357-64	7.5	40
71	Distinct roles of the Y1 and Y2 receptors on neuropeptide Y-induced sensitization to sedation. <i>Journal of Neurochemistry</i> , 2001 , 78, 1201-7	6	39
70	The beta-chemokines CCL2 and CCL7 are two novel differentiation factors for midbrain dopaminergic precursors and neurons. <i>Experimental Cell Research</i> , 2008 , 314, 2123-30	4.2	37
69	Ca ²⁺ and cAMP signaling in human embryonic stem cell-derived dopamine neurons. <i>Stem Cells and Development</i> , 2010 , 19, 1355-64	4.4	34
68	Stem-cell-based strategies for the treatment of Parkinson's disease. <i>Neurodegenerative Diseases</i> , 2007 , 4, 339-47	2.3	34
67	Stromal factors SDF1 β SFRP1, and VEGFD induce dopaminergic neuron differentiation of human pluripotent stem cells. <i>Journal of Neuroscience Research</i> , 2012 , 90, 1367-81	4.4	33

66	Tiam1 regulates the Wnt/Dvl/Rac1 signaling pathway and the differentiation of midbrain dopaminergic neurons. <i>Molecular and Cellular Biology</i> , 2013 , 33, 59-70	4.8	33
65	A small synthetic cripto blocking Peptide improves neural induction, dopaminergic differentiation, and functional integration of mouse embryonic stem cells in a rat model of Parkinson's disease. <i>Stem Cells</i> , 2010 , 28, 1326-37	5.8	31
64	Microarray analyses support a role for Nurr1 in resistance to oxidative stress and neuronal differentiation in neural stem cells. <i>Stem Cells</i> , 2007 , 25, 511-9	5.8	31
63	Region-specific effects of glia on neuronal induction and differentiation with a focus on dopaminergic neurons. <i>Glia</i> , 2003 , 43, 47-51	9	31
62	Niche-derived laminin-511 promotes midbrain dopaminergic neuron survival and differentiation through YAP. <i>Science Signaling</i> , 2017 , 10,	8.8	30
61	Targeted lipidomic analysis of oxysterols in the embryonic central nervous system. <i>Molecular BioSystems</i> , 2009 , 5, 529-41		30
60	Engineering a dopaminergic phenotype in stem/precursor cells: role of Nurr1, glia-derived signals, and Wnts. <i>Annals of the New York Academy of Sciences</i> , 2005 , 1049, 51-66	6.5	30
59	Delayed dopaminergic neuron differentiation in Lrp6 mutant mice. <i>Developmental Dynamics</i> , 2010 , 239, 211-21	2.9	29
58	A proteomic analysis of LRRK2 binding partners reveals interactions with multiple signaling components of the WNT/PCP pathway. <i>Molecular Neurodegeneration</i> , 2017 , 12, 54	19	28
57	Oriented clonal cell dynamics enables accurate growth and shaping of vertebrate cartilage. <i>ELife</i> , 2017 , 6,	8.9	27
56	The antimicrobial peptide rCRAMP is present in the central nervous system of the rat. <i>Journal of Neurochemistry</i> , 2005 , 93, 1132-40	6	27
55	Nerve growth factor and basic fibroblast growth factor protect cholinergic neurons against quinolinic acid excitotoxicity in rat neostriatum. <i>European Journal of Neuroscience</i> , 1994 , 6, 706-11	3.5	27
54	Inhibition of canonical Wnt signaling promotes gliogenesis in P0-NSCs. <i>Biochemical and Biophysical Research Communications</i> , 2009 , 386, 628-33	3.4	25
53	Efficient expansion and dopaminergic differentiation of human fetal ventral midbrain neural stem cells by midbrain morphogens. <i>Neurobiology of Disease</i> , 2013 , 49, 118-27	7.5	24
52	Genetic interaction between Lrp6 and Wnt5a during mouse development. <i>Developmental Dynamics</i> , 2010 , 239, 237-45	2.9	24
51	Additional pathways of sterol metabolism: Evidence from analysis of Cyp27a1 ^{-/-} mouse brain and plasma. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2019 , 1864, 191-211	5	24
50	Dopamine Receptor Antagonists Enhance Proliferation and Neurogenesis of Midbrain Lmx1a-expressing Progenitors. <i>Scientific Reports</i> , 2016 , 6, 26448	4.9	22
49	Alpha-chemokines regulate proliferation, neurogenesis, and dopaminergic differentiation of ventral midbrain precursors and neurospheres. <i>Stem Cells</i> , 2008 , 26, 1891-900	5.8	22

48	Dickkopf 3 Promotes the Differentiation of a Rostralateral Midbrain Dopaminergic Neuronal Subset In Vivo and from Pluripotent Stem Cells In Vitro in the Mouse. <i>Journal of Neuroscience</i> , 2015 , 35, 13385-401	6.6	21
47	Foxa2: the rise and fall of dopamine neurons. <i>Cell Stem Cell</i> , 2008 , 2, 110-2	18	21
46	24(),25-Epoxycholesterol and () overexpression promote midbrain dopaminergic neurogenesis. <i>Journal of Biological Chemistry</i> , 2019 , 294, 4169-4176	5.4	20
45	Spatio-temporal expression pattern of frizzled receptors after contusive spinal cord injury in adult rats. <i>PLoS ONE</i> , 2012 , 7, e50793	3.7	20
44	BMP-2 and cAMP elevation confer locus coeruleus neurons responsiveness to multiple neurotrophic factors. <i>Journal of Neurobiology</i> , 2002 , 50, 291-304		20
43	Neurturin is a neuritogenic but not a survival factor for developing and adult central noradrenergic neurons. <i>Journal of Neurochemistry</i> , 2002 , 81, 1318-27	6	20
42	Regulation of dopamine D2 receptor affinity by cholecystokinin octapeptide in fibroblast cells cotransfected with human CCKB and D2L receptor cDNAs. <i>Molecular Brain Research</i> , 1996 , 36, 292-9		20
41	Peptide-presenting two-dimensional protein matrix on supported lipid bilayers: an efficient platform for cell adhesion. <i>Biointerphases</i> , 2007 , 2, 165-72	1.8	19
40	Wnt/ β Catenin Stimulation and Laminins Support Cardiovascular Cell Progenitor Expansion from Human Fetal Cardiac Mesenchymal Stromal Cells. <i>Stem Cell Reports</i> , 2016 , 6, 607-617	8	18
39	Dkk1 regulates ventral midbrain dopaminergic differentiation and morphogenesis. <i>PLoS ONE</i> , 2011 , 6, e15786	3.7	17
38	The Ryk receptor is expressed in glial and fibronectin-expressing cells after spinal cord injury. <i>Journal of Neurotrauma</i> , 2013 , 30, 806-17	5.4	16
37	WNT unrelated activities in commercially available preparations of recombinant WNT3a. <i>Journal of Cellular Biochemistry</i> , 2010 , 111, 1077-9	4.7	16
36	Single-cell RNA-seq analysis reveals the platinum resistance gene COX7B and the surrogate marker CD63. <i>Cancer Medicine</i> , 2018 , 7, 6193-6204	4.8	16
35	Increased Wnt levels in the neural tube impair the function of adherens junctions during neurulation. <i>Molecular and Cellular Neurosciences</i> , 2005 , 30, 437-51	4.8	15
34	GABAA and GABAB antagonists prevent the opioid inhibition of endogenous acetylcholine release evoked by glutamate from rat neostriatal slices. <i>Neuroscience Letters</i> , 1990 , 120, 201-4	3.3	15
33	Translation of WNT developmental programs into stem cell replacement strategies for the treatment of Parkinson's disease. <i>British Journal of Pharmacology</i> , 2017 , 174, 4716-4724	8.6	13
32	The tyrosine Y250 in Frizzled 4 defines a conserved motif important for structural integrity of the receptor and recruitment of Disheveled. <i>Cellular Signalling</i> , 2017 , 38, 85-96	4.9	13
31	BMPs, FGF8 and Wnts regulate the differentiation of locus coeruleus noradrenergic neuronal precursors. <i>Journal of Neurochemistry</i> , 2006 , 99, 343-52	6	13

30	Selective resistance of tachykinin-responsive cholinergic neurons in the quinolinic acid lesioned neostriatum. <i>Brain Research</i> , 1993 , 603, 317-20	3.7	13
29	Liver X receptors and cholesterol metabolism: role in ventral midbrain development and neurodegeneration. <i>F1000prime Reports</i> , 2015 , 7, 37		13
28	Laminin α controls mouse and human stem cell behaviour during midbrain dopaminergic neuron development. <i>Development (Cambridge)</i> , 2019 , 146,	6.6	11
27	Striatopallidal neurons are selectively protected by neurturin in an excitotoxic model of Huntington's disease. <i>Journal of Neurobiology</i> , 2002 , 50, 323-32		11
26	Nerve growth factor and its receptor are differentially modified by chronic naltrexone treatment during rat brain development. <i>Neuroscience Letters</i> , 1993 , 149, 47-50	3.3	11
25	Midbrain Dopaminergic Neuron Development at the Single Cell Level: and in Stem Cells. <i>Frontiers in Cell and Developmental Biology</i> , 2020 , 8, 463	5.7	10
24	A Zeb2-miR-200c loop controls midbrain dopaminergic neuron neurogenesis and migration. <i>Communications Biology</i> , 2018 , 1, 75	6.7	10
23	Fibroblast-like cells from rat plantar skin and neurotrophin-transfected 3T3 fibroblasts influence neurite growth from rat sensory neurons in vitro. <i>Journal of Neurocytology</i> , 2000 , 29, 653-63		10
22	Molecular architecture of the mouse nervous system		10
21	Genetic Identification of Cell Types Underlying Brain Complex Traits Yields Novel Insights Into the Etiology of Parkinson's Disease		9
20	Transcriptional synergy as an emergent property defining cell subpopulation identity enables population shift. <i>Nature Communications</i> , 2018 , 9, 2595	17.4	9
19	Mapping genes for calcium signaling and their associated human genetic disorders. <i>Bioinformatics</i> , 2017 , 33, 2547-2554	7.2	8
18	Transplantable midbrain dopamine neurons: a moving target. <i>Experimental Neurology</i> , 2010 , 222, 173-8	5.7	8
17	Novel isoforms of the TFIID subunit TAF4 modulate nuclear receptor-mediated transcriptional activity. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 325, 574-9	3.4	8
16	Cxcl12/Cxcr4 signaling controls the migration and process orientation of A9-A10 dopaminergic neurons. <i>Journal of Cell Science</i> , 2013 , 126, e1-e1	5.3	8
15	Mining for Oxysterols in Mouse Brain and Plasma: Relevance to Spastic Paraplegia Type 5. <i>Biomolecules</i> , 2019 , 9,	5.9	7
14	The Matricellular Protein R-Spondin 2 Promotes Midbrain Dopaminergic Neurogenesis and Differentiation. <i>Stem Cell Reports</i> , 2018 , 11, 651-664	8	7
13	WNT signaling in midbrain dopaminergic neuron development and cell replacement therapies for Parkinson's disease. <i>SpringerPlus</i> , 2015 , 4, L49		6

12	Graphene Oxide and Reduced Derivatives, as Powder or Film Scaffolds, Differentially Promote Dopaminergic Neuron Differentiation and Survival. <i>Frontiers in Neuroscience</i> , 2020 , 14, 570409	5.1	6
11	The T-type Ca Channel Ca _v 3.2 Regulates Differentiation of Neural Progenitor Cells during Cortical Development via Caspase-3. <i>Neuroscience</i> , 2019 , 402, 78-89	3.9	5
10	Srebf1 Controls Midbrain Dopaminergic Neurogenesis. <i>Cell Reports</i> , 2020 , 31, 107601	10.6	5
9	Neostriatal dopaminergic terminals prevent the GABAergic involvement in the mu- and delta-opioid inhibition of KCl-evoked endogenous acetylcholine release. <i>Brain Research</i> , 1991 , 556, 349-357	3.7	4
8	Control of tachykinin-evoked acetylcholine release from rat striatal slices by dopaminergic neurons. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1993 , 348, 445-9	3.4	3
7	MEIS-WNT5A axis regulates development of 4th ventricle choroid plexus		3
6	MEIS-WNT5A axis regulates development of fourth ventricle choroid plexus. <i>Development (Cambridge)</i> , 2021 , 148,	6.6	3
5	Combinatorial ECM Arrays Identify Cooperative Roles for Matricellular Proteins in Enhancing the Generation of TH+ Neurons From Human Pluripotent Cells.. <i>Frontiers in Cell and Developmental Biology</i> , 2021 , 9, 755406	5.7	1
4	The Cerebrospinal Fluid Profile of Cholesterol Metabolites in Parkinson's Disease and Their Association With Disease State and Clinical Features. <i>Frontiers in Aging Neuroscience</i> , 2021 , 13, 685594	5.3	0
3	Parkinson's disease in the single-cell era.. <i>Nature Neuroscience</i> , 2022 , 25, 536-538	25.5	0
2	Neurotrophins in Development of the Nervous System 1999 , 447-461		
1	Vang-like protein 2 and Rac1 interact to regulate adherens junctions. <i>Development (Cambridge)</i> , 2010 , 137, e406-e406	6.6	