

Kristian L Funck

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7282828/publications.pdf>

Version: 2024-02-01

14
papers

153
citations

1162367

8
h-index

1199166

12
g-index

15
all docs

15
docs citations

15
times ranked

317
citing authors

#	ARTICLE	IF	CITATIONS
1	Evaluation of renal oxygenation by BOLD-MRI in high-risk patients with type 2 diabetes and matched controls. <i>Nephrology Dialysis Transplantation</i> , 2023, 38, 691-699.	0.4	4
2	Coronary plaque burden in Turner syndrome a coronary computed tomography angiography study. <i>Heart and Vessels</i> , 2021, 36, 14-23.	0.5	3
3	Real-world use of cardioprotective glucose-lowering drugs in patients with type 2 diabetes and cardiovascular disease: A Danish nationwide cohort study, 2012 to 2019. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 520-529.	2.2	19
4	Arterial stiffness and progression of cerebral white matter hyperintensities in patients with type 2 diabetes and matched controls: a 5-year cohort study. <i>Diabetology and Metabolic Syndrome</i> , 2021, 13, 71.	1.2	5
5	Low physical activity is associated with impaired endothelial function in patients with type 2 diabetes and controls after 5 years of follow-up. <i>BMC Endocrine Disorders</i> , 2021, 21, 189.	0.9	6
6	Retinal vessel reactivity is not attenuated in patients with type 2 diabetes compared with matched controls and is associated with peripheral endothelial function in controls. <i>Journal of Diabetes and Its Complications</i> , 2019, 33, 641-647.	1.2	7
7	Differential vascular effects of aspirin in people with Type 2 diabetes without cardiovascular disease and matched controls without diabetes. <i>Diabetic Medicine</i> , 2019, 36, 1141-1148.	1.2	2
8	Volumes of coronary plaque disease in relation to body mass index, waist circumference, truncal fat mass and epicardial adipose tissue in patients with type 2 diabetes mellitus and controls. <i>Diabetes and Vascular Disease Research</i> , 2019, 16, 328-336.	0.9	16
9	Antiplatelet effect of aspirin during 24 h in patients with type 2 diabetes without cardiovascular disease. <i>Thrombosis Research</i> , 2018, 161, 1-6.	0.8	11
10	Detection of biomarkers using a novel proximity extension assay in patients with ST-elevation myocardial infarction. <i>Thrombosis Research</i> , 2018, 172, 21-28.	0.8	17
11	Increased high-risk coronary plaque burden is associated with arterial stiffness in patients with type 2 diabetes without clinical signs of coronary artery disease. <i>Journal of Hypertension</i> , 2017, 35, 1235-1243.	0.3	16
12	Plasma levels of MASP-1, MASP-3 and MAP44 in patients with type 2 diabetes: influence of glycaemic control, body composition and polymorphisms in the <i>MASP1</i> gene. <i>Clinical and Experimental Immunology</i> , 2017, 189, 103-112.	1.1	17
13	High burden of coronary atherosclerosis in patients with a new diagnosis of type 2 diabetes. <i>Diabetes and Vascular Disease Research</i> , 2017, 14, 468-476.	0.9	10
14	Low Physical Activity Is Associated With Increased Arterial Stiffness in Patients Recently Diagnosed With Type 2 Diabetes. <i>American Journal of Hypertension</i> , 2016, 29, 882-888.	1.0	20