

Chiara Massera

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/7290768/chiara-massera-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

118
papers

2,775
citations

31
h-index

47
g-index

125
ext. papers

2,922
ext. citations

4.1
avg, IF

4.57
L-index

#	Paper	IF	Citations
118	Combined Approach of Mechanochemistry and Electron Crystallography for the Discovery of 1D and 2D Coordination Polymers. <i>Crystal Growth and Design</i> , 2021 , 21, 6660-6664	3.5	1
117	Reusable Cavitand-Based Electrospun Membranes for the Removal of Polycyclic Aromatic Hydrocarbons from Water. <i>Small</i> , 2021 , 18, e2104946	11	0
116	Design and Synthesis of Piperazine-Based Compounds Conjugated to Humanized Ferritin as Delivery System of siRNA in Cancer Cells. <i>Bioconjugate Chemistry</i> , 2021 , 32, 1105-1116	6.3	2
115	Multidentate, V-Shaped Pyridine Building Blocks as Tectons for Crystal Engineering. <i>Chemistry - A European Journal</i> , 2021 , 27, 4660-4669	4.8	1
114	Thioureidocalix[6]arenes Pseudorotaxanes. <i>European Journal of Organic Chemistry</i> , 2021 , 2021, 5788	3.2	2
113	Mechanically-Driven Vase-Kite Conformational Switch in Cavitand Cross-Linked Polyurethanes. <i>ChemistryOpen</i> , 2020 , 9, 261-268	2.3	4
112	Synthesis of Distally-Bridged Chiral Resorcinarene Crowns. <i>European Journal of Organic Chemistry</i> , 2020 , 2020, 5695-5708	3.2	
111	Synthesis, structural and DFT interpretation of a Schiff base assisted Mn(III) derivative. <i>Journal of Molecular Structure</i> , 2020 , 1199, 126985	3.4	1
110	Solvent-responsive cavitand lanthanum complex. <i>Dalton Transactions</i> , 2019 , 48, 13732-13739	4.3	1
109	Partial functionalisation of C4-symmetric tetramethoxyresorcinarenes. <i>Supramolecular Chemistry</i> , 2019 , 31, 442-450	1.8	1
108	Structural aspects of a trimetallic CuII derivative: cytotoxicity and anti-proliferative activity on human cancer cell lines. <i>Journal of Coordination Chemistry</i> , 2019 , 72, 920-940	1.6	4
107	Spectral, electrochemical and DFT studies of a trimetallic CuII Derivative: Antimycobacterial and cytotoxic activity. <i>Inorganica Chimica Acta</i> , 2019 , 490, 155-162	2.7	4
106	New Geometries for Calix[6]arene-Based Rotaxanes. <i>European Journal of Organic Chemistry</i> , 2019 , 2019, 3513-3524	3.2	8
105	A new, deep quinoxaline-based cavitand receptor for the complexation of benzene. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2019 , 75, 103-108	0.7	
104	Crystal structure of a host-guest complex between mephedrone hydro-chloride and a tetra-phospho-nate cavitand. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2019 , 75, 277-283	0.7	
103	Structural diversity, topology and luminescent properties of a two-dimensional Cd(II) coordination polymer incorporating 4,4'-dipyridyl and 4,4'-sulfonyldibenzoic acid. <i>Journal of Molecular Structure</i> , 2019 , 1179, 618-622	3.4	4
102	A new Ni(II) coordination polymer formed by bulky bis(imidazole) and 4,4'-oxybis(benzoic acid): Topological and spectral elucidation. <i>Inorganica Chimica Acta</i> , 2018 , 469, 478-483	2.7	8

101	Sensing of halogenated aromatic hydrocarbons in water with a cavitand coated piezoelectric device. <i>Sensors and Actuators B: Chemical</i> , 2018 , 276, 340-348	8.5	8
100	Cavitand-Decorated Silicon Columnar Nanostructures for the Surface Recognition of Volatile Nitroaromatic Compounds. <i>ACS Omega</i> , 2018 , 3, 9172-9181	3.9	7
99	Metal ion complexation by tetrakisphosphonate cavitands: The influence of the ionic radius. <i>Inorganica Chimica Acta</i> , 2018 , 470, 250-253	2.7	2
98	Synthesis of Bromoundecyl Resorc[4]arenes and Applications of the Cone Stereoisomer as Selector for Liquid Chromatography. <i>Journal of Organic Chemistry</i> , 2018 , 83, 7683-7693	4.2	4
97	A zig-zag end-to-end azido bridged Mn(III) 1-D coordination polymer: Spectral elucidation, magnetism, redox study and biological activity. <i>Polyhedron</i> , 2017 , 127, 315-322	2.7	4
96	Zn and Cd-Based Coordination Networks: Highly Selective Naked Eye Sensing of Pyridine. <i>ChemistrySelect</i> , 2017 , 2, 3831-3837	1.8	1
95	Structural elucidation, EPR and photoluminescent property of a Ce(III) two dimensional coordination polymer with 4,4'-bipyridine and 2-sulfoterephthalate. <i>Journal of Molecular Structure</i> , 2017 , 1139, 390-394	3.4	6
94	In Search of the Ultimate Benzene Sensor: The EtQxBox Solution. <i>ACS Sensors</i> , 2017 , 2, 590-598	9.2	23
93	Formation of Imidazo[1,5-a]pyridine Derivatives Due to the Action of Fe on Dynamic Libraries of Imines. <i>Journal of Organic Chemistry</i> , 2017 , 82, 3820-3825	4.2	15
92	EPR interpretation, magnetism and biological study of a Cu(II) dinuclear complex assisted by a schiff base precursor. <i>Journal of Biological Inorganic Chemistry</i> , 2017 , 22, 481-495	3.7	13
91	A Deep Insight into the Photoluminescence Properties of Schiff Base Cd and Zn Complexes. <i>Inorganic Chemistry</i> , 2017 , 56, 12893-12901	5.1	26
90	Urea derivatives from carbon dioxide and amines by guanidine catalysis: Easy access to imidazolidin-2-ones under solvent-free conditions. <i>Journal of CO2 Utilization</i> , 2017 , 21, 553-561	7.6	33
89	Nitro-sonium complexation by the tetra-phospho-nate cavitand 5,11,17,23-tetra-methyl-6,10:12,16:18,22:24,4-tetra-kis-(phenyl-phospho-nato- β)-resorcin(4)arene. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2017 , 73, 1801-1805	0.7	
88	Structural elucidation, EPR and magnetic interpretation of lanthanide (Ln = La, Nd, Sm) compounds with 4,4'-bipyridine and 2-sulfoterephthalate. <i>Inorganica Chimica Acta</i> , 2016 , 453, 128-134	2.7	6
87	Synthesis and recognition properties of calix[4]arene semitubes as ditopic hosts for N-alkylpyridinium ion pairs. <i>CrystEngComm</i> , 2016 , 18, 5017-5027	3.3	5
86	Structure and properties of arylnorbornyl palladacycles as stable models for catalytic intermediates. <i>Inorganica Chimica Acta</i> , 2015 , 431, 230-233	2.7	5
85	An enolato-bridged dinuclear Cu(II) complex with a coumarin-assisted precursor: a spectral, magnetic and biological study. <i>New Journal of Chemistry</i> , 2015 , 39, 7309-7321	3.6	12
84	A Ni(II) dinuclear complex bridged by end-on azide-N and phenolate-O atoms: spectral interpretation, magnetism and biological study. <i>Inorganic Chemistry Frontiers</i> , 2015 , 2, 749-762	6.8	17

83	Chemoselective recognition with phosphonate cavitands: the ephedrine over pseudoephedrine case. <i>Chemical Communications</i> , 2015 , 51, 3426-9	5.8	6
82	A 2-D coordination polymer incorporating cobalt(II), 2-sulfoterephthalate and the flexible bridging ligand 1,3-di(4-pyridyl)propane. <i>Inorganic Chemistry Frontiers</i> , 2015 , 2, 157-163	6.8	14
81	Triply phenoxo bridged Eu(III) and Sm(III) complexes with 2,6-diformyl-4-methylphenol-di(benzoylhydrazone): structure, spectra and biological study in human cell lines. <i>New Journal of Chemistry</i> , 2015 , 39, 1101-1114	3.6	21
80	The effect of number and position of P=O/P=S bridging units on cavitand selectivity toward methyl ammonium salts. <i>Molecules</i> , 2015 , 20, 4460-72	4.8	3
79	Exploration of CH π interactions involving the π system of pseudohalide coligands in metal complexes of a Schiff-base ligand. <i>CrystEngComm</i> , 2015 , 17, 4680-4690	3.3	71
78	Discrete and polymeric complexes formed from cobalt(II), 4,4'-bipyridine and 2-sulfoterephthalate: synthetic, crystallographic and magnetic studies. <i>CrystEngComm</i> , 2015 , 17, 4502-4511	3.3	8
77	Influence of para substituents in controlling photophysical behavior and different non-covalent weak interactions in zinc complexes of a phenol based "end-off" compartmental ligand. <i>Dalton Transactions</i> , 2015 , 44, 20032-44	4.3	14
76	Design and synthesis of a cavitand pillar for MOFs. <i>Supramolecular Chemistry</i> , 2014 , 26, 151-156	1.8	2
75	Highly cytotoxic DNA-interacting copper(II) coordination compounds. <i>Metallomics</i> , 2014 , 6, 1853-68	4.5	28
74	A novel one-pot synthesis of oxazolidinones through direct introduction of CO ₂ into allylamine derivatives. <i>Tetrahedron Letters</i> , 2014 , 55, 1379-1383	2	26
73	Cavitand-grafted silicon microcantilevers as a universal probe for illicit and designer drugs in water. <i>Angewandte Chemie - International Edition</i> , 2014 , 53, 9183-8	16.4	39
72	A mixed valent heterometallic Cu(II)/Na(I) coordination polymer with sodium-phenyl bonds. <i>Dalton Transactions</i> , 2014 , 43, 5558-63	4.3	14
71	Cavitand-Grafted Silicon Microcantilevers as a Universal Probe for Illicit and Designer Drugs in Water. <i>Angewandte Chemie</i> , 2014 , 126, 9337-9342	3.6	6
70	One-Pot Multiple Metal/Organic Framework Formation: Concomitant Generation of Structural Isomers or of Drastically Distinct Materials. <i>European Journal of Inorganic Chemistry</i> , 2014 , 2014, 4385-4393	2.3	8
69	A thiourea calix[6]arene-based synthon that generates a supramolecular porous crystal structure. <i>Supramolecular Chemistry</i> , 2013 , 25, 703-708	1.8	4
68	Anion Exchange in Coordination-Network Materials. <i>European Journal of Inorganic Chemistry</i> , 2013 , 2013, 4812-4822	2.3	6
67	Analysis of the contribution of the acidity of the s-tetrazine ring in the crystal packing of coordination polymers. <i>CrystEngComm</i> , 2013 , 15, 3031	3.3	29
66	Detection of amphetamine precursors with quinoxaline-bridged cavitands. <i>Supramolecular Chemistry</i> , 2013 , 25, 682-687	1.8	8

65	Exclusive recognition of sarcosine in water and urine by a cavitand-functionalized silicon surface. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 2263-8	11.5	54
64	Switching from separated to contact ion-pair binding modes with diastereomeric calix[4]pyrrole bis-phosphonate receptors. <i>Journal of the American Chemical Society</i> , 2012 , 134, 13121-32	16.4	36
63	Supramolecular control of single-crystal-to-single-crystal transformation through selective guest exchange. <i>Chemistry - A European Journal</i> , 2011 , 17, 3064-8	4.8	35
62	Thermodynamics of host-guest interactions between methylpyridinium salts and phosphonate cavitands. <i>Supramolecular Chemistry</i> , 2010 , 22, 768-775	1.8	30
61	Formation of tetrameric water clusters driven by a cavitand template. <i>Chemical Communications</i> , 2010 , 46, 88-90	5.8	29
60	Tetraphosphonate cavitands: interplay between metal coordination and H-bonding in the formation of dimeric capsules. <i>CrystEngComm</i> , 2010 , 12, 2057	3.3	11
59	Cascade Reactions: Catalytic Synthesis of Functionalized 1,3-Dihydroisobenzofuran and Tetrahydrofuran Derivatives by Sequential Nucleophilic Ring Opening-Heterocyclization-Oxidative Carbonylation of Alkynyloxiranes. <i>Advanced Synthesis and Catalysis</i> , 2009 , 351, 2423-2432	5.6	42
58	Solid-phase synthesis of linear and cyclic peptides containing a calix[4]arene amino acid. <i>Tetrahedron Letters</i> , 2009 , 50, 3450-3453	2	7
57	Molecular recognition on a cavitand-functionalized silicon surface. <i>Journal of the American Chemical Society</i> , 2009 , 131, 7447-55	16.4	56
56	3-D lanthanide metal-organic frameworks: structure, photoluminescence, and magnetism. <i>Inorganic Chemistry</i> , 2009 , 48, 1062-8	5.1	117
55	Selective, catalytic aerobic oxidation of alcohols using CuBr(2) and bifunctional triazine-based ligands containing both a bipyridine and a TEMPO group. <i>Dalton Transactions</i> , 2009 , 3559-70	4.3	56
54	Self-assembly of heteroditopic calix[4]arene capsules through ion-pair recognition. <i>CrystEngComm</i> , 2009 , 11, 239-241	3.3	14
53	Monotopic and heteroditopic calix[4]arene receptors as hosts for pyridinium and viologen ion pairs: a solution and solid-state study. <i>Organic and Biomolecular Chemistry</i> , 2009 , 7, 3698-708	3.9	54
52	Influence of Coordinating and Non-Coordinating Anions and of a Methoxy Substituent on the Formation of Copper-Based Coordination Assemblies. <i>Crystal Growth and Design</i> , 2008 , 8, 1005-1012	3.5	33
51	Synthesis of Partially Bridged Phosphonate and Thiophosphonate Resorcinarenes. <i>Supramolecular Chemistry</i> , 2008 , 20, 29-34	1.8	14
50	Structure, cytotoxicity, and DNA-cleavage properties of the complex [Cu(II)(pbt)Br ₂]. <i>Inorganic Chemistry</i> , 2008 , 47, 3719-27	5.1	114
49	A New Sulfur-containing Schiff-Base Ligand and Binding to Copper(II) and Cobalt(II). <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , 2008 , 634, 2477-2482	1.3	3
48	Self-assembly of a double calix[6]arene pseudorotaxane in oriented channels. <i>Chemistry - A European Journal</i> , 2008 , 14, 98-106	4.8	44

47	Supramolecular sensing with phosphonate cavitands. <i>Chemistry - A European Journal</i> , 2008 , 14, 5772-9	4.8	71
46	Self-Assembled Chiral Dimeric Capsules from Difunctionalized N,C-Linked Peptidocalix[4]arenes: Scope and Limitations. <i>European Journal of Organic Chemistry</i> , 2008 , 2008, 869-886	3.2	23
45	Unique ligand-based oxidative DNA cleavage by zinc(II) complexes of hpyramol and hpyrimol. <i>Chemistry - A European Journal</i> , 2007 , 13, 5213-22	4.8	63
44	Designing nanoporous crystalline materials by self-assembly: 2D hydrogen-bonded networks from upper rim calix[4]arene diamide derivatives. <i>Inorganica Chimica Acta</i> , 2007 , 360, 970-980	2.7	9
43	Controlled copper-mediated chlorination of phenol rings under mild conditions. <i>Inorganic Chemistry</i> , 2007 , 46, 4944-50	5.1	34
42	Conformational behavior of pyrazine-bridged and mixed-bridged cavitands: a general model for solvent effects on thermal "vase-kite" switching. <i>Chemistry - A European Journal</i> , 2006 , 12, 4775-84	4.8	68
41	Coordination Dependence of Magnetic Properties within a Family of Related [Fe(II)] Complexes of a Triazine-Based Ligand. <i>European Journal of Inorganic Chemistry</i> , 2006 , 2006, 1353-1361	2.3	28
40	Cyanoresorc[5]arenes: Isolation, Conformation and Crystal Structure. <i>European Journal of Organic Chemistry</i> , 2006 , 2006, 3652-3660	3.2	7
39	Organic guests inclusion by tungsten-calix[4]arene hosts. <i>New Journal of Chemistry</i> , 2006 , 30, 952	3.6	8
38	Crystallographic evidence of theoretically novel anion π interactions. <i>New Journal of Chemistry</i> , 2006 , 30, 1561-1566	3.6	55
37	Metal-directed self-assembly of cavitand frameworks. <i>Journal of Organic Chemistry</i> , 2006 , 71, 2617-24	4.2	37
36	Crystallographic evidence of nitrate- π interactions involving the electron-deficient 1,3,5-triazine ring. <i>Inorganic Chemistry</i> , 2006 , 45, 6637-45	5.1	90
35	Crystallographic and Theoretical Evidence of Acetonitrile π Interactions with the Electron-Deficient 1,3,5-Triazine Ring. <i>Crystal Growth and Design</i> , 2006 , 6, 1569-1574	3.5	76
34	Solvent control in the synthesis of Zn(II) and Cd(II) supramolecular compounds with N,N'-(2,4-di-[(di-pyridin-2-yl)amine]-1,3,5-triazine)ethylenediamine. <i>Polyhedron</i> , 2006 , 25, 2959-2966	2.7	5
33	Intermolecular interactions between benzene and 1,3,5-triazine: a new tool for crystal engineering and molecular recognition. <i>CrystEngComm</i> , 2005 , 7, 121	3.3	21
32	Synthesis, structural characterisation and solvatochromism of a Ni(II) and a Co(II) compound with 2-aminopyrimidine as a ligand. <i>Journal of Molecular Structure</i> , 2005 , 734, 83-88	3.4	14
31	Chloro-bridged Cu(II) pairs linked into a 1D coordination polymer through a dinucleating imidazole-based ligand: 3D structure and magnetism. <i>Polyhedron</i> , 2005 , 24, 1599-1605	2.7	3
30	A one-dimensional copper (II) coordination polymer [Cu ₃ (ampym) ₂ (μ ,1-N ₃) ₄ (μ ,3-N ₃) ₂ (dmf) ₂] (ampym = 2-aminopyrimidine) containing both end-on and end-to-end azido bridges. <i>Inorganica Chimica Acta</i> , 2005 , 358, 1171-1178	2.7	28

29	N-Donor Ligand Coordination Polymers of CuII, ZnII, and CdII Obtained at Elevated Temperature and Pressure. <i>European Journal of Inorganic Chemistry</i> , 2005 , 2005, 2902-2908	2.3	21
28	An integrated approach to the study of the recognition of guests containing CH ₃ and CH ₂ acidic groups by differently rigidified cone p-tert-butylcalix[4]arene derivatives. <i>New Journal of Chemistry</i> , 2004 , 28, 56	3.6	26
27	A New Trinuclear Linear Copper(II) Complex: Unusual Crystal Structure with Semi-Coordinated Thiophene Moieties and Weak Antiferromagnetic Coupling Through the Bridging Imidazolate Rings. <i>European Journal of Inorganic Chemistry</i> , 2004 , 2004, 4566-4571	2.3	16
26	Stabilization of a Copper(II) Complex by Hydrogen Bonds between the Axial Tetrafluoroborate Anion and the Ammonium Group from the Coordinated Cationic Ligand Bis(imidazol-2-yl)methyl(isopropyl)ammonium. <i>European Journal of Inorganic Chemistry</i> , 2004 , 2004, 3025-3029	2.3	9
25	Calix[6]arene-based pseudorotaxanes: a solid state structural investigation. <i>CrystEngComm</i> , 2004 , 6, 227	3.3	24
24	An unusual open cubane structure in a $\mu(1,1)$ -azido- and alkoxo-bridged tetranuclear copper(II) complex, [Cu ₄ L ₂ ($\mu(1,1)$ -N ₃) ₂].5H ₂ O (H ₃ L = N,NP(2-Hydroxypropyl-1,3-diyl)bis-salicylideneimine). <i>Inorganic Chemistry</i> , 2004 , 43, 6842-7	5.1	125
23	Synthesis of 4H-3,1-benzoxazines, quinazolin-2-ones, and quinoline-4-ones by palladium-catalyzed oxidative carbonylation of 2-ethynylaniline derivatives. <i>Journal of Organic Chemistry</i> , 2004 , 69, 2469-77	4.2	99
22	Design and Preparation of Mesogenic Cavitands. <i>Collection of Czechoslovak Chemical Communications</i> , 2004 , 69, 1362-1380		4
21	Synthesis of New Calix[4]arene-Based Ionophores. <i>Collection of Czechoslovak Chemical Communications</i> , 2004 , 69, 1309-1324		2
20	Reactivity of Ph ₂ (2-C ₅ H ₄ N)PSe towards Ru ₃ (CO) ₁₂ and mononuclear MCl ₂ (PhCN) ₂ (M=Pd or Pt) complexes. <i>Inorganica Chimica Acta</i> , 2003 , 350, 421-427	2.7	7
19	Reactions between Cu(II) halide complexes of 2,2'-dipyridylamine and nitrophenolates. X-ray diffraction structure of the [bis(2,2'-dipyridylamine)di(p-nitrophenolato)di(η -methoxy)dicopper(II)], [Cu(dipyam)(pnp)(η -OCH ₃) ₂] ₂ , a dimeric complex containing axial phenoxo ligands in square-pyramidal arrangements. <i>Inorganica Chimica Acta</i> , 2003 , 349, 128-134	2.7	6
18	Energy Partitioning Analysis of the Bonding in L ₂ TM η ² H ₂ and L ₂ TM η ² H ₄ (TM = Ni, Pd, Pt; L ₂ = (PH ₃) ₂ , (PMe ₃) ₂ , H ₂ PCH ₂ PH ₂ , H ₂ P(CH ₂) ₂ PH ₂) <i>Organometallics</i> , 2003 , 22, 2758-2765	3.8	82
17	Rational design of cavitand receptors for mass sensors. <i>Journal of the American Chemical Society</i> , 2003 , 125, 12068-9	16.4	47
16	Reactivity of Ph ₂ P(Se)CH ₂ C ₆ H ₄ CH ₂ P(Se)Ph ₂ with [M ₃ (CO) ₁₂] (M=Ru or Fe). Synthesis and characterization of the new carbene cluster [Ru ₃ (μ ₃ -Se)(μ ₃ -H)(μ ₂ -PPh ₂)(CO) ₆ (μ ₂ -CHC ₆ H ₄ CH ₂ PPh ₂)]. <i>Inorganic Chemistry</i> , 2003 , 42, 8509-18	5.1	17
15	A novel self-assembled supramolecular architecture involving cation, anion and a calix[4]arene heteroditopic receptor. <i>Tetrahedron Letters</i> , 2002 , 43, 7311-7314	2	31
14	Reaction of [Pt{Fe(CO) ₃ (NO)} ₂ (PhCN) ₂] with diphenyl(2-pyridyl)phosphine selenide. Crystal structure of [(CO) ₃ Fe(η -Se){Pt(CO)P(2-C ₅ H ₄ N)Ph ₂ } ₂] and its theoretical study. <i>Inorganica Chimica Acta</i> , 2002 , 330, 95-102	2.7	15
13	[Cu(phdpa)Cl] ⁺ (phdpa=bis(2-pyridylmethyl)aniline): a moiety of unusual stability in some 1:1 Cu(II) complexes of phdpa. Synthesis and X-ray crystal structures of [Cu(phdpa)Cl ₂] and [Cu ₂ (phdpa) ₂ Cl ₃]PF ₆ ·0.5MeOH. <i>Inorganica Chimica Acta</i> , 2002 , 340, 97-104	2.7	25
12	Design and self-assembly of wide and robust coordination cages. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 4911-5	11.5	85

11	Reactivity of the heterometallic clusters $[\text{HMCo}_3(\text{CO})_{12}]$ and $[\text{Et}_4\text{N}][\text{MCo}_3(\text{CO})_{12}]$ ($\text{M} = \text{Fe}, \text{Ru}$) toward phosphine selenides, including selenium transfer. Crystal structures of $[\text{HRuCo}_3(\text{CO})_7(\mu\text{-CO})_3(\mu\text{-dppy})]$, $[\text{MCo}_2(\mu_3\text{-Se})(\text{CO})_7(\mu\text{-dppy})]$, and $[\text{RuCo}_2(\mu_3\text{-Se})(\text{CO})_7(\mu\text{-dppm})]$ [$\text{dppy} = \text{Ph}_2(2\text{-C}_5\text{H}_4\text{N})\text{P}$, $\text{dppm} = \text{Ph}_2\text{PCH}_2\text{PPh}_2$]. <i>Inorganic Chemistry</i> , 2001 , 40, 1718-1723	5.1	34
10	Interaction between benzene and model neutral organic molecules bearing acid CH groups. <i>New Journal of Chemistry</i> , 2002 , 26, 1718-1723	3.6	34
9	Iodo Derivatives of Bimetallic FePd and FePt Complexes. Crystal Structures of. <i>Monatshefte für Chemie</i> , 2001 , 132, 885-896	1.4	9
8	Multiple Oxidative Addition of $\text{Ph}_2(\text{C}_5\text{H}_4\text{N})\text{PSe}$ to $[\text{Ru}_3(\text{CO})_{12}]$: Structural Characterization of $[\text{Ru}_3(\mu_3\text{-Se})(\mu\text{-PPh}_2)_2(\mu\text{-C}_5\text{H}_4\text{N})(\mu_3\text{-C}_5\text{H}_4\text{N})(\text{CO})_6]$ Containing Two Differently Metalated 2-Pyridyl Fragments. <i>European Journal of Inorganic Chemistry</i> , 2001 , 2001, 721-723	2.3	19
7	Synthesis and crystal structure of some (2,2'-dipyridylamine)(carboxylato)(phenoxo)copper(II) complexes with rather unusual phenoxo ligands. <i>Inorganica Chimica Acta</i> , 2001 , 324, 162-172	2.7	6
6	Easy Access to Phosphido-Selenido Clusters. Reaction of $[\text{Ru}_3(\text{CO})_{12}]$ with $\text{Ph}_2(\text{pyth})\text{PSe}$ { $\text{pyth} = 5\text{-}(2\text{-Pyridyl})\text{-}2\text{-Thienyl}$ } and Crystal Structure of $[\text{Ru}_3(\mu_3\text{-Se})(\mu\text{-PPh}_2)(\mu\text{-pyth})(\text{CO})_6\{\text{P}(\text{pyth})\text{Ph}_2\}]$. <i>Journal of Cluster Science</i> , 2001 , 12, 259-271	3	17
5	Copper(II)nitrito complexes with 2,2'-dipyridylamine. Crystal structures of the [(acetato)(2,2'-dipyridylamine)(nitrito-O,O')copper(II)] and [(2,2'-dipyridylamine)(nitrito-O,O')(nitrito-O)copper(II)] \cdot 2D(acetonitrile). <i>Inorganica Chimica Acta</i> , 2000 , 309, 1-9	2.7	21
4	Multiple oxidative addition of diphenyl-2-thienylphosphine selenide to $[\text{Ru}_3(\text{CO})_{12}]$: crystal and molecular structure of $[\text{Ru}_3(\mu_3\text{-Se})(\mu\text{-PPh}_2)(\mu_3\text{-C}_4\text{H}_3\text{S})(\text{CO})_6\{\text{P}(\text{C}_4\text{H}_3\text{S})\text{Ph}_2\}]$. <i>Inorganica Chimica Acta</i> , 2000 , 300-302, 471-476	2.7	20
3	Chelating versus bridging behaviour and NMR fluxionality of dppf in the nido clusters $[\text{M}_3\text{Se}_2(\text{CO})_7(\text{dppf})]$ [$\text{M} = \text{Fe}$ or Ru , $\text{dppf} = \text{Fe}(\mu\text{-C}_5\text{H}_4\text{PPh}_2)_2$]. Crystal structure of the chelated ruthenium derivative. <i>Journal of the Chemical Society Dalton Transactions</i> , 1999 , 3515-3521		34
2	Cluster growth reactions with selenido-carbonyl clusters. Synthesis and structural characterization of $[\text{M}_2\text{Ru}_2(\mu_4\text{-Se})_2(\mu\text{-CO})_4(\text{CO})_6(\text{PPh}_3)_2]$ ($\text{M} = \text{Mo}$ or W). <i>Journal of the Chemical Society Dalton Transactions</i> , 1998 , 321-322		13
1	Synthesis of quinoxaline cavitand baskets. <i>Supramolecular Chemistry</i> , 1-10	1.8	0