

# Cahit Birdir

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7296477/publications.pdf>

Version: 2024-02-01

30  
papers

984  
citations

759233

12  
h-index

477307

29  
g-index

34  
all docs

34  
docs citations

34  
times ranked

1127  
citing authors

#	ARTICLE	IF	CITATIONS
1	A review of image processing methods for fetal head and brain analysis in ultrasound images. <i>Computer Methods and Programs in Biomedicine</i> , 2022, 215, 106629.	4.7	25
2	Placenta accreta spectrum disordersâ€™ experience of management in a German tertiary perinatal centre. <i>Archives of Gynecology and Obstetrics</i> , 2021, 303, 1451-1460.	1.7	11
3	Evaluation of carcinoembryonic antigenâ€™related cell adhesion molecule 1 blood serum levels in women at high risk for preeclampsia. <i>American Journal of Reproductive Immunology</i> , 2021, 85, e13375.	1.2	1
4	Are two children at once better than one? Risk analysis of twin pregnancies and births after assisted reproduction. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2021, 264, 76-82.	1.1	3
5	MR-proANP, a cardiovascular biomarker to predict late-onset preeclampsia and intrauterine growth restricted fetuses. <i>Pregnancy Hypertension</i> , 2020, 22, 54-58.	1.4	0
6	Prenatal Prediction of Outcome by Fetal Gastroschisis in a Tertiary Referral Center. <i>Diagnostics</i> , 2020, 10, 540.	2.6	8
7	Predictive value of sFlt-1, PlGF, sFlt-1/PlGF ratio and PAPP-A for late-onset preeclampsia and IUGR between 32 and 37â€™weeks of pregnancy. <i>Pregnancy Hypertension</i> , 2018, 12, 124-128.	1.4	54
8	Afamin: an early predictor of preeclampsia. <i>Archives of Gynecology and Obstetrics</i> , 2018, 298, 1009-1016.	1.7	18
9	Soluble B7â€™H4 blood serum levels are elevated in women at high risk for preeclampsia in the first trimester, as well as in patients with confirmed preeclampsia. <i>American Journal of Reproductive Immunology</i> , 2018, 80, e12988.	1.2	11
10	Impact of maternal serum levels of Visfatin, AFP, PAPP-A, sFlt-1 and PlGF at 11â€™13 weeks gestation on small for gestational age births. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2017, 30, 629-634.	1.5	12
11	Follistatin during pregnancy and its potential role as an ovarian suppressing agent. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2017, 212, 150-154.	1.1	8
12	Preeclampsia in Fetal Medicine. <i>Ultraschall in Der Medizin</i> , 2017, 38, 589-590.	1.5	0
13	Serum concentrations of soluble B7â€™H4 in early pregnancy are elevated in women with preterm premature rupture of fetal membranes. <i>American Journal of Reproductive Immunology</i> , 2016, 76, 149-154.	1.2	11
14	Detection of circulating trophoblast particles in maternal blood using density gradient centrifugation in preeclampsia and in normotensive pregnancies. <i>Hypertension in Pregnancy</i> , 2016, 35, 323-329.	1.1	6
15	Maternal serum copeptin, MR-proANP and procalcitonin levels at 11â€™13 weeks gestation in the prediction of preeclampsia. <i>Archives of Gynecology and Obstetrics</i> , 2015, 292, 1033-1042.	1.7	21
16	Change of anti-Mullerian-hormone levels during follicular phase in PCOS patients. <i>Gynecological Endocrinology</i> , 2015, 31, 26-30.	1.7	6
17	Maternal serum anti-Müllerian hormone at 11â€™13 weeksâ€™ gestation in the prediction of preeclampsia. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2015, 28, 865-868.	1.5	15
18	Idiopathic polyhydramnios and fetal gender. <i>Archives of Gynecology and Obstetrics</i> , 2015, 291, 987-991.	1.7	9

#	ARTICLE	IF	CITATIONS
19	Reply to "Questions about and speculations on the incidence of idiopathic polyhydramnios by fetal gender". Archives of Gynecology and Obstetrics, 2015, 291, 1197-1197.	1.7	0
20	Embryo Reduction in Dichorionic Triplets to Dichorionic Twins by Intrafetal Laser. Fetal Diagnosis and Therapy, 2014, 35, 83-86.	1.4	17
21	Anti-Mullerian-hormone levels during pregnancy and postpartum. Reproductive Biology and Endocrinology, 2013, 11, 60.	3.3	67
22	Maternal serum tumour necrosis factor receptor 1 (TNF-R1) at 30-33 weeks in the prediction of preeclampsia. Journal of Maternal-Fetal and Neonatal Medicine, 2013, 26, 763-767.	1.5	3
23	Noninvasive Prenatal Testing for Fetal Trisomies in a Routinely Screened First-Trimester Population. Obstetrical and Gynecological Survey, 2013, 68, 173-175.	0.4	13
24	First-Trimester Screening for Neural Tube Defects Using Alpha-Fetoprotein. Fetal Diagnosis and Therapy, 2012, 31, 109-114.	1.4	38
25	Early Detection of Maternal Risk for Preeclampsia. ISRN Obstetrics & Gynecology, 2012, 2012, 1-7.	1.2	20
26	Noninvasive prenatal testing for fetal trisomies in a routinely screened first-trimester population. American Journal of Obstetrics and Gynecology, 2012, 207, 374.e1-374.e6.	1.3	323
27	Chromosome-selective sequencing of maternal plasma cell-free DNA for first-trimester detection of trisomy 21 and trisomy 18. American Journal of Obstetrics and Gynecology, 2012, 206, 322.e1-322.e5.	1.3	245
28	First-Trimester Screening for Trisomy 21 with Adjustment for Biochemical Results of Previous Pregnancies. Fetal Diagnosis and Therapy, 2011, 30, 194-202.	1.4	8
29	Placental Growth Factor: A Predictive Marker for Preeclampsia?. Gynakologisch-geburtshilfliche Rundschau, 2009, 49, 94-99.	0.9	19
30	Altered angiogenesis in preeclampsia: evaluation of a new test system for measuring placental growth factor. Clinical Chemistry and Laboratory Medicine, 2007, 45, 1504-10.	2.3	9