

# Anna Wasilewska

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7303813/publications.pdf>

Version: 2024-02-01

74  
papers

941  
citations

471061

17  
h-index

500791

28  
g-index

75  
all docs

75  
docs citations

75  
times ranked

1427  
citing authors

#	ARTICLE	IF	CITATIONS
1	Treatment and long-term outcome in primary nephrogenic diabetes insipidus. <i>Nephrology Dialysis Transplantation</i> , 2023, 38, 2120-2130.	0.4	9
2	Untargeted Metabolomics Analysis of the Serum Metabolic Signature of Childhood Obesity. <i>Nutrients</i> , 2022, 14, 214.	1.7	14
3	An open-label phase 2 trial to assess the efficacy, safety and pharmacokinetics of lanthanum carbonate in hyperphosphatemic children and adolescents with chronic kidney disease undergoing dialysis. <i>BMC Nephrology</i> , 2022, 23, 84.	0.8	0
4	Health-related quality of life in children with immunoglobulin A nephropathy – results of a multicentre national study. <i>Archives of Medical Science</i> , 2021, 17, 84-91.	0.4	4
5	Urine NGAL and KIM-1 – Tubular Injury Biomarkers in Long-Term Survivors of Childhood Solid Tumors: A Cross-Sectional Study. <i>Journal of Clinical Medicine</i> , 2021, 10, 399.	1.0	4
6	Tumor Necrosis Factor-Like Weak Inducer of Apoptosis and Selected Cytokines – Potential Biomarkers in Children with Solitary Functioning Kidney. <i>Journal of Clinical Medicine</i> , 2021, 10, 497.	1.0	3
7	Is Urinary Netrin-1 a Good Marker of Tubular Damage in Preterm Newborns?. <i>Journal of Clinical Medicine</i> , 2021, 10, 847.	1.0	0
8	Galectin-3 – A New Player of Kidney Damage or an Innocent Bystander in Children with a Single Kidney?. <i>Journal of Clinical Medicine</i> , 2021, 10, 2012.	1.0	1
9	Urinary excretion of renin and angiotensinogen in hypertensive children and adolescents. <i>Archives of Medical Science</i> , 2021, 17, 1325-1331.	0.4	0
10	The Role of Complement Component C3 Activation in the Clinical Presentation and Prognosis of IgA Nephropathy – A National Study in Children. <i>Journal of Clinical Medicine</i> , 2021, 10, 4405.	1.0	5
11	Efficacy and safety of valsartan in children aged 1–5 years with hypertension, with or without chronic kidney disease: a randomized, double-blind study followed by open-label phase. <i>Current Medical Research and Opinion</i> , 2021, 37, 1-23.	0.9	3
12	Urinary Levels of Cathepsin B in Preterm Newborns. <i>Journal of Clinical Medicine</i> , 2021, 10, 4254.	1.0	0
13	Urinary netrin-1 concentration in healthy full-term newborns. <i>Archives of Medical Science</i> , 2021, 17, 47-52.	0.4	0
14	Urinary Beta-2-Microglobulin and Late Nephrotoxicity in Childhood Cancer Survivors. <i>Journal of Clinical Medicine</i> , 2021, 10, 5279.	1.0	3
15	Clinical and Epidemiological Analysis of Children’s Urinary Tract Infections in Accordance with Antibiotic Resistance Patterns of Pathogens. <i>Journal of Clinical Medicine</i> , 2021, 10, 5260.	1.0	3
16	Analysis of Indications for Voiding Cystography in Children. <i>Journal of Clinical Medicine</i> , 2021, 10, 5809.	1.0	2
17	Are low birth weight children predisposed to renal loss of carnitine?. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2020, 33, 2612-2617.	0.7	0
18	P182125 YEARS OF GROWTH HORMONE TREATMENT IN CHILDREN WITH CHRONIC KIDNEY DISEASE IN POLAND - RESULTS OF NATIONAL MULTICENTER STUDY. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.4	0

#	ARTICLE	IF	CITATIONS
19	Urine NGAL and KIM-1: tubular injury markers in acute lymphoblastic leukemia survivors. <i>Cancer Chemotherapy and Pharmacology</i> , 2020, 86, 741-749.	1.1	7
20	Salivary Gland Dysfunction, Protein Glycooxidation and Nitrosative Stress in Children with Chronic Kidney Disease. <i>Journal of Clinical Medicine</i> , 2020, 9, 1285.	1.0	28
21	A Case-Control Study of Salivary Redox Homeostasis in Hypertensive Children. Can Salivary Uric Acid be a Marker of Hypertension?. <i>Journal of Clinical Medicine</i> , 2020, 9, 837.	1.0	40
22	Compound heterozygous IFT140 variants in two Polish families with Sensenbrenner syndrome and early onset end-stage renal disease. <i>Orphanet Journal of Rare Diseases</i> , 2020, 15, 36.	1.2	12
23	Dysfunction of Salivary Glands, Disturbances in Salivary Antioxidants and Increased Oxidative Damage in Saliva of Overweight and Obese Adolescents. <i>Journal of Clinical Medicine</i> , 2020, 9, 548.	1.0	34
24	The tubular damage markers: neutrophil gelatinase-associated lipocalin and kidney injury molecule-1 in newborns with exposure to maternal diabetes during pregnancy. <i>Archives of Medical Science</i> , 2020, , .	0.4	0
25	Salivary FRAP as A Marker of Chronic Kidney Disease Progression in Children. <i>Antioxidants</i> , 2019, 8, 409.	2.2	34
26	Urinary procollagen III aminoterminal propeptide and $\beta$ -catenin – New diagnostic biomarkers in solitary functioning kidney?. <i>Advances in Medical Sciences</i> , 2019, 64, 189-194.	0.9	3
27	GP221 – Influence of controlled physical activity on serum adipokines concentration in obese children. , 2019, , .		0
28	Twenty years of growth hormone treatment in dialyzed children in Poland – Results of national multicenter study. <i>Advances in Medical Sciences</i> , 2019, 64, 90-99.	0.9	0
29	The role of appetite-regulating hormones: Ghrelin and leptin in the nutritional status of children with neurogenic bladder due to myelomeningocele. <i>Journal of Paediatrics and Child Health</i> , 2019, 55, 928-931.	0.4	0
30	The Tubular Damage Markers: Neutrophil Gelatinase-Associated Lipocalin and Kidney Injury Molecule-1 in Newborns with Intrauterine Growth Restriction. <i>Neonatology</i> , 2019, 115, 169-174.	0.9	12
31	Activity of lysosomal exoglycosidases in the urine of healthy normotensive and pre-hypertensive children. <i>Advances in Medical Sciences</i> , 2019, 64, 24-31.	0.9	0
32	Upper metastable limit osmolality of urine as a predictor of kidney stone formation in children. <i>Urolithiasis</i> , 2019, 47, 155-163.	1.2	14
33	Tubular and Glomerular Biomarkers of Acute Kidney Injury in Newborns. <i>Current Drug Metabolism</i> , 2019, 20, 332-349.	0.7	10
34	Response to Letter to the Editor re – Urinary tract infection in children: Diagnosis, treatment, imaging – Comparison of current guidelines™. <i>Journal of Pediatric Urology</i> , 2018, 14, 301-302.	0.6	0
35	Urinary exoglycosidases, reference values in healthy children. <i>Advances in Medical Sciences</i> , 2018, 63, 224-229.	0.9	3
36	Pediatric reference data on activity of urinary N-acetyl- $\beta$ -D-hexosaminidase and its isoenzymes. <i>Advances in Medical Sciences</i> , 2018, 63, 94-99.	0.9	4

#	ARTICLE	IF	CITATIONS
37	FP778GDIGA1 AND GDIGA1/C3 SERUM RATIO IN CHILDREN WITH IGA NEPHROPATHY AND HENOCH-SCHÄ¶NLEIN NEPHRITIS. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i307-i307.	0.4	0
38	Serum Renalase Levels in Adolescents with Primary Hypertension. <i>Pediatric Cardiology</i> , 2018, 39, 1258-1264.	0.6	11
39	Serum GDIGA1 levels in children with IgA nephropathy and Henoch-Sch¶nlein nephritis. <i>Central-European Journal of Immunology</i> , 2018, 43, 162-167.	0.4	16
40	Salivary Biomarkers of Oxidative Stress in Children with Chronic Kidney Disease. <i>Journal of Clinical Medicine</i> , 2018, 7, 209.	1.0	63
41	Citrate usage in the leading causes of blindness: new possibilities for the old metabolite. <i>Metabolomics</i> , 2018, 14, 82.	1.4	1
42	Correlation of Salusin Beta with hs-CRP and ADMA in Hypertensive Children and Adolescents. <i>Current Pharmaceutical Design</i> , 2018, 24, 3551-3557.	0.9	6
43	The Possible Impact of Hyperuricemia on Serum Soluble Receptor for Advanced Glycation end Products (sRAGE) Levels in Teenagers: A Case Control Study. <i>Current Pharmaceutical Design</i> , 2018, 24, 3232-3239.	0.9	1
44	Determining normal values of urinary phosphorus excretion in 3913 healthy children aged 2¶18 to aid early diagnosis and treatment for urolithiasis. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2017, 106, 1170-1175.	0.7	9
45	Long-term renal outcome in children with OCRL mutations: retrospective analysis of a large international cohort. <i>Nephrology Dialysis Transplantation</i> , 2016, 33, gfw350.	0.4	27
46	Is urine intercellular adhesion molecule-1 a marker of renal disorder in children with ureteropelvic junction obstruction?. <i>Biomarkers</i> , 2016, 21, 123-128.	0.9	2
47	Health term¶born girls had higher levels of urine neutrophil gelatinase¶associated lipocalin than boys during the first postnatal days. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2016, 105, 1105-1108.	0.7	10
48	Salusins in Hypertension and Related Cardiovascular Diseases. <i>Current Drug Metabolism</i> , 2016, 17, 827-833.	0.7	7
49	Urine exoglycosidases are potential markers of renal tubular injury in children with ureteropelvic junction obstruction. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2015, 104, e518-23.	0.7	7
50	An association between kidney stone composition and urinary metabolic disturbances in children. <i>Journal of Pediatric Urology</i> , 2014, 10, 130-135.	0.6	38
51	Disease-related social situation in family of children with chronic kidney disease ¶ parents¶ assessment. A multicentre study. <i>Annals of Agricultural and Environmental Medicine</i> , 2014, 21, 876-881.	0.5	6
52	The assessment of thiol status in children with neurogenic bladder caused by meningomyelocele. <i>Urology Journal</i> , 2014, 11, 1400-5.	0.3	1
53	Asymmetric and Symmetric Dimethylarginine in Adolescents with Hyperuricemia. <i>Disease Markers</i> , 2013, 35, 407-412.	0.6	11
54	Urinary MMP-9/NGAL Ratio as a Potential Marker of FSGS in Nephrotic Children. <i>Disease Markers</i> , 2013, 34, 357-362.	0.6	20

#	ARTICLE	IF	CITATIONS
55	Is plasma symmetric dimethylarginine a suitable marker of renal function in children and adolescents?. <i>Scandinavian Journal of Urology and Nephrology</i> , 2012, 46, 58-64.	1.4	7
56	Urinary cytokine profiles in unilateral congenital hydronephrosis. <i>Pediatric Nephrology</i> , 2012, 27, 2107-2113.	0.9	36
57	Markers of systemic inflammation in children with hyperuricemia. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2012, 101, 497-500.	0.7	29
58	KIM-1 and NGAL: new markers of obstructive nephropathy. <i>Pediatric Nephrology</i> , 2011, 26, 579-586.	0.9	119
59	Urinary monocyte chemoattractant protein-1 excretion in children with glomerular proteinuria. <i>Scandinavian Journal of Urology and Nephrology</i> , 2011, 45, 52-59.	1.4	17
60	Neutrophil gelatinase-associated lipocalin (NGAL): a new marker of cyclosporine nephrotoxicity?. <i>Pediatric Nephrology</i> , 2010, 25, 889-897.	0.9	63
61	High-sensitivity C-reactive protein and mean platelet volume in paediatric hypertension. <i>Pediatric Nephrology</i> , 2010, 25, 1519-1527.	0.9	10
62	Serum RANKL, osteoprotegerin (OPG), and RANKL/OPG ratio in nephrotic children. <i>Pediatric Nephrology</i> , 2010, 25, 2067-2075.	0.9	25
63	Urinary transforming growth factor beta1 in children and adolescents with congenital solitary kidney. <i>Pediatric Nephrology</i> , 2009, 24, 753-759.	0.9	13
64	IgA Nephropathy in a Girl with Psoriasis and Seronegative Arthritis. <i>Pediatric Dermatology</i> , 2008, 25, 408-409.	0.5	7
65	MDR-1 gene polymorphisms and clinical course of steroid-responsive nephrotic syndrome in children. <i>Pediatric Nephrology</i> , 2007, 22, 44-51.	0.9	47
66	High-sensitivity C-reactive protein (hs-CRP) level in children with nephrotic syndrome. <i>Pediatric Nephrology</i> , 2007, 22, 403-408.	0.9	8
67	Expression of multidrug resistance P-glycoprotein on lymphocytes from nephrotic children treated with cyclosporine A and ACE-inhibitor. <i>European Journal of Pediatrics</i> , 2007, 166, 447-452.	1.3	14
68	Assessment of serum cystatin C in children with congenital solitary kidney. <i>Pediatric Nephrology</i> , 2006, 21, 688-693.	0.9	25
69	Expression of P-glycoprotein in lymphocytes from children with nephrotic syndrome, depending on their steroid response. <i>Pediatric Nephrology</i> , 2006, 21, 1274-1280.	0.9	18
70	Vascular endothelial growth factor in children with nephrotic syndrome treated with cyclosporine A. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2006, 95, 291-296.	0.7	3
71	Vascular endothelial growth factor in children with nephrotic syndrome treated with cyclosporine A. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2006, 95, 291-296.	0.7	0
72	Glucocorticoid receptor and vascular endothelial growth factor in nephrotic syndrome. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2006, 95, 587-593.	0.7	0

#	ARTICLE	IF	CITATIONS
73	Glucocorticoid receptor and vascular endothelial growth factor in nephrotic syndrome. Acta Paediatrica, International Journal of Paediatrics, 2006, 95, 587-593.	0.7	7
74	Expression of Glucocorticoid Receptors in Nephrotic Children Depending on Total Prednisone Dose. Journal of Pediatric Endocrinology and Metabolism, 2005, 18, 799-806.	0.4	3