
Kazuki Terada

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/7303875/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Inhibition of Nerve Growth Factor-Induced Neurite Outgrowth from PC12 Cells by Dexamethasone: Signaling Pathways through the Glucocorticoid Receptor and Phosphorylated Akt and ERK1/2. PLoS ONE, 2014, 9, e93223.	2.5	35
2	Cholinesterase inhibitor rivastigmine enhances nerve growth factor-induced neurite outgrowth in PC12 cells via sigma-1 and sigma-2 receptors. PLoS ONE, 2018, 13, e0209250.	2.5	23
3	Fluvoxamine moderates reduced voluntary activity following chronic dexamethasone infusion in mice via recovery of BDNF signal cascades. Neurochemistry International, 2014, 69, 9-13.	3.8	13
4	Sigma-2 receptor as a potential therapeutic target for treating central nervous system disorders. Neural Regeneration Research, 2019, 14, 1893.	3.0	13
5	Prodrugs for Skin Delivery of Menahydroquinone-4, an Active Form of Vitamin K2(20), Could Overcome the Photoinstability and Phototoxicity of Vitamin K2(20). International Journal of Molecular Sciences, 2019, 20, 2548.	4.1	10
6	Atypical Antipsychotic Drug Ziprasidone Protects against Rotenone-Induced Neurotoxicity: An In Vitro Study. Molecules, 2020, 25, 4206.	3.8	9
7	Oxytocin treatment improves dexamethasoneâ€induced depressionâ€like symptoms associated with enhancement ofÂhippocampal <scp>CREBâ€BDNF</scp> signaling in female mice. Neuropsychopharmacology Reports, 2022, 42, 356-361.	2.3	9
8	The Kampo medicine Yokukansan (YKS) enhances nerve growth factor (NGF)-induced neurite outgrowth in PC12 cells. Bosnian Journal of Basic Medical Sciences, 2018, 18, 224-233.	1.0	8
9	Antitumor Effects and Delivery Profiles of Menahydroquinone-4 Prodrugs with Ionic or Nonionic Promoiety to Hepatocellular Carcinoma Cells. Molecules, 2018, 23, 1738.	3.8	6