

# Daming Zhang

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7314624/publications.pdf>

Version: 2024-02-01

38  
papers

1,201  
citations

361045

20  
h-index

377514

34  
g-index

38  
all docs

38  
docs citations

38  
times ranked

1988  
citing authors

#	ARTICLE	IF	CITATIONS
1	Novel long noncoding RNA OTUD6B-AS1 indicates poor prognosis and inhibits clear cell renal cell carcinoma proliferation via the Wnt/ $\beta$ -catenin signaling pathway. <i>Molecular Cancer</i> , 2019, 18, 15.	7.9	107
2	The DNA methylation-regulated miR-193a-3p dictates the multi-chemoresistance of bladder cancer via repression of SRSF2/PLAU/HIC2 expression. <i>Cell Death and Disease</i> , 2014, 5, e1402-e1402.	2.7	89
3	miR-193a-3p regulates the multi-drug resistance of bladder cancer by targeting the LOXL4 gene and the Oxidative Stress pathway. <i>Molecular Cancer</i> , 2014, 13, 234.	7.9	68
4	Recommendations for Surgery During the Novel Coronavirus (COVID-19) Epidemic. <i>Indian Journal of Surgery</i> , 2020, 82, 124-128.	0.2	67
5	PERK silencing inhibits glioma cell growth under low glucose stress by blockage of p-AKT and subsequent HK2's mitochondria translocation. <i>Scientific Reports</i> , 2015, 5, 9065.	1.6	65
6	MicroRNA-153 is tumor suppressive in glioblastoma stem cells. <i>Molecular Biology Reports</i> , 2013, 40, 2789-2798.	1.0	56
7	The miR-193a-3p-regulated ING5 gene activates the DNA damage response pathway and inhibits multi-chemoresistance in bladder cancer. <i>Oncotarget</i> , 2015, 6, 10195-10206.	0.8	56
8	miR-577 inhibits glioblastoma tumor growth via the Wnt signaling pathway. <i>Molecular Carcinogenesis</i> , 2016, 55, 575-585.	1.3	53
9	MiR-196a exerts its oncogenic effect in glioblastoma multiforme by inhibition of $\beta$ -catenin both in vitro and in vivo. <i>Neuro-Oncology</i> , 2014, 16, 652-661.	0.6	52
10	MiR-193a-3p promotes the multi-chemoresistance of bladder cancer by targeting the HOXC9 gene. <i>Cancer Letters</i> , 2015, 357, 105-113.	3.2	50
11	The miR-193a-3p regulated PSEN1 gene suppresses the multi-chemoresistance of bladder cancer. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2015, 1852, 520-528.	1.8	45
12	Nitazoxanide, an antiprotozoal drug, inhibits late-stage autophagy and promotes ING1-induced cell cycle arrest in glioblastoma. <i>Cell Death and Disease</i> , 2018, 9, 1032.	2.7	45
13	Methionine and cystine double deprivation stress suppresses glioma proliferation via inducing ROS/autophagy. <i>Toxicology Letters</i> , 2015, 232, 349-355.	0.4	41
14	The effect of Hsa_circ_0001451 in clear cell renal cell carcinoma cells and its relationship with clinicopathological features. <i>Journal of Cancer</i> , 2018, 9, 3269-3277.	1.2	36
15	MiR-106a is an independent prognostic marker in patients with glioblastoma. <i>Neuro-Oncology</i> , 2013, 15, 707-717.	0.6	32
16	mir-300 Promotes Self-Renewal and Inhibits the Differentiation of Glioma Stem-Like Cells. <i>Journal of Molecular Neuroscience</i> , 2014, 53, 637-644.	1.1	31
17	Exosomal miR-2276-5p in Plasma Is a Potential Diagnostic and Prognostic Biomarker in Glioma. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 671202.	1.8	27
18	MicroRNA-195 protection against focal cerebral ischemia by targeting CX3CR1. <i>Journal of Neurosurgery</i> , 2019, 131, 1445-1454.	0.9	25

#	ARTICLE	IF	CITATIONS
19	Circulating MicroRNAs as Potential Noninvasive Biomarkers of Spontaneous Intracerebral Hemorrhage. <i>World Neurosurgery</i> , 2020, 133, e369-e375.	0.7	25
20	Postoperative pneumonia after craniotomy: incidence, risk factors and prediction with a nomogram. <i>Journal of Hospital Infection</i> , 2020, 105, 167-175.	1.4	22
21	Long non-coding RNA urothelial carcinoma-associated 1 as a tumor biomarker for the diagnosis of urinary bladder cancer. <i>Tumor Biology</i> , 2017, 39, 101042831770999.	0.8	21
22	Plasma D-dimer predicts poor outcome and mortality after spontaneous intracerebral hemorrhage. <i>Brain and Behavior</i> , 2021, 11, 462-468.	1.0	21
23	BMP8A promotes survival and drug resistance via Nrf2/TRIM24 signaling pathway in clear cell renal cell carcinoma. <i>Cancer Science</i> , 2020, 111, 1555-1566.	1.7	20
24	The downregulated long noncoding RNA <i>DHRS4-AS1</i> is protumoral and associated with the prognosis of clear cell renal cell carcinoma. <i>OncoTargets and Therapy</i> , 2018, Volume 11, 5631-5646.	1.0	19
25	Prognostic and clinicopathological role of long non-coding RNA UCA1 in various carcinomas. <i>Oncotarget</i> , 2017, 8, 28373-28384.	0.8	19
26	Preclinical optimization of a broad-spectrum anti-bladder cancer tri-drug regimen via the Feedback System Control (FSC) platform. <i>Scientific Reports</i> , 2015, 5, 11464.	1.6	17
27	$\alpha$ -1,2-Mannosidase MAN1C1 Inhibits Proliferation and Invasion of Renal Clear Cell Carcinoma. <i>Journal of Cancer</i> , 2018, 9, 4618-4626.	1.2	16
28	Moxidectin inhibits glioma cell viability by inducing G0/G1 $\frac{1}{2}$ cell cycle arrest and apoptosis. <i>Oncology Reports</i> , 2018, 40, 1348-1358.	1.2	15
29	Prognostic and clinicopathological role of long non-coding RNA taurine upregulated 1 in various human malignancies: A systemic review and meta-analysis. <i>Tumor Biology</i> , 2017, 39, 101042831771436.	0.8	13
30	Inhibition of MicroRNA-381 Promotes Tumor Cell Growth and Chemoresistance in Clear-Cell Renal Cell Carcinoma. <i>Medical Science Monitor</i> , 2019, 25, 5181-5190.	0.5	10
31	Association between the Epidermal Growth Factor +61G/A Polymorphism and Glioma Risk: A Meta-Analysis. <i>PLoS ONE</i> , 2014, 9, e95139.	1.1	9
32	Acetylcholine plays an antinociceptive role by modulating pain-induced discharges of pain-related neurons in the caudate putamen of rats. <i>NeuroReport</i> , 2014, 25, 164-170.	0.6	8
33	Functions of the bone morphogenetic protein signaling pathway through non-coding RNAs. <i>Non-coding RNA Research</i> , 2022, 7, 178-183.	2.4	8
34	Left ventricular ejection fraction as an independent predictor of poor outcome in acute intracerebral hemorrhage. <i>Brain and Behavior</i> , 2020, 10, e01643.	1.0	5
35	Total intracranial volume as a covariate for predicting prognosis in patients with primary intracerebral hemorrhage. <i>Clinical Neurology and Neurosurgery</i> , 2022, 214, 107135.	0.6	3
36	Irregular shape as an independent predictor of prognosis in patients with primary intracerebral hemorrhage. <i>Scientific Reports</i> , 2022, 12, .	1.6	3

#	ARTICLE	IF	CITATIONS
37	Multiple Meningiomas Characterized by Benign and Malignant Tumor Entities. <i>CNS Neuroscience and Therapeutics</i> , 2013, 19, 984-986.	1.9	1
38	Roles of Loss of Chromosome 14q Allele in the Prognosis of Renal Cell Carcinoma with C-reactive Protein Abnormality. <i>Chinese Medical Journal</i> , 2017, 130, 2176-2182.	0.9	1