

Daming Zhang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7314624/publications.pdf>

Version: 2024-02-01

38
papers

1,201
citations

361045

20
h-index

377514

34
g-index

38
all docs

38
docs citations

38
times ranked

1988
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 1 | Total intracranial volume as a covariate for predicting prognosis in patients with primary intracerebral hemorrhage. <i>Clinical Neurology and Neurosurgery</i> , 2022, 214, 107135. | 0.6 | 3 |
| 2 | Irregular shape as an independent predictor of prognosis in patients with primary intracerebral hemorrhage. <i>Scientific Reports</i> , 2022, 12, . | 1.6 | 3 |
| 3 | Functions of the bone morphogenetic protein signaling pathway through non-coding RNAs. <i>Non-coding RNA Research</i> , 2022, 7, 178-183. | 2.4 | 8 |
| 4 | Plasma D-dimer predicts poor outcome and mortality after spontaneous intracerebral hemorrhage. <i>Brain and Behavior</i> , 2021, 11, 462-468. | 1.0 | 21 |
| 5 | Exosomal miR-2276-5p in Plasma Is a Potential Diagnostic and Prognostic Biomarker in Glioma. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 671202. | 1.8 | 27 |
| 6 | Circulating MicroRNAs as Potential Noninvasive Biomarkers of Spontaneous Intracerebral Hemorrhage. <i>World Neurosurgery</i> , 2020, 133, e369-e375. | 0.7 | 25 |
| 7 | Left ventricular ejection fraction as an independent predictor of poor outcome in acute intracerebral hemorrhage. <i>Brain and Behavior</i> , 2020, 10, e01643. | 1.0 | 5 |
| 8 | Postoperative pneumonia after craniotomy: incidence, risk factors and prediction with a nomogram. <i>Journal of Hospital Infection</i> , 2020, 105, 167-175. | 1.4 | 22 |
| 9 | BMP8A promotes survival and drug resistance via Nrf2/TRIM24 signaling pathway in clear cell renal cell carcinoma. <i>Cancer Science</i> , 2020, 111, 1555-1566. | 1.7 | 20 |
| 10 | Recommendations for Surgery During the Novel Coronavirus (COVID-19) Epidemic. <i>Indian Journal of Surgery</i> , 2020, 82, 124-128. | 0.2 | 67 |
| 11 | Novel long noncoding RNA OTUD6B-AS1 indicates poor prognosis and inhibits clear cell renal cell carcinoma proliferation via the Wnt/ β -catenin signaling pathway. <i>Molecular Cancer</i> , 2019, 18, 15. | 7.9 | 107 |
| 12 | MicroRNA-195 protection against focal cerebral ischemia by targeting CX3CR1. <i>Journal of Neurosurgery</i> , 2019, 131, 1445-1454. | 0.9 | 25 |
| 13 | Inhibition of MicroRNA-381 Promotes Tumor Cell Growth and Chemoresistance in Clear-Cell Renal Cell Carcinoma. <i>Medical Science Monitor</i> , 2019, 25, 5181-5190. | 0.5 | 10 |
| 14 | α -1,2-Mannosidase MAN1C1 Inhibits Proliferation and Invasion of Renal Clear Cell Carcinoma. <i>Journal of Cancer</i> , 2018, 9, 4618-4626. | 1.2 | 16 |
| 15 | The downregulated long noncoding RNA α -DHS4-AS1 is protumoral and associated with the prognosis of clear cell renal cell carcinoma. <i>OncoTargets and Therapy</i> , 2018, Volume 11, 5631-5646. | 1.0 | 19 |
| 16 | The effect of Hsa_circ_0001451 in clear cell renal cell carcinoma cells and its relationship with clinicopathological features. <i>Journal of Cancer</i> , 2018, 9, 3269-3277. | 1.2 | 36 |
| 17 | Nitazoxanide, an antiprotozoal drug, inhibits late-stage autophagy and promotes ING1-induced cell cycle arrest in glioblastoma. <i>Cell Death and Disease</i> , 2018, 9, 1032. | 2.7 | 45 |
| 18 | Moxidectin inhibits glioma cell viability by inducing G0/G1 $\frac{1}{2}$ cell cycle arrest and apoptosis. <i>Oncology Reports</i> , 2018, 40, 1348-1358. | 1.2 | 15 |

| # | ARTICLE | IF | CITATIONS |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 19 | Prognostic and clinicopathological role of long non-coding RNA taurine upregulated 1 in various human malignancies: A systemic review and meta-analysis. <i>Tumor Biology</i> , 2017, 39, 101042831771436. | 0.8 | 13 |
| 20 | Long non-coding RNA urothelial carcinoma-associated 1 as a tumor biomarker for the diagnosis of urinary bladder cancer. <i>Tumor Biology</i> , 2017, 39, 101042831770999. | 0.8 | 21 |
| 21 | Roles of Loss of Chromosome 14q Allele in the Prognosis of Renal Cell Carcinoma with C-reactive Protein Abnormality. <i>Chinese Medical Journal</i> , 2017, 130, 2176-2182. | 0.9 | 1 |
| 22 | Prognostic and clinicopathological role of long non-coding RNA UCA1 in various carcinomas. <i>Oncotarget</i> , 2017, 8, 28373-28384. | 0.8 | 19 |
| 23 | miR-577 inhibits glioblastoma tumor growth via the Wnt signaling pathway. <i>Molecular Carcinogenesis</i> , 2016, 55, 575-585. | 1.3 | 53 |
| 24 | Preclinical optimization of a broad-spectrum anti-bladder cancer tri-drug regimen via the Feedback System Control (FSC) platform. <i>Scientific Reports</i> , 2015, 5, 11464. | 1.6 | 17 |
| 25 | PERK silence inhibits glioma cell growth under low glucose stress by blockage of p-AKT and subsequent HK2's mitochondria translocation. <i>Scientific Reports</i> , 2015, 5, 9065. | 1.6 | 65 |
| 26 | MiR-193a-3p promotes the multi-chemoresistance of bladder cancer by targeting the HOXC9 gene. <i>Cancer Letters</i> , 2015, 357, 105-113. | 3.2 | 50 |
| 27 | The miR-193a-3p regulated PSEN1 gene suppresses the multi-chemoresistance of bladder cancer. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2015, 1852, 520-528. | 1.8 | 45 |
| 28 | Methionine and cystine double deprivation stress suppresses glioma proliferation via inducing ROS/autophagy. <i>Toxicology Letters</i> , 2015, 232, 349-355. | 0.4 | 41 |
| 29 | The miR-193a-3p-regulated ING5 gene activates the DNA damage response pathway and inhibits multi-chemoresistance in bladder cancer. <i>Oncotarget</i> , 2015, 6, 10195-10206. | 0.8 | 56 |
| 30 | Association between the Epidermal Growth Factor +61G/A Polymorphism and Glioma Risk: A Meta-Analysis. <i>PLoS ONE</i> , 2014, 9, e95139. | 1.1 | 9 |
| 31 | The DNA methylation-regulated miR-193a-3p dictates the multi-chemoresistance of bladder cancer via repression of SRSF2/PLAU/HIC2 expression. <i>Cell Death and Disease</i> , 2014, 5, e1402-e1402. | 2.7 | 89 |
| 32 | miR-193a-3p regulates the multi-drug resistance of bladder cancer by targeting the LOXL4 gene and the Oxidative Stress pathway. <i>Molecular Cancer</i> , 2014, 13, 234. | 7.9 | 68 |
| 33 | MiR-196a exerts its oncogenic effect in glioblastoma multiforme by inhibition of β both in vitro and in vivo. <i>Neuro-Oncology</i> , 2014, 16, 652-661. | 0.6 | 52 |
| 34 | mir-300 Promotes Self-Renewal and Inhibits the Differentiation of Glioma Stem-Like Cells. <i>Journal of Molecular Neuroscience</i> , 2014, 53, 637-644. | 1.1 | 31 |
| 35 | Acetylcholine plays an antinociceptive role by modulating pain-induced discharges of pain-related neurons in the caudate putamen of rats. <i>NeuroReport</i> , 2014, 25, 164-170. | 0.6 | 8 |
| 36 | MiR-106a is an independent prognostic marker in patients with glioblastoma. <i>Neuro-Oncology</i> , 2013, 15, 707-717. | 0.6 | 32 |

| # | ARTICLE | IF | CITATIONS |
|----|------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 37 | MicroRNA-153 is tumor suppressive in glioblastoma stem cells. <i>Molecular Biology Reports</i> , 2013, 40, 2789-2798. | 1.0 | 56 |
| 38 | Multiple Meningiomas Characterized by Benign and Malignant Tumor Entities. <i>CNS Neuroscience and Therapeutics</i> , 2013, 19, 984-986. | 1.9 | 1 |