

Heng-Chen Yao

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/7320196/publications.pdf>

Version: 2024-02-01

22
papers

238
citations

1051969

10
h-index

1113639

15
g-index

22
all docs

22
docs citations

22
times ranked

386
citing authors

#	ARTICLE	IF	CITATIONS
1	Gender-Related Difference in D-Dimer Level Predicts In-Hospital Heart Failure after Primary PCI for ST-Segment Elevation Myocardial Infarction. <i>Disease Markers</i> , 2021, 2021, 1-8.	0.6	5
2	Leukocyte telomere length: a potential biomarker for the prognosis of coronary artery disease. <i>Biomarkers in Medicine</i> , 2020, 14, 933-941.	0.6	2
3	Prognostic efficacy of high-sensitivity C-reactive protein to albumin ratio in patients with acute coronary syndrome. <i>Biomarkers in Medicine</i> , 2019, 13, 811-820.	0.6	39
4	Plasma miR-208b and miR-499: Potential Biomarkers for Severity of Coronary Artery Disease. <i>Disease Markers</i> , 2019, 2019, 1-7.	0.6	13
5	HMGB1 Protects the Heart Against Ischemia-Induced Reperfusion Injury via PI3K/Akt Pathway-Mediated Upregulation of VEGF Expression. <i>Frontiers in Physiology</i> , 2019, 10, 1595.	1.3	20
6	HGF Reduces Disease Severity and Inflammation by Attenuating the NF- κ B Signaling in a Rat Model of Pulmonary Artery Hypertension. <i>Inflammation</i> , 2018, 41, 924-931.	1.7	25
7	Dynamic evaluation of depression and anxiety may be beneficial in the prediction of long term mortality in patients with coronary artery disease. <i>International Journal of Cardiology</i> , 2018, 258, 30.	0.8	0
8	Statins therapy might be beneficial in the treatment of myocardial ischemia reperfusion injury. <i>International Journal of Cardiology</i> , 2017, 239, 15.	0.8	1
9	Autophagy in myocardial ischemia reperfusion injury: Friend or foe?. <i>International Journal of Cardiology</i> , 2017, 239, 10.	0.8	10
10	Inhibition of IL-17 might be a novel therapeutic target in the treatment of myocardial ischemia reperfusion injury. <i>International Journal of Cardiology</i> , 2017, 239, 29.	0.8	2
11	Association between statins use and erectile dysfunction. <i>International Journal of Cardiology</i> , 2017, 239, 16.	0.8	0
12	High mobility group box 1 protein attenuates myocardial ischemia reperfusion injury via inhibition of the p38 mitogen-activated protein kinase signaling pathway. <i>Experimental and Therapeutic Medicine</i> , 2017, 14, 1582-1588.	0.8	18
13	Red cell distribution width is a predictor of mortality in patients with chronic heart failure. <i>International Journal of Cardiology</i> , 2016, 212, 79-81.	0.8	5
14	Serum albumin levels might be an adverse predictor of long term mortality in patients with acute myocardial infarction. <i>International Journal of Cardiology</i> , 2016, 223, 647-648.	0.8	7
15	Intravenous high mobility group box 1 upregulates the expression of HIF-1 α in the myocardium via a protein kinase B-dependent pathway in rats following acute myocardial ischemia. <i>Molecular Medicine Reports</i> , 2016, 13, 1211-1219.	1.1	14
16	Simvastatin protects the heart against ischemia reperfusion injury via inhibiting HMGB1 expression through PI3K/Akt signal pathways. <i>International Journal of Cardiology</i> , 2015, 201, 568-569.	0.8	14
17	Postconditioning with simvastatin decreases myocardial injury in rats following acute myocardial ischemia. <i>Experimental and Therapeutic Medicine</i> , 2015, 9, 1166-1170.	0.8	11
18	Response to letter regarding "Increased serum HMGB1 level may predict the fatal outcomes in patients with chronic heart failure". <i>International Journal of Cardiology</i> , 2015, 187, 434-435.	0.8	1

#	ARTICLE	IF	CITATIONS
19	Protection of intravenous HMGB1 on myocardial ischemia reperfusion injury. <i>International Journal of Cardiology</i> , 2015, 184, 280-282.	0.8	8
20	Effect of Basic Fibroblast Growth Factor on the Myocardial Expression of Hypoxia-inducible Factor-1 α and Vascular Endothelial Growth Factor Following Acute Myocardial Infarction. <i>Heart Lung and Circulation</i> , 2013, 22, 946-951.	0.2	15
21	Correlation between serum high-mobility group box-1 levels and high-sensitivity C-reactive protein and troponin I in patients with coronary artery disease. <i>Experimental and Therapeutic Medicine</i> , 2013, 6, 121-124.	0.8	24
22	Interleukin-2 enhances the cytotoxic activity of circulating natural killer cells in patients with chronic heart failure. <i>Heart and Vessels</i> , 2009, 24, 283-286.	0.5	4